

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

THIS PAGE BLANK (USPTO)

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07D 261/00		A2	(11) International Publication Number: WO 98/28282 (43) International Publication Date: 2 July 1998 (02.07.98)
(21) International Application Number: PCT/US97/23470 (22) International Filing Date: 18 December 1997 (18.12.97) (30) Priority Data: 08/771,814 23 December 1996 (23.12.96) US 08/879,763 20 June 1997 (20.06.97) US (71) Applicant: THE DU PONT MERCK PHARMACEUTICAL COMPANY [US/US]; 1007 Market Street, Wilmington, DE 19898 (US). (72) Inventors: PRUITT, James, Russell; 237 Skycrest Drive, Landenberg, PA 19350 (US). FEVIG, John, Matthew; 987 Church Road, Lincoln University, PA 19352 (US). QUAN, Mimi, Lifen; 113 Venus Drive, Newark, DE 19711 (US). PINTO, Donald, Joseph, Phillip; 39 Whitson Road, Newark, DE 19702 (US). (74) Agent: VANCE, David, H.; The Du Pont Merck Pharmaceutical Company, Legal Patent Records Center, 1007 Market Street, Wilmington, DE 19898 (US).			(81) Designated States: AM, AU, AZ, BR, BY, CA, CN, CZ, EE, HU, IL, JP, KG, KR, KZ, LT, LV, MD, MX, NO, NZ, PL, RO, RU, SG, SI, SK, TJ, TM, UA, VN, Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: OXYGEN OR SULFUR CONTAINING HETEROAROMATICS AS FACTOR Xa INHIBITORS			
(57) Abstract <p>The present application describes oxygen and sulfur containing heteroaromatics and derivatives thereof of formula (I), or pharmaceutically acceptable salt or prodrug forms thereof, wherein J is O or S and D may be C(=NH)NH₂, which are useful as inhibitors of factor Xa.</p>			
<div style="text-align: center;"><p style="text-align: right;">(I)</p></div>			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

TITLE

Oxygen or Sulfur Containing Heteroaromatics as Factor Xa
Inhibitors


5

FIELD OF THE INVENTION

This invention relates generally to oxygen or sulfur
containing 5-membered ring heteroaromatics which are
inhibitors of trypsin-like serine protease enzymes, especially
10 factor Xa, pharmaceutical compositions containing the same,
and methods of using the same as anticoagulant agents for
treatment and prevention of thromboembolic disorders.

BACKGROUND OF THE INVENTION


15 WO 95/13155 and PCT International Application US 96/07692
describe isoxazoline and isoxazole fibrinogen receptor
antagonists of the formula:



20

wherein R¹ may be a basic group, U-V may be a six-membered
aromatic ring, W-X may be a variety of linear or cyclic
groups, and Y is an oxy group. Thus, these compounds all
contain an acid functionality (i.e., W-X-C(=O)-Y). In
25 contrast, the presently claimed compounds do not contain such
an acid functionality.

EP 0,513,387 depicts active oxygen inhibitors which are
oxazoles or thiazoles of the formula:




30


wherein X is O or S, R² is preferably hydrogen, and both R¹ and
R³ are substituted cyclic groups, with at least one being

WO 95/18111 addresses fibrinogen receptor antagonists, containing basic and acidic termini, of the formula:


5



In U.S. Patent No. 5,463,071, Himmelsbach et al depict cell aggregation inhibitors which are 5-membered heterocycles of the formula:



25 Baker et al, in U.S. Patent No. 5,317,103, discuss 5-HT₁
agonists which are indole substituted five-membered
heteroaromatic compounds of the formula:



30

wherein R¹ may be pyrrolidine or piperidine and A may be a basic group including amino and amidino. Baker et al,

however, do not indicate that A can be a substituted ring system like that contained in the presently claimed heteroaromatics.

5 Tidwell et al, in *J. Med. Chem.* **1978**, 21(7), 613-623, describe a series of diarylamidine derivatives including 3,5-bis(4-amidinophenyl)isoxazole. This series of compounds was tested against thrombin, trypsin, and pancreatic kallikrein. The presently claimed invention does not include these types of compounds.

10 Activated factor Xa, whose major practical role is the generation of thrombin by the limited proteolysis of prothrombin, holds a central position that links the intrinsic and extrinsic activation mechanisms in the final common pathway of blood coagulation. The generation of thrombin, the
15 final serine protease in the pathway to generate a fibrin clot, from its precursor is amplified by formation of prothrombinase complex (factor Xa, factor V, Ca^{2+} and phospholipid). Since it is calculated that one molecule of factor Xa can generate 138 molecules of thrombin (Elodi, S.,
20 Varadi, K.: *Optimization of conditions for the catalytic effect of the factor IXa-factor VIII Complex: Probable role of the complex in the amplification of blood coagulation. Thromb. Res.* **1979**, 15, 617-629), inhibition of factor Xa may be more efficient than inactivation of thrombin in
25 interrupting the blood coagulation system.

30 Therefore, efficacious and specific inhibitors of factor Xa are needed as potentially valuable therapeutic agents for the treatment of thromboembolic disorders. It is thus desirable to discover new factor Xa inhibitors.

SUMMARY OF THE INVENTION


Accordingly, one object of the present invention is to provide novel oxygen or sulfur containing aromatic heterocycles which are useful as factor Xa inhibitors or
35 pharmaceutically acceptable salts or prodrugs thereof.

It is another object of the present invention to provide pharmaceutical compositions comprising a pharmaceutically acceptable carrier and a therapeutically effective amount of

at least one of the compounds of the present invention or a pharmaceutically acceptable salt or prodrug form thereof.

It is another object of the present invention to provide a method for treating thromboembolic disorders comprising administering to a host in need of such treatment a therapeutically effective amount of at least one of the compounds of the present invention or a pharmaceutically acceptable salt or prodrug form thereof.

These and other objects, which will become apparent during the following detailed description, have been achieved by the inventors' discovery that compounds of formula (I):




I

or pharmaceutically acceptable salt or prodrug forms thereof, wherein A, B, D, E, G, J, M, R^{1a}, R^{1b}, s and Z are defined below, are effective factor Xa inhibitors.

DETAILED DESCRIPTION OF PREFERRED EMBODIMENTS

[1] Thus, in a first embodiment, the present invention provides novel compounds of formula I:



I

or a stereoisomer or pharmaceutically acceptable salt thereof, wherein;

ring M contains, in addition to J, 0-2 N atoms;

J is O or S;

D is selected from CN, C(=NR⁸)NR⁷R⁹, NHC(=NR⁸)NR⁷R⁹, NR⁸CH(=NR⁷), C(O)NR⁷R⁸, and (CR⁸R⁹)_tNR⁷R⁸, provided that D is substituted meta or para to G on E;

5 E is selected from phenyl, pyridyl, pyrimidyl, pyrazinyl, pyridazinyl, and piperidinyl substituted with 1 R;

alternatively, D-E-G together represent pyridyl substituted with 1 R;

10

R is selected from H, halogen, (CH₂)_tOR³, C₁₋₄ alkyl, OCF₃, and CF₃;

G is absent or is selected from NHCH₂, OCH₂, and SCH₂;

15

Z is selected from a C₁₋₄ alkylene, (CH₂)_rO(CH₂)_r, (CH₂)_rNR³(CH₂)_r, (CH₂)_rC(O)(CH₂)_r, (CH₂)_rC(O)O(CH₂)_r, (CH₂)_rOC(O)(CH₂)_r, (CH₂)_rC(O)NR³(CH₂)_r, (CH₂)_rNR³C(O)(CH₂)_r, (CH₂)_rOC(O)O(CH₂)_r, (CH₂)_rOC(O)NR³(CH₂)_r, (CH₂)_rNR³C(O)O(CH₂)_r, (CH₂)_rNR³C(O)NR³(CH₂)_r, (CH₂)_rS(O)_p(CH₂)_r, (CH₂)_rSO₂NR³(CH₂)_r, (CH₂)_rNR³SO₂(CH₂)_r, and (CH₂)_rNR³SO₂NR³(CH₂)_r, provided that Z does not form a N-N, N-O, N-S, NCH₂N, NCH₂O, or NCH₂S bond with ring M or group A;

25

R^{1a} and R^{1b} are independently absent or selected from -(CH₂)_r-R^{1'}, NCH₂R^{1'}, OCH₂R^{1'}, SCH₂R^{1'}, N(CH₂)₂(CH₂)_tR^{1'}, O(CH₂)₂(CH₂)_tR^{1'}, and S(CH₂)₂(CH₂)_tR^{1'}, or combined to form a 5-8 membered saturated, partially saturated or unsaturated ring substituted with 0-2 R⁴ and which contains from 0-2 heteroatoms selected from the group consisting of N, O, and S;

30

35 R^{1'} is selected from H, C₁₋₃ alkyl, halo, (CF₂)_rCF₃, OR², NR²R^{2a}, C(O)R^{2c}, OC(O)R², (CF₂)_rCO₂R^{2c}, S(O)_pR^{2b}, NR²(CH₂)_rOR², NR²C(O)R^{2b}, NR²C(O)NHR^{2b}, NR²C(O)₂R^{2a}, OC(O)NR^{2b}, C(O)NR²R^{2a}, SO₂NR²R^{2a}, NR²SO₂R^{2b}, C₃₋₆

carbocyclic residue substituted with 0-2 R⁴, and 5-10 membered heterocyclic system containing from 1-4 heteroatoms selected from the group consisting of N, O, and S substituted with 0-2 R⁴;

5

R¹ is selected from H, C(O)R^{2b}, C(O)NR²R^{2a}, S(O)R^{2b}, S(O)₂R^{2b}, and SO₂NR²R^{2a};

10

R², at each occurrence, is selected from H, CF₃, C₁₋₆ alkyl, benzyl, C₃₋₆ carbocyclic residue substituted with 0-2 R^{4b}, and 5-6 membered heterocyclic system containing from 1-4 heteroatoms selected from the group consisting of N, O, and S substituted with 0-2 R^{4b};

15

R^{2a}, at each occurrence, is selected from H, CF₃, C₁₋₆ alkyl, benzyl, C₃₋₆ carbocyclic residue substituted with 0-2 R^{4b}, and 5-6 membered heterocyclic system containing from 1-4 heteroatoms selected from the group consisting of N, O, and S substituted with 0-2 R^{4b};

20

R^{2b}, at each occurrence, is selected from CF₃, C₁₋₄ alkoxy, C₁₋₆ alkyl, benzyl, C₃₋₆ carbocyclic residue substituted with 0-2 R^{4b}, and 5-6 membered heterocyclic system containing from 1-4 heteroatoms selected from the group consisting of N, O, and S substituted with 0-2 R^{4b};

25

R^{2c}, at each occurrence, is selected from CF₃, OH, C₁₋₄ alkoxy, C₁₋₆ alkyl, benzyl, C₃₋₆ carbocyclic residue substituted with 0-2 R^{4b}, and 5-6 membered heterocyclic system containing from 1-4 heteroatoms selected from the group consisting of N, O, and S substituted with 0-2 R^{4b};

30

alternatively, R² and R^{2a} combine to form a 5 or 6 membered saturated, partially saturated or unsaturated ring substituted with 0-2 R^{4b} which contains from 0-1 additional heteroatoms selected from the group consisting of N, O, and S;

35

R³, at each occurrence, is selected from H, C₁₋₄ alkyl, and phenyl;

5 R^{3a}, at each occurrence, is selected from H, C₁₋₄ alkyl, and phenyl;

A is selected from:

C₃₋₁₀ carbocyclic residue substituted with 0-2 R⁴, and
5-10 membered heterocyclic system containing from 1-4
10 heteroatoms selected from the group consisting of N, O, and S substituted with 0-2 R⁴;

B is selected from:

X-Y, NR²R^{2a}, C(=NR²)NR²R^{2a}, NR²C(=NR²)NR²R^{2a},
15 C₃₋₁₀ carbocyclic residue substituted with 0-2 R^{4a}, and
5-10 membered heterocyclic system containing from 1-4
heteroatoms selected from the group consisting of N, O, and S substituted with 0-2 R^{4a};

20 X is selected from C₁₋₄ alkylene, -CR²(CR²R^{2b})(CH₂)_t-, -C(O)-, -C(=NR)-, -CR²(NR¹R²)-, -CR²(OR²)-, -CR²(SR²)-, -C(O)CR²R^{2a}-, -CR²R^{2a}C(O)-, -S(O)_p-, -S(O)_pCR²R^{2a}-, -CR²R^{2a}S(O)_p-, -S(O)₂NR²-, -NR²S(O)₂-, -NR²S(O)₂CR²R^{2a}-, -CR²R^{2a}S(O)₂NR²-, -NR²S(O)₂NR²-, -C(O)NR²-, -NR²C(O)-, 25 -C(O)NR²CR²R^{2a}-, -NR²C(O)CR²R^{2a}-, -CR²R^{2a}C(O)NR²-, -CR²R^{2a}NR²C(O)-, -NR²C(O)O-, -OC(O)NR²-, -NR²C(O)NR²-, -NR²-, -NR²CR²R^{2a}-, -CR²R^{2a}NR²-, O-, -CR²R^{2a}O-, and -OCR²R^{2a}-;

30 Y is selected from:

(CH₂)_rNR²R^{2a}, provided that X-Y do not form a N-N, O-N, or S-N bond,
C₃₋₁₀ carbocyclic residue substituted with 0-2 R^{4a}, and
5-10 membered heterocyclic system containing from 1-4
35 heteroatoms selected from the group consisting of N, O, and S substituted with 0-2 R^{4a};

R⁴, at each occurrence, is selected from =O, (CH₂)_rOR², halo, C₁₋₄ alkyl, -CN, NO₂, (CH₂)_rNR²R^{2a}, (CH₂)_rC(O)R^{2b}, NR²C(O)R^{2b}, C(O)NR²R^{2a}, NR²C(O)NR²R^{2a}, CH(=NR²)NR²R^{2a}, NHC(=NR²)NR²R^{2a}, SO₂NR²R^{2a}, NR²SO₂NR²R^{2a}, NR²SO₂-C₁₋₄ alkyl, NR²SO₂R⁵, S(O)_pR⁵, (CF₂)_rCF₃, NCH₂R^{1''}, OCH₂R^{1''}, SCH₂R^{1''}, N(CH₂)₂(CH₂)_tR^{1'}, O(CH₂)₂(CH₂)_tR^{1'}, and S(CH₂)₂(CH₂)_tR^{1'},

alternatively, one R⁴ is a 5-6 membered aromatic heterocycle containing from 1-4 heteroatoms selected from the group consisting of N, O, and S;

R^{4a}, at each occurrence, is selected from =O, (CH₂)_rOR², halo, C₁₋₄ alkyl, -CN, NO₂, (CH₂)_rNR²R^{2a}, (CH₂)_rC(O)R^{2b}, NR²C(O)R^{2b}, C(O)NR²R^{2a}, NR²C(O)NR²R^{2a}, CH(=NR²)NR²R^{2a}, NHC(=NR²)NR²R^{2a}, SO₂NR²R^{2a}, NR²SO₂NR²R^{2a}, NR²SO₂-C₁₋₄ alkyl, NR²SO₂R⁵, S(O)_pR⁵, and (CF₂)_rCF₃;

alternatively, one R^{4a} is a 5-6 membered aromatic heterocycle containing from 1-4 heteroatoms selected from the group consisting of N, O, and S substituted with 0-1 R⁵;

R^{4b}, at each occurrence, is selected from =O, (CH₂)_rOR³, halo, C₁₋₄ alkyl, -CN, NO₂, (CH₂)_rNR³R^{3a}, (CH₂)_rC(O)R³, NR³C(O)R^{3a}, C(O)NR³R^{3a}, NR³C(O)NR³R^{3a}, CH(=NR³)NR³R^{3a}, NH³C(=NR³)NR³R^{3a}, SO₂NR³R^{3a}, NR³SO₂NR³R^{3a}, NR³SO₂-C₁₋₄ alkyl, NR³SO₂CF₃, NR³SO₂-phenyl, S(O)_pCF₃, S(O)_p-C₁₋₄ alkyl, S(O)_p-phenyl, and (CF₂)_rCF₃;

R⁵, at each occurrence, is selected from CF₃, C₁₋₆ alkyl, phenyl substituted with 0-2 R⁶, and benzyl substituted with 0-2 R⁶;

R⁶, at each occurrence, is selected from H, OH, (CH₂)_rOR², halo, C₁₋₄ alkyl, CN, NO₂, (CH₂)_rNR²R^{2a}, (CH₂)_rC(O)R^{2b}, NR²C(O)R^{2b}, NR²C(O)NR²R^{2a}, CH(=NH)NH₂, NHC(=NH)NH₂, SO₂NR²R^{2a}, NR²SO₂NR²R^{2a}, and NR²SO₂C₁₋₄ alkyl;

R⁷, at each occurrence, is selected from H, OH, C₁₋₆ alkyl, C₁₋₆ alkylcarbonyl, C₁₋₆ alkoxy, C₁₋₄ alkoxycarbonyl, (CH₂)_n-phenyl, C₆₋₁₀ aryloxy, C₆₋₁₀ aryloxycarbonyl, C₆₋₁₀ arylmethylcarbonyl, C₁₋₄ alkylcarbonyloxy C₁₋₄ alkoxy
5 alkoxy carbonyl, C₆₋₁₀ arylcarbonyloxy C₁₋₄ alkoxy carbonyl, C₁₋₆ alkylaminocarbonyl, phenylaminocarbonyl, and phenyl C₁₋₄ alkoxy carbonyl;

R⁸, at each occurrence, is selected from H, C₁₋₆ alkyl and
10 (CH₂)_n-phenyl;

alternatively, R⁷ and R⁸ combine to form a 5 or 6 membered saturated, ring which contains from 0-1 additional heteroatoms selected from the group consisting of N, O,
15 and S;

R⁹, at each occurrence, is selected from H, C₁₋₆ alkyl and (CH₂)_n-phenyl;

20 n, at each occurrence, is selected from 0, 1, 2, and 3;

m, at each occurrence, is selected from 0, 1, and 2;

p, at each occurrence, is selected from 0, 1, and 2;
25 r, at each occurrence, is selected from 0, 1, 2, and 3;


s, at each occurrence, is selected from 0, 1, and 2; and,

30 t, at each occurrence, is selected from 0 and 1;


provided that D-E-G-(CH₂)_s- and -Z-A-B are not both benzamidines.

35


[2] In a preferred embodiment, the present invention provides novel compounds of formulae Ia-If:




1a




1b




1c



1d



1e



1f

wherein, groups D-E- and -Z-A-B are attached to adjacent atoms on the ring;

- 5 Z is selected from a CH_2O , OCH_2 , CH_2NH , NHCH_2 , $\text{C}(\text{O})$, $\text{CH}_2\text{C}(\text{O})$, $\text{C}(\text{O})\text{CH}_2$, $\text{NHC}(\text{O})$, $\text{C}(\text{O})\text{NH}$, $\text{CH}_2\text{S}(\text{O})_2$, $\text{S}(\text{O})_2(\text{CH}_2)$, SO_2NH , and NHSO_2 , provided that Z does not form a N-N, N-O, NCH_2N , or NCH_2O bond with ring M or group A;
- 10 A is selected from one of the following carbocyclic and heterocyclic systems which are substituted with 0-2 R^4 ;
phenyl, piperidinyl, piperazinyl, pyridyl,
pyrimidyl, furanyl, morpholinyl, thiophenyl, pyrrolyl,
pyrrolidinyl, oxazolyl, isoxazolyl, thiazolyl,
15 isothiazolyl, pyrazolyl, imidazolyl, oxadiazolyl,
thiadiazolyl, triazolyl, 1,2,3-oxadiazolyl,
1,2,4-oxadiazolyl, 1,2,5-oxadiazolyl, 1,3,4-oxadiazolyl,
1,2,3-thiadiazolyl, 1,2,4-thiadiazolyl,
1,2,5-thiadiazolyl, 1,3,4-thiadiazolyl, 1,2,3-triazolyl,
20 1,2,4-triazolyl, 1,2,5-triazolyl, 1,3,4-triazolyl,
benzofuranyl, benzothiofuranyl, indolyl, benzimidazolyl,
benzoxazolyl, benzthiazolyl, indazolyl, benzisoxazolyl,
benzisothiazolyl, and isoindazolyl;
- 25 B is selected from: Y, X-Y, NR^2R^{2a} , $\text{C}(=\text{NR}^2)\text{NR}^2\text{R}^{2a}$, and $\text{NR}^2\text{C}(=\text{NR}^2)\text{NR}^2\text{R}^{2a}$;
- X is selected from C_{1-4} alkylene, $-\text{C}(\text{O})-$, $-\text{C}(=\text{NR})-$,
 $-\text{CR}^2(\text{NR}^2\text{R}^{2a})-$, $-\text{C}(\text{O})\text{CR}^2\text{R}^{2a}-$, $-\text{CR}^2\text{R}^{2a}\text{C}(\text{O})-$, $-\text{C}(\text{O})\text{NR}^2-$,
30 $-\text{NR}^2\text{C}(\text{O})-$, $-\text{C}(\text{O})\text{NR}^2\text{CR}^2\text{R}^{2a}-$, $-\text{NR}^2\text{C}(\text{O})\text{CR}^2\text{R}^{2a}-$,

$-\text{CR}^2\text{R}^{2a}\text{C}(\text{O})\text{NR}^2-$, $-\text{CR}^2\text{R}^{2a}\text{NR}^2\text{C}(\text{O})-$, $-\text{NR}^2\text{C}(\text{O})\text{NR}^2-$, $-\text{NR}^2-$,
 $-\text{NR}^2\text{CR}^2\text{R}^{2a}-$, $-\text{CR}^2\text{R}^{2a}\text{NR}^2-$, O , $-\text{CR}^2\text{R}^{2a}\text{O}-$, and $-\text{OCR}^2\text{R}^{2a}-$;


Y is NR^2R^{2a} , provided that X-Y do not form a N-N or O-N bond;

5

alternatively, Y is selected from one of the following carbocyclic and heterocyclic systems which are substituted with 0-2 R^{4a} ;







cyclopropyl, cyclopentyl, cyclohexyl, phenyl,
 10 piperidinyl, piperazinyl, pyridyl, pyrimidyl, furanyl, morpholinyl, thiophenyl, pyrrolyl, pyrrolidinyl, oxazolyl, isoxazolyl, isoxazolinyl, thiazolyl, isothiazolyl, pyrazolyl, imidazolyl, oxadiazolyl, thiadiazolyl, triazolyl, 1,2,3-oxadiazolyl,
 15 1,2,4-oxadiazolyl, 1,2,5-oxadiazolyl, 1,3,4-oxadiazolyl, 1,2,3-thiadiazolyl, 1,2,4-thiadiazolyl, 1,2,5-thiadiazolyl, 1,3,4-thiadiazolyl, 1,2,3-triazolyl, 1,2,4-triazolyl, 1,2,5-triazolyl, 1,3,4-triazolyl, benzofuranyl, benzothiofuranyl, indolyl, benzimidazolyl,
 20 benzoxazolyl, benzthiazolyl, indazolyl, benzisoxazolyl, benzisothiazolyl, and isoindazolyl;

alternatively, Y is selected from the following bicyclic heteroaryl ring systems:



K is selected from O, S, NH, and N.

30 [3] In a more preferred embodiment, the present invention provides novel compounds of formulae Ib and Ic:

Ib₁Ib₂Ib₃Ib₄Ic₁Ic₂

wherein;

5 J is O or S; and,

Z is selected from a C(O), CH₂C(O), C(O)CH₂, NHC(O), C(O)NH,
C(O)N(CH₃), CH₂S(O)₂, S(O)₂(CH₂), SO₂NH, and NHSO₂,
provided that Z does not form a N-N or NCH₂N bond with
10 ring M or group A.

[4] In an even more preferred embodiment, the present
invention provides novel compounds of formulae Ib and Ic,
15 wherein;

E is phenyl substituted with R or 2-pyridyl substituted with
R;

20 D is selected from NH₂, C(O)NH₂, C(=NH)NH₂, CH₂NH₂, CH₂NHCH₃,
CH(CH₃)NH₂, and C(CH₃)₂NH₂, provided that D is substituted
meta or para to ring M on E; and,

R is selected from H, OCH₃, Cl, and F.

[5] In a further preferred embodiment, the present invention
provides novel compounds of formulae Ib and Ic, wherein;

D-E is selected from 3-aminophenyl, 3-amidinophenyl, 3-aminomethylphenyl, 3-aminocarbonylphenyl, 3-(methylaminomethyl)phenyl, 3-(1-aminoethyl)phenyl, 3-(2-amino-2-propyl)phenyl, 4-chloro-3-aminophenyl, 4-chloro-3-amidinophenyl, 4-chloro-3-aminomethylphenyl, 4-chloro-3-(methylaminomethyl)phenyl, 4-fluoro-3-aminophenyl, 4-fluoro-3-amidinophenyl, 4-fluoro-3-aminomethylphenyl, 4-fluoro-3-(methylaminomethyl)phenyl, 6-aminopyrid-2-yl, 6-amidinopyrid-2-yl, 6-aminomethylpyrid-2-yl, 6-aminocarbonylpyrid-2-yl, 6-(methylaminomethyl)pyrid-2-yl, 6-(1-aminoethyl)pyrid-2-yl, and 6-(2-amino-2-propyl)pyrid-2-yl.

[6] In another even more preferred embodiment, the present invention provides novel compounds of formulae Ib and Ic, wherein;

Z is $C(O)CH_2$ and $CONH$, provided that Z does not form a N-N bond with group A;

A is selected from phenyl, pyridyl, and pyrimidyl, and is substituted with 0-2 R^4 ; and,

B is selected from X-Y, phenyl, pyrrolidino, morpholino, 1,2,3-triazolyl, and imidazolyl, and is substituted with 0-1 R^{4a} ;

R^4 , at each occurrence, is selected from OH, $(CH_2)_rOR^2$, halo, C_{1-4} alkyl, $(CH_2)_rNR^2R^{2a}$, and $(CF_2)_rCF_3$;

R^{4a} is selected from C_{1-4} alkyl, CF_3 , $S(O)_pR^5$, $SO_2NR^2R^{2a}$, and 1- CF_3 -tetrazol-2-yl;

R^5 , at each occurrence, is selected from CF_3 , C_{1-6} alkyl, phenyl, and benzyl;

X is CH_2 or $C(O)$; and,

Y is selected from pyrrolidino and morpholino.

- 5 [7] In another further preferred embodiment, the present invention provides novel compounds of formulae Ib and Ic, wherein;

10 A is selected from the group: phenyl, 2-pyridyl, 3-pyridyl, 2-pyrimidyl, 2-Cl-phenyl, 3-Cl-phenyl, 2-F-phenyl, 3-F-phenyl, 2-methylphenyl, 2-aminophenyl, and 2-methoxyphenyl; and,

15 B is selected from the group: 2-CF₃-phenyl, 2-(aminosulfonyl)phenyl, 2-(methylaminosulfonyl)phenyl, 2-(dimethylaminosulfonyl)phenyl, 1-pyrrolidinocarbonyl, 2-(methylsulfonyl)phenyl, 4-morpholino, 2-(1'-CF₃-tetrazol-2-yl)phenyl, 4-morpholinocarbonyl, 2-methyl-1-imidazolyl, 5-methyl-1-imidazolyl, 2-methylsulfonyl-1-imidazolyl and, 20 5-methyl-1,2,3-triazolyl.

- 25 [8] In another even more preferred embodiment, the present invention provides novel compounds of formulae Ib and Ic, wherein;

E is phenyl substituted with R or 2-pyridyl substituted with R;

30 D is selected from NH₂, C(O)NH₂, C(=NH)NH₂, CH₂NH₂, CH₂NHCH₃, CH(CH₃)NH₂, and C(CH₃)₂NH₂, provided that D is substituted meta or para to ring M on E; and,

R is selected from H, OCH₃, Cl, and F;

35 Z is C(O)CH₂ and CONH, provided that Z does not form a N-N bond with group A;

A is selected from phenyl, pyridyl, and pyrimidyl, and is substituted with 0-2 R^4 ; and,

5 B is selected from X-Y, phenyl, pyrrolidino, morpholino, 1,2,3-triazolyl, and imidazolyl, and is substituted with 0-1 R^{4a} ;

R^4 , at each occurrence, is selected from OH, $(CH_2)_rOR^2$, halo, C_{1-4} alkyl, $(CH_2)_rNR^2R^{2a}$, and $(CF_2)_rCF_3$;

10 R^{4a} is selected from C_{1-4} alkyl, CF_3 , $S(O)_pR^5$, $SO_2NR^2R^{2a}$, and 1- CF_3 -tetrazol-2-yl;

15 R^5 , at each occurrence, is selected from CF_3 , C_{1-6} alkyl, phenyl, and benzyl;

X is CH_2 or $C(O)$; and,

20 Y is selected from pyrrolidino and morpholino.

[9] In another further preferred embodiment, the present invention provides novel compounds of formulae Ib and Ic, wherein;

25 D-E is selected from 3-aminophenyl, 3-amidinophenyl, 3-aminomethylphenyl, 3-aminocarbonylphenyl, 3-(methylaminomethyl)phenyl, 3-(1-aminoethyl)phenyl, 3-(2-amino-2-propyl)phenyl, 4-chloro-3-aminophenyl, 4-chloro-3-amidinophenyl, 4-chloro-3-aminomethylphenyl, 4-chloro-3-(methylaminomethyl)phenyl, 4-fluoro-3-aminophenyl, 4-fluoro-3-amidinophenyl, 4-fluoro-3-aminomethylphenyl, 4-fluoro-3-(methylaminomethyl)phenyl, 6-aminopyrid-2-yl, 6-amidinopyrid-2-yl, 6-aminomethylpyrid-2-yl, 6-aminocarbonylpyrid-2-yl, 6-(methylaminomethyl)pyrid-2-yl, 30 6-(1-aminoethyl)pyrid-2-yl, 6-(2-amino-2-propyl)pyrid-2-yl;
35

A is selected from the group: phenyl, 2-pyridyl, 3-pyridyl, 2-pyrimidyl, 2-Cl-phenyl, 3-Cl-phenyl, 2-F-phenyl, 3-F-phenyl, 2-methylphenyl, 2-aminophenyl, and 2-methoxyphenyl; and,

5

B is selected from the group: 2-CF₃-phenyl, 2-(aminosulfonyl)phenyl, 2-(methylaminosulfonyl)phenyl, 2-(dimethylaminosulfonyl)phenyl, 1-pyrrolidinocarbonyl, 2-(methylsulfonyl)phenyl, 4-morpholino, 2-(1'-CF₃-tetrazol-2-yl)phenyl, 4-morpholinocarbonyl, 2-methyl-1-imidazolyl, 5-methyl-1-imidazolyl, 2-methylsulfonyl-1-imidazolyl and, 5-methyl-1,2,3-triazolyl.

10

15 [10] In a still further preferred embodiment, the present invention provides a novel compound of formula Ib₁.

20 [11] In another still further preferred embodiment, the present invention provides a novel compound of formula Ib₂.

[12] In another still further preferred embodiment, the present invention provides a novel compound of formula Ib₃.

25

[13] In another still further preferred embodiment, the present invention provides a novel compound of formula Ib₄.

30

[14] In another still further preferred embodiment, the present invention provides a novel compound of formula Ic₁.

35 [15] In another still further preferred embodiment, the present invention provides a novel compound of formula Ic₂.

[16] In another even more preferred embodiment, the present invention provides novel compounds of formulae Ib and Ic, wherein;

- 5 D is selected from $C(=NR^8)NR^7R^9$, $C(O)NR^7R^8$, NR^7R^8 , and $CH_2NR^7R^8$, provided that D is substituted meta or para to ring M on E;

E is phenyl substituted with R or pyridyl substituted with R;

10

R is selected from H, Cl, F, OR^3 , CH_3 , CH_2CH_3 , OCF_3 , and CF_3 ;

Z is selected from $C(O)$, $CH_2C(O)$, $C(O)CH_2$, $NHC(O)$, and $C(O)NH$, provided that Z does not form a N-N bond with ring M or group A;

15

R^{1a} and R^{1b} are independently absent or selected from $-(CH_2)_r-R^{1'}$, $NCH_2R^{1''}$, $OCH_2R^{1''}$, $SCH_2R^{1''}$, $N(CH_2)_2(CH_2)_tR^{1'}$, $O(CH_2)_2(CH_2)_tR^{1'}$, and $S(CH_2)_2(CH_2)_tR^{1'}$, or combined to form a 5-8 membered saturated, partially saturated or unsaturated ring substituted with 0-2 R^4 and which contains from 0-2 heteroatoms selected from the group consisting of N, O, and S;

20

- 25 $R^{1'}$, at each occurrence, is selected from H, C_{1-3} alkyl, halo, $(CF_2)_rCF_3$, OR^2 , NR^2R^{2a} , $C(O)R^{2c}$, $(CF_2)_rCO_2R^{2c}$, $S(O)_pR^{2b}$, $NR^2(CH_2)_rOR^2$, $NR^2C(O)R^{2b}$, $NR^2C(O)_2R^{2b}$, $C(O)NR^2R^{2a}$, $SO_2NR^2R^{2a}$, and $NR^2SO_2R^{2b}$;

- 30 A is selected from one of the following carbocyclic and heterocyclic systems which are substituted with 0-2 R^4 ; phenyl, piperidinyl, piperazinyl, pyridyl, pyrimidyl, furanyl, morpholinyl, thiophenyl, pyrrolyl, pyrrolidinyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, pyrazolyl, and imidazolyl;

35

B is selected from: Y, X-Y, NR^2R^{2a} , $C(=NR^2)NR^2R^{2a}$, and $NR^2C(=NR^2)NR^2R^{2a}$;

X is selected from CH_2 , $-\text{CR}^2(\text{CR}^2\text{R}^{2b})(\text{CH}_2)_t-$, $-\text{C}(\text{O})-$, $-\text{C}(=\text{NR})-$,
5 $-\text{CH}(\text{NR}^2\text{R}^{2a})-$, $-\text{C}(\text{O})\text{NR}^2-$, $-\text{NR}^2\text{C}(\text{O})-$, $-\text{NR}^2\text{C}(\text{O})\text{NR}^2-$, $-\text{NR}^2-$,
and O;

Y is NR^2R^{2a} , provided that X-Y do not form a N-N or O-N bond;

alternatively, Y is selected from one of the following
carbocyclic and heterocyclic systems which are
10 substituted with 0-2 R^{4a} ;

phenyl, piperidinyl, piperazinyl, pyridyl,
pyrimidyl, furanyl, morpholinyl, thiophenyl, pyrrolyl,
pyrrolidinyl, oxazolyl, isoxazolyl, isoxazolinyl,
thiazolyl, isothiazolyl, pyrazolyl, imidazolyl,
15 oxadiazolyl, thiadiazolyl, triazolyl, 1,2,3-oxadiazolyl,
1,2,4-oxadiazolyl, 1,2,5-oxadiazolyl, 1,3,4-oxadiazolyl,
1,2,3-thiadiazolyl, 1,2,4-thiadiazolyl,
1,2,5-thiadiazolyl, 1,3,4-thiadiazolyl, 1,2,3-triazolyl,
20 1,2,4-triazolyl, 1,2,5-triazolyl, and 1,3,4-triazolyl;

R^4 , at each occurrence, is selected from =O, OH, Cl, F, C_{1-4}
alkyl, $(\text{CH}_2)_r\text{NR}^2\text{R}^{2a}$, $(\text{CH}_2)_r\text{C}(\text{O})\text{R}^{2b}$, $\text{NR}^2\text{C}(\text{O})\text{R}^{2b}$, $\text{C}(\text{O})\text{NR}^2\text{R}^{2a}$,
 $\text{CH}(=\text{NH})\text{NH}_2$, $\text{NHC}(=\text{NH})\text{NH}_2$, $\text{SO}_2\text{NR}^2\text{R}^{2a}$, $\text{NR}^2\text{SO}_2-\text{C}_{1-4}$ alkyl,
25 $\text{NR}^2\text{SO}_2\text{R}^5$, $\text{S}(\text{O})_p\text{R}^5$, and $(\text{CF}_2)_r\text{CF}_3$;

R^{4a} , at each occurrence, is selected from =O, OH, Cl, F, C_{1-4}
alkyl, $(\text{CH}_2)_r\text{NR}^2\text{R}^{2a}$, $(\text{CH}_2)_r\text{C}(\text{O})\text{R}^{2b}$, $\text{NR}^2\text{C}(\text{O})\text{R}^{2b}$, $\text{C}(\text{O})\text{NR}^2\text{R}^{2a}$,
 $\text{CH}(=\text{NH})\text{NH}_2$, $\text{NHC}(=\text{NH})\text{NH}_2$, $\text{SO}_2\text{NR}^2\text{R}^{2a}$, $\text{NR}^2\text{SO}_2-\text{C}_{1-4}$ alkyl,
30 $\text{NR}^2\text{SO}_2\text{R}^5$, $\text{S}(\text{O})_p\text{R}^5$, $(\text{CF}_2)_r\text{CF}_3$, and 1- CF_3 -tetrazol-2-yl;

R^5 , at each occurrence, is selected from CF_3 , C_{1-6} alkyl,
phenyl substituted with 0-2 R^6 , and benzyl substituted
with 0-2 R^6 ;

35 R^6 , at each occurrence, is selected from H, =O, OH, OR^2 , Cl, F,
 CH_3 , CN, NO_2 , $(\text{CH}_2)_r\text{NR}^2\text{R}^{2a}$, $(\text{CH}_2)_r\text{C}(\text{O})\text{R}^{2b}$, $\text{NR}^2\text{C}(\text{O})\text{R}^{2b}$,
 $\text{CH}(=\text{NH})\text{NH}_2$, $\text{NHC}(=\text{NH})\text{NH}_2$, and $\text{SO}_2\text{NR}^2\text{R}^{2a}$;

R⁷, at each occurrence, is selected from H, OH, C₁₋₆ alkyl, C₁₋₆ alkylcarbonyl, C₁₋₆ alkoxy, C₁₋₄ alkoxycarbonyl, benzyl, C₆₋₁₀ aryloxy, C₆₋₁₀ aryloxycarbonyl, C₆₋₁₀ arylmethylcarbonyl, C₁₋₄ alkylcarbonyloxy C₁₋₄ alkoxycarbonyl, C₆₋₁₀ arylcarbonyloxy C₁₋₄ alkoxycarbonyl, C₁₋₆ alkylaminocarbonyl, phenylaminocarbonyl, and phenyl C₁₋₄ alkoxycarbonyl;

R⁸, at each occurrence, is selected from H, C₁₋₆ alkyl and benzyl; and

alternatively, R⁷ and R⁸ combine to form a morpholino group; and,

R⁹, at each occurrence, is selected from H, C₁₋₆ alkyl and benzyl.

[17] In a another further preferred embodiment, the present invention provides novel compounds of formulae Ib and Ic, wherein;

E is phenyl substituted with R or 2-pyridyl substituted with R;

R is selected from H, Cl, F, OCH₃, CH₃, OCF₃, and CF₃;

Z is selected from a C(O)CH₂ and C(O)NH, provided that Z does not form a N-N bond with group A;

R^{1a} is selected from H, CH₃, CH₂CH₃, Cl, F, CF₃, OCH₃, NR²R^{2a}, S(O)_pR^{2b}, CH₂S(O)_pR^{2b}, CH₂NR²S(O)_pR^{2b}, C(O)R^{2c}, CH₂C(O)R^{2c}, C(O)NR²R^{2a}, and SO₂NR²R^{2a};

R^{1b} is selected from H, CH₃, CH₂CH₃, Cl, F, CF₃, OCH₃, NR²R^{2a}, S(O)_pR^{2b}, CH₂S(O)_pR^{2b}, CH₂NR²S(O)_pR^{2b}, C(O)R^{2c}, CH₂C(O)R^{2c}, C(O)NR²R^{2a}, and SO₂NR²R^{2a};

A is selected from one of the following carbocyclic and heterocyclic systems which are substituted with 0-2 R⁴; phenyl, pyridyl, pyrimidyl, furanyl, thiophenyl, pyrrolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, pyrazolyl, and imidazolyl;

B is selected from: Y and X-Y;

X is selected from CH₂, -CR²(CR²R^{2b})-, -C(O)-, -C(=NR)-, -CH(NR²R^{2a})-, -C(O)NR²-, -NR²C(O)-, -NR²C(O)NR²-, -NR²-, and O;

Y is NR²R^{2a}, provided that X-Y do not form a N-N or O-N bond;

alternatively, Y is selected from one of the following carbocyclic and heterocyclic systems which are substituted with 0-2 R^{4a};

phenyl, piperidinyl, piperazinyl, pyridyl, pyrimidyl, furanyl, morpholinyl, thiophenyl, pyrrolyl, pyrrolidinyl, oxazolyl, isoxazolyl, isoxazolinyl, thiazolyl, isothiazolyl, pyrazolyl, imidazolyl, oxadiazolyl, thiadiazolyl, triazolyl, 1,2,3-oxadiazolyl, 1,2,4-oxadiazolyl, 1,2,5-oxadiazolyl, 1,3,4-oxadiazolyl, 1,2,3-thiadiazolyl, 1,2,4-thiadiazolyl, 1,2,5-thiadiazolyl, 1,3,4-thiadiazolyl, 1,2,3-triazolyl, 1,2,4-triazolyl, 1,2,5-triazolyl, and 1,3,4-triazolyl;

R², at each occurrence, is selected from H, CF₃, CH₃, benzyl, and phenyl;

R^{2a}, at each occurrence, is selected from H, CF₃, CH₃, benzyl, and phenyl;

R^{2b}, at each occurrence, is selected from CF₃, OCH₃, CH₃, benzyl, and phenyl;

R^{2c}, at each occurrence, is selected from CF₃, OH, OCH₃, CH₃, benzyl, and phenyl;

alternatively, R^2 and R^{2a} combine to form a 5 or 6 membered saturated, partially unsaturated, or unsaturated ring which contains from 0-1 additional heteroatoms selected from the group consisting of N, O, and S;

R^3 , at each occurrence, is selected from H, CH_3 , CH_2CH_3 , and phenyl;

R^{3a} , at each occurrence, is selected from H, CH_3 , CH_2CH_3 , and phenyl;

R^4 , at each occurrence, is selected from OH, Cl, F, CH_3 , CH_2CH_3 , NR^2R^{2a} , $CH_2NR^2R^{2a}$, $C(O)R^{2b}$, $NR^2C(O)R^{2b}$, $C(O)NR^2R^{2a}$, and CF_3 ;

R^{4a} , at each occurrence, is selected from OH, Cl, F, CH_3 , CH_2CH_3 , NR^2R^{2a} , $CH_2NR^2R^{2a}$, $C(O)R^{2b}$, $C(O)NR^2R^{2a}$, $SO_2NR^2R^{2a}$, $S(O)_pR^5$, CF_3 , and 1- CF_3 -tetrazol-2-yl;

R^5 , at each occurrence, is selected from CF_3 , C_{1-6} alkyl, phenyl substituted with 0-2 R^6 , and benzyl substituted with 1 R^6 ;

R^6 , at each occurrence, is selected from H, OH, OCH_3 , Cl, F, CH_3 , CN, NO_2 , NR^2R^{2a} , $CH_2NR^2R^{2a}$, and $SO_2NR^2R^{2a}$;

R^7 , at each occurrence, is selected from H, OH, C_{1-3} alkyl, C_{1-3} alkylcarbonyl, C_{1-3} alkoxy, C_{1-4} alkoxy carbonyl, benzyl, phenoxy, phenoxycarbonyl, benzylcarbonyl, C_{1-4} alkylcarbonyloxy, C_{1-4} alkoxy carbonyl, phenylcarbonyloxy, C_{1-4} alkoxy carbonyl, C_{1-6} alkylaminocarbonyl, phenylaminocarbonyl, and phenyl C_{1-4} alkoxy carbonyl;

R^8 , at each occurrence, is selected from H, CH_3 , and benzyl; and,

alternatively, R^7 and R^8 combine to form a morpholino group;

R⁹, at each occurrence, is selected from H, CH₃, and benzyl.

5 [18] In a another still further preferred embodiment, the present invention provides novel compounds of formulae Ib and Ic, wherein;

10 R^{1a} is absent or is selected from H, CH₃, CH₂CH₃, Cl, F, CF₃, OCH₃, NR²R^{2a}, S(O)_pR^{2b}, C(O)NR²R^{2a}, CH₂S(O)_pR^{2b}, CH₂NR²S(O)_pR^{2b}, C(O)R^{2c}, CH₂C(O)R^{2c}, and SO₂NR²R^{2a};

15 R^{1b} is absent or is selected from H, CH₃, CH₂CH₃, Cl, F, CF₃, OCH₃, NR²R^{2a}, S(O)_pR^{2b}, C(O)NR²R^{2a}, CH₂S(O)_pR^{2b}, CH₂NR²S(O)_pR^{2b}, C(O)R^{2b}, CH₂C(O)R^{2b}, and SO₂NR²R^{2a};

20 A is selected from one of the following carbocyclic and heterocyclic systems which are substituted with 0-2 R⁴; phenyl, pyridyl, and pyrimidyl;

B is selected from: Y and X-Y;

X is selected from -C(O)- and O;

25 Y is NR²R^{2a}, provided that X-Y do not form a O-N bond;

alternatively, Y is selected from one of the following carbocyclic and heterocyclic systems which are substituted with 0-2 R^{4a};

30 phenyl, piperazinyl, pyridyl, pyrimidyl, morpholinyl, pyrrolidinyl, imidazolyl, and 1,2,3-triazolyl;

35 R², at each occurrence, is selected from H, CF₃, CH₃, benzyl, and phenyl;

R^{2a}, at each occurrence, is selected from H, CF₃, CH₃, benzyl, and phenyl;

R^{2b}, at each occurrence, is selected from CF₃, OCH₃, CH₃, benzyl, and phenyl;

5 R^{2c}, at each occurrence, is selected from CF₃, OH, OCH₃, CH₃, benzyl, and phenyl;

alternatively, R² and R^{2a} combine to form a ring system selected from pyrrolidinyl, piperazinyl and morpholino;

10

R⁴, at each occurrence, is selected from Cl, F, CH₃, NR²R^{2a}, and CF₃;

15 R^{4a}, at each occurrence, is selected from Cl, F, CH₃, SO₂NR²R^{2a}, S(O)_pR⁵, and CF₃; and,

R⁵, at each occurrence, is selected from CF₃ and CH₃.

20 [19] Specifically preferred compounds of the present invention invention is selected from the group:

3-(3-amidinophenyl)-4-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-5-(hydroxymethyl)isoxazole;

25

3-(3-amidinophenyl)-4-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]isoxazole;

30

3-(3-amidinophenyl)-4-[(2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]isoxazole;

3-(3-amidinophenyl)-4-[5-(2-aminosulfonyl)phenylpyrid-2-yl)aminocarbonyl]-5-(methoxymethyl)isoxazole;

35 3-(3-amidinophenyl)-4-[(2'-trifluoromethyl-[1,1']-biphen-4-yl)aminocarbonyl]isoxazole;

3-(3-amidinophenyl)-4-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-5-(trifluoromethyl)isoxazole;

2-acetylamino-4-(3-amidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole;

2-amino-4-(3-amidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole;

2-methyl-4-(3-amidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole;

5-(3-amidinophenyl)-4-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]oxazole;

3-(3-amidinophenyl)-4-[(2'-t-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]isoxazole;

3-(3-amidinophenyl)-4-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-5-(methoxymethyl)-isoxazole;

3-(3-amidinophenyl)-4-[(2'-t-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]isoxazole;

3-(3-amidinophenyl)-4-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-5-(methoxymethyl)isoxazole;

2-methyl-4-(3-amidinophenyl)-5-[(2'-trifluoromethyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole;

2-phenyl-4-(3-amidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole;

3-(3-amidinophenyl)-4-[(3-fluoro-2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]isoxazole;

3-(3-amidinophenyl)-4-[(2'-trifluoromethylthio-[1,1']-biphen-4-yl)aminocarbonyl]isoxazole;

3-(3-amidinophenyl)-5-amino-4-[(2'-aminosulfonyl-[1,1']-
biphen-4-yl)aminocarbonyl]isoxazole;

5 2-(phenylamino)-4-(3-amidinophenyl)-5-[(2'-aminosulfonyl-
[1,1']-biphen-4-yl)aminocarbonyl]thiazole;

2-(benzylamino)-4-(3-amidinophenyl)-5-[(2'-aminosulfonyl-
[1,1']-biphen-4-yl)aminocarbonyl]thiazole;

10

2-(methylamino)-4-(3-amidinophenyl)-5-[(2'-aminosulfonyl-
[1,1']-biphen-4-yl)aminocarbonyl]thiazole;

15

2-(methylamino)-4-(3-carboxamidophenyl)-5-[(2'-aminosulfonyl-
[1,1']-biphen-4-yl)aminocarbonyl]thiazole;

2-methyl-4-(3-amidinophenyl)-5-[[5-(2'-aminosulfonylphenyl-1-
yl)pyridin-2-yl]aminocarbonyl]thiazole;

20

2-methyl-4-(3-(carboxamido)phenyl)-5-[[5-(2'-
aminosulfonylphenyl-1-yl)pyridin-2-
yl]aminocarbonyl]thiazole;

25

2-(3-pyridyl)-4-(3-amidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-
biphen-4-yl)aminocarbonyl]thiazole;

2-(3-pyridyl)-4-(3-carboxamidophenyl)-5-[(2'-aminosulfonyl-
[1,1']-biphen-4-yl)aminocarbonyl]thiazole;

30

2-chloro-4-(3-amidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-
biphen-4-yl)aminocarbonyl]thiazole;

2-chloro-4-(3-carboxamidophenyl)-5-[(2'-aminosulfonyl-[1,1']-
biphen-4-yl)aminocarbonyl]thiazole;

35

2-chloro-4-(3-amidinophenyl)-5-[[5-(2'-aminosulfonylphenyl-1-
yl)pyridin-2-yl]aminocarbonyl]thiazole;

2-chloro-4-(3-(carboxamido)phenyl)-5-[[5-(2'-aminosulfonylphenyl-1-yl)pyridin-2-yl]aminocarbonyl]thiazole;

5 2-hydroxy-4-(3-amidinophenyl)-5-[[5-(2'-aminosulfonylphenyl-1-yl)pyridin-2-yl]aminocarbonyl]thiazole;

2-chloro-4-(3-aminophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole;

10

2-amino-4-[(3-amino-4-chloro)phenyl]-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole;

15

2-chloro-4-[(3-amino-4-chloro)phenyl]-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole; and,

2-amino-4-[(3-aminomethyl)phenyl]-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole;

20

and a pharmaceutically acceptable salt thereof.

In a second embodiment, the present invention provides novel pharmaceutical compositions, comprising: a
25 pharmaceutically acceptable carrier and a therapeutically effective amount of a compound of formula (I) or a pharmaceutically acceptable salt form thereof.

30

In a third embodiment, the present invention provides a novel method for treating or preventing a thromboembolic disorder, comprising: administering to a patient in need thereof a therapeutically effective amount of a compound of formula (I) or a pharmaceutically acceptable salt form thereof.

35

DEFINITIONS

The compounds herein described may have asymmetric centers. Compounds of the present invention containing an

asymmetrically substituted atom may be isolated in optically active or racemic forms. It is well known in the art how to prepare optically active forms, such as by resolution of racemic forms or by synthesis from optically active starting materials. Many geometric isomers of olefins, C=N double bonds, and the like can also be present in the compounds described herein, and all such stable isomers are contemplated in the present invention. Cis and trans geometric isomers of the compounds of the present invention are described and may be isolated as a mixture of isomers or as separated isomeric forms. All chiral, diastereomeric, racemic forms and all geometric isomeric forms of a structure are intended, unless the specific stereochemistry or isomeric form is specifically indicated.

The term "substituted," as used herein, means that any one or more hydrogens on the designated atom is replaced with a selection from the indicated group, provided that the designated atom's normal valency is not exceeded, and that the substitution results in a stable compound. When a substituent is keto (i.e., =O), then 2 hydrogens on the atom are replaced.

When any variable (e.g., R^6) occurs more than one time in any constituent or formula for a compound, its definition at each occurrence is independent of its definition at every other occurrence. Thus, for example, if a group is shown to be substituted with 0-2 R^6 , then said group may optionally be substituted with up to two R^6 groups and R^6 at each occurrence is selected independently from the definition of R^6 . Also, combinations of substituents and/or variables are permissible only if such combinations result in stable compounds.

When a bond to a substituent is shown to cross a bond connecting two atoms in a ring, then such substituent may be bonded to any atom on the ring. When a substituent is listed without indicating the atom via which such substituent is bonded to the rest of the compound of a given formula, then such substituent may be bonded via any atom in such substituent. Combinations of substituents and/or variables are permissible only if such combinations result in stable compounds.

As used herein, "C₁₋₆ alkyl" is intended to include both branched and straight-chain saturated aliphatic hydrocarbon groups having the specified number of carbon atoms, examples of which include, but are not limited to, methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl, sec-butyl, t-butyl, pentyl, and hexyl; "Alkenyl" is intended to include hydrocarbon chains of either a straight or branched configuration and one or more unsaturated carbon-carbon bonds which may occur in any stable point along the chain, such as ethenyl, propenyl, and the like.

"Halo" or "halogen" as used herein refers to fluoro, chloro, bromo, and iodo; and "counterion" is used to represent a small, negatively charged species such as chloride, bromide, hydroxide, acetate, sulfate, and the like.

As used herein, "carbocycle" or "carbocyclic residue" is intended to mean any stable 3- to 10-membered monocyclic or bicyclic or 7- to 13-membered bicyclic or tricyclic, any of which may be saturated, partially unsaturated, or aromatic. Examples of such carbocycles include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, adamantyl, cyclooctyl, [3.3.0]bicyclooctane, [4.3.0]bicyclononane, [4.4.0]bicyclodecane (decalin), [2.2.2]bicyclooctane, fluorenyl, phenyl, naphthyl, indanyl, adamantyl, or tetrahydronaphthyl (tetralin).

As used herein, the term "heterocycle" or "heterocyclic system" is intended to mean a stable 5- to 7- membered monocyclic or bicyclic or 7- to 10-membered bicyclic heterocyclic ring which is saturated partially unsaturated or unsaturated (aromatic), and which consists of carbon atoms and from 1 to 4 heteroatoms independently selected from the group consisting of N, O and S and including any bicyclic group in which any of the above-defined heterocyclic rings is fused to a benzene ring. The nitrogen and sulfur heteroatoms may optionally be oxidized. The heterocyclic ring may be attached to its pendant group at any heteroatom or carbon atom which results in a stable structure. The heterocyclic rings described herein may be substituted on carbon or on a nitrogen atom if the resulting compound is stable. If specifically

noted, a nitrogen in the heterocycle may optionally be quaternized. It is preferred that when the total number of S and O atoms in the heterocycle exceeds 1, then these heteroatoms are not adjacent to one another. It is preferred that the total number of S and O atoms in the heterocycle is not more than 1. As used herein, the term "aromatic heterocyclic system" is intended to mean a stable 5- to 7-membered monocyclic or bicyclic or 7- to 10-membered bicyclic heterocyclic aromatic ring which consists of carbon atoms and from 1 to 4 heteroatoms independently selected from the group consisting of N, O and S. It is preferred that the total number of S and O atoms in the aromatic heterocycle is not more than 1.

Examples of heterocycles include, but are not limited to, 1H-indazole, 2-pyrrolidinyl, 2H,6H-1,5,2-dithiazinyl, 2H-pyrrolyl, 3H-indolyl, 4-piperidinyl, 4aH-carbazole, 4H-quinolizinyl, 6H-1,2,5-thiadiazinyl, acridinyl, azocinyl, benzimidazolyl, benzofuranyl, benzothiofuranyl, benzothiophenyl, benzoxazolyl, benzthiazolyl, benztriazolyl, benztetrazolyl, benzisoxazolyl, benzisothiazolyl, benzimidazolyl, carbazolyl, 4aH-carbazolyl, β -carboline, chromanyl, chromenyl, cinnolinyl, decahydroquinolinyl, 2H,6H-1,5,2-dithiazinyl, dihydrofuro[2,3-b]tetrahydrofuran, furanyl, furazanyl, imidazolidinyl, imidazolinyl, imidazolyl, 1H-indazolyl, indolenyl, indolinyl, indolizinyl, indolyl, isobenzofuranyl, isochromanyl, isoindazolyl, isoindolinyl, isoindolyl, isoquinolinyl, isothiazolyl, isoxazolyl, morpholinyl, naphthyridinyl, octahydroisoquinolinyl, oxadiazolyl, 1,2,3-oxadiazolyl, 1,2,4-oxadiazolyl, 1,2,5-oxadiazolyl, 1,3,4-oxadiazolyl, oxazolidinyl, oxazolyl, oxazolidinylperimidinyl, phenanthridinyl, phenanthrolinyl, phenarsazinyl, phenazinyl, phenothiazinyl, phenoxathiinyl, phenoxazinyl, phthalazinyl, piperazinyl, piperidinyl, pteridinyl, piperidinyl, 4-piperidinyl, pteridinyl, purinyl, pyran, pyrazinyl, pyrazolidinyl, pyrazolinyl, pyrazolyl, pyridazinyl, pyridoaxazole, pyridoimidazole, pyridothiazole, pyridinyl, pyridyl, pyrimidinyl, pyrrolidinyl, pyrrolinyl, pyrrolyl, quinazolinyl, quinolinyl, 4H-quinolizinyl,

quinoxaliny1, quinuclidiny1, carboliny1, tetrahydrofurany1,
tetrahydroisoquinoliny1, tetrahydroquinoliny1, 6H-1,2,5-
thiadiaziny1, 1,2,3-thiadiazoly1, 1,2,4-thiadiazoly1, 1,2,5-
thiadiazoly1, 1,3,4-thiadiazoly1, thianthreny1, thiazoly1,
5 thieny1, thienothiazoly1, thienooxazoly1, thienoimidazoly1,
thiopheny1, triaziny1, 1,2,3-triazoly1, 1,2,4-triazoly1,
1,2,5-triazoly1, 1,3,4-triazoly1, xantheny1. Preferred
heterocycles include, but are not limited to, pyridiny1,
furany1, thieny1, pyrroly1, pyrazoly1, imidazoly1, indoly1,
10 benzimidazoly1, 1H-indazoly1, oxazolidiny1, benzotriazoly1,
benzisoxazoly1, oxindoly1, benzoxazolinyl, or isatinoy1. Also
included are fused ring and spiro compounds containing, for
example, the above heterocycles.

The phrase "pharmaceutically acceptable" is employed
15 herein to refer to those compounds, materials, compositions,
and/or dosage forms which are, within the scope of sound
medical judgment, suitable for use in contact with the tissues
of human beings and animals without excessive toxicity,
irritation, allergic response, or other problem or
20 complication, commensurate with a reasonable benefit/risk
ratio.

As used herein, "pharmaceutically acceptable salts" refer
to derivatives of the disclosed compounds wherein the parent
compound is modified by making acid or base salts thereof.
25 Examples of pharmaceutically acceptable salts include, but are
not limited to, mineral or organic acid salts of basic
residues such as amines; alkali or organic salts of acidic
residues such as carboxylic acids; and the like. The
pharmaceutically acceptable salts include the conventional
30 non-toxic salts or the quaternary ammonium salts of the parent
compound formed, for example, from non-toxic inorganic or
organic acids. For example, such conventional non-toxic salts
include those derived from inorganic acids such as
hydrochloric, hydrobromic, sulfuric, sulfamic, phosphoric,
35 nitric and the like; and the salts prepared from organic acids
such as acetic, propionic, succinic, glycolic, stearic,
lactic, malic, tartaric, citric, ascorbic, pamoic, maleic,
hydroxymaleic, phenylacetic, glutamic, benzoic, salicylic,

sulfanilic, 2-acetoxybenzoic, fumaric, toluenesulfonic, methanesulfonic, ethane disulfonic, oxalic, isethionic, and the like.

5 The pharmaceutically acceptable salts of the present invention can be synthesized from the parent compound which contains a basic or acidic moiety by conventional chemical methods. Generally, such salts can be prepared by reacting the free acid or base forms of these compounds with a stoichiometric amount of the appropriate base or acid in water
10 or in an organic solvent, or in a mixture of the two; generally, nonaqueous media like ether, ethyl acetate, ethanol, isopropanol, or acetonitrile are preferred. Lists of suitable salts are found in *Remington's Pharmaceutical Sciences*, 17th ed., Mack Publishing Company, Easton, PA, 1985,
15 p. 1418, the disclosure of which is hereby incorporated by reference.

"Prodrugs" are intended to include any covalently bonded carriers which release the active parent drug according to formula (I) *in vivo* when such prodrug is administered to a
20 mammalian subject. Prodrugs of a compound of formula (I) are prepared by modifying functional groups present in the compound in such a way that the modifications are cleaved, either in routine manipulation or *in vivo*, to the parent compound. Prodrugs include compounds of formula (I) wherein a
25 hydroxy, amino, or sulfhydryl group is bonded to any group that, when the prodrug or compound of formula (I) is administered to a mammalian subject, cleaves to form a free hydroxyl, free amino, or free sulfhydryl group, respectively. Examples of prodrugs include, but are not limited to, acetate,
30 formate and benzoate derivatives of alcohol and amine functional groups in the compounds of formula (I), and the like. Preferred prodrugs are amidine prodrugs wherein D is $C(=NR^7)NH_2$, and R^7 is selected from OH, C_{1-4} alkoxy, C_{6-10} aryloxy, C_{1-4} alkoxycarbonyl, C_{6-10} aryloxycarbonyl, C_{6-10}
35 arylmethylcarbonyl, C_{1-4} alkylcarbonyloxy C_{1-4} alkoxycarbonyl, and C_{6-10} arylcarbonyloxy C_{1-4} alkoxycarbonyl. More preferred prodrugs are where R^7 is OH, methoxy, ethoxy,

benzyloxycarbonyl, methoxycarbonyl, and methylcarbonyloxymethoxycarbonyl.

"Stable compound" and "stable structure" are meant to indicate a compound that is sufficiently robust to survive
5 isolation to a useful degree of purity from a reaction mixture, and formulation into an efficacious therapeutic agent.


SYNTHESIS

10 The compounds of the present invention can be prepared in a number of ways known to one skilled in the art of organic synthesis. The compounds of the present invention can be synthesized using the methods described below, together with
15 synthetic methods known in the art of synthetic organic chemistry, or by variations thereon as appreciated by those skilled in the art. Preferred methods include, but are not limited to, those described below. The reactions are performed in a solvent appropriate to the reagents and materials employed and suitable for the transformations being
20 effected. It will be understood by those skilled in the art of organic synthesis that the functionality present on the molecule should be consistent with the transformations proposed. This will sometimes require a judgment to modify the order of the synthetic steps or to select one particular
25 process scheme over another in order to obtain a desired compound of the invention. It will also be recognized that another major consideration in the planning of any synthetic route in this field is the judicious choice of the protecting group used for protection of the reactive functional groups
30 present in the compounds described in this invention. An authoritative account describing the many alternatives to the trained practitioner is Greene and Wuts (*Protective Groups In Organic Synthesis*, Wiley and Sons, 1991). All references cited herein are hereby incorporated in their entirety herein
35 by reference.

Schemes 1-4 describe the synthesis of compounds wherein M is furan and Q is a protected precursor of group D of formula I. Each scheme represents a different substitution pattern

for the furan ring. In Scheme 1, an alpha-substituted carboxylic acid, wherein V is a nitro, protected sulfonamide or ester group, can be treated with two equivalents of base to activate it, quenched with an appropriate aldehyde electrophile as described by Wierenga (*J. Org. Chem.*, 44(2), 310, 1979) and then oxidized by pyridinium dichromate to a ketone. Treatment with base and acetic anhydride should give the enol acetate which can react with a vinyl sulfoxide to give a dihydrofuran as shown by Chan (*J. Chem. Soc., Perkins Trans. 1* **1992**, 945). This sulfoxide can then be oxidized and eliminated to give the desired furan.

Scheme 1




15

In Scheme 2, the readily available bromides Q-E-Br are coupled to a terminal acetylene, to give a disubstituted alkyne as shown by Padwa (*J. Org. Chem.* **1991**, 56(7), 2523). Also shown in Scheme 2, a carboxylic acid can be homologated into a ketone and then converted into a diazoketone. Rhodium catalyzed cyclization can provide the desired furan as in Davies (*Tetrahedron* **1988**, 44(11), 3343).


25

Scheme 2




Addition of a grignard reagent to the appropriate aldehyde, oxidation and O-methylation should give the required enol ether as shown in Scheme 3. Diazoketone formation of the acetyl derivative, AcW, and copper catalyzed cyclization can be done like Alonos (*J. Org. Chem.* **1991**, 56(7), 2523) followed by standard deprotection should give the desired furan.

Scheme 3




Scheme 4 describes a synthesis of a different substitution pattern on a furan ring. The carboxylic acid from above can be converted into a ketone in a two-step process by conversion to the activated acid chloride and reacting with a cuprate (*Tetr. Lett.* **1971**, 829). Piperidine-catalyzed condensation with an appropriate aldehyde should give the unsaturated ketone as shown by Taylor (*J. Het. Chem.* **1989**, 26, 1353). Conjugate addition of a dithiane to the unsaturated ketone should give the required substitution pattern. N-bromosuccinimide deprotection of the dithiane followed by acid-catalyzed cyclization can provide the furan.

Scheme 4



- 5 Schemes 5 and 6 describe the synthesis of compounds wherein ring M is thiophene. The appropriate aldehyde in Scheme 5 can be oxidized to a carboxylic acid and converted to an acid chloride. Reaction of this acid chloride with methyl ketone and lithium bis(trimethylsilyl)amide as shown by
- 10 Cushman (*Tetr. Lett.* **1990**, 45, 6497). Further treatment with diazomethane can provide a mixture of two regioisomers which need not be separated at this time. Treatment of the commercially available bromide with sodium sulfide followed by the unsaturated ketone should give a mixture of thiophene
- 15 regioisomers which can be separated according to Alberola (*Synth. Comm.* **1990**, 20, 2537).

Scheme 5



20


Alternatively, in Scheme 6, ethyl acetate can be diazotized by tosyl azide and carbene insertion into the E-Br bond as in D'yakonov (*J. Gen. Chem. USSR*, **1951**, 21, 851).

Nucleophilic displacement with a thiocarboxylic acid (*Org. Syn. Coll.* **1963**, 4, 924) should give the appropriate carboxylic acid after basic hydrolysis as shown by Masuda (*Chem. Pharm. Bull.* **1977**, 25, 1471). Reaction with a

- 5 disubstituted alkyne with trifluoroacetic anhydride can give a mixture of regioisomers. By analogy switching the position of V and Q-E in the reagents can give a different set of regioisomers.

10

Scheme 6




- Schemes 7 and 8 provide routes to compounds of Formula I
 15 wherein ring M is isoxazole. Scheme 7 shows one possible
 synthesis of isoxazoles. Substituted benzaldehydes can be
 reacted with hydroxyl amine then chlorinated to give the
 hydroximinoyl chloride (see *J. Org. Chem.* **1980**, 45, 3916).
 Preparation of the nitrile oxide in situ with triethylamine
 20 and cycloaddition with a substituted alkyne can give a mixture
 of regioisomeric isoxazoles as shown by H. Kawakami (*Chem.*
Lett. **1987**, 1, 85). Preparation of the disubstituted alkyne
 can be achieved by nucleophilic attack of the alkynyl anion on
 an electrophile as shown by L. N. Jungheim (*J. Org. Chem.*
 25 **1987**, 57, 4007). Alternatively, one could make the

hydroxyiminoyl chloride of the R^{1a} piece and react it with an appropriately substituted alkyne to give another set of regioisomeric isoxazoles which can be separated chromatographically.

5

Scheme 7




10 An alternate procedure which should produce only one isoxazole regioisomer is described in Scheme 8. The methylated form of substituent V can be deprotonated and silylated. Chlorination with carbon tetrachloride or fluorination with difluorodibromomethane under triethylborane
15 catalysis can give the geminal dihalo compound as shown by Sugimoto (*Chem. Lett.* **1991**, 1319). Cuprate-mediated conjugate addition-elimination give the desired alkene as in Harding (*J. Org. Chem.* **1978**, 43, 3874).


Alternatively, one can acylate with an acid chloride to
20 form a ketone as in Andrews (*Tetr. Lett.* **1991**, 7731) followed by diazomethane to form the enol ether. Each of these compounds can be reacted with a hydroximinoyl chloride in the presence of triethylamine to give one regioisomeric isoxazole as shown by Stevens (*Tetr. Lett.* **1984**, 4587).

25

Scheme 8



- 5 The following is a reaction grid for the synthesis of the Z linkage. The following coupling reaction would be readily known to those skilled in the art of organic synthesis.



When Z =


- 10 **CONH** then use V = CO₂CH₃ and W = NH₂ under AlMe₃ catalysis
- SO₂NH** then make heterocycle after sulfonamide formation
- CH₂NH** then reduce V = CO₂CH₃ with DIBAL to CH₂OH and couple with W = NH₂ using PPh₃
- 15 **CH₂S** then reduce V = CO₂CH₃ with DIBAL to CH₂OH and couple with W = SH using MsCl
- CH₂O** then reduce V = CO₂CH₃ with DIBAL to CH₂OH and couple with W = OH using PPh₃
- 20 **NHCO** then reduce V = NO₂ to NH₂ using H₂/Pd and couple with W = CO₂CH₃ using AlMe₃
- NHSO₂** then reduce V = NO₂ to NH₂ using H₂/Pd and couple with W = SO₂Cl
- NHCH₂** then reduce V = NO₂ to NH₂ using H₂/Pd and couple with W = CH₂Br
- 25 **OCH₂** then reduce, then diazotize V = NO₂ and couple with W = CH₂Br
- SCH₂** then reduce V = SO₂NR₂ with LAH and couple with W = CH₂Br.

To complete the final reaction sequence, substituent Q can be deprotected or reacted to give an amine or amide. The

amine can be converted into an amidine, guanidine or formamidine under standard conditions as outlined in Scheme 9. From the nitrile, imidate formation followed by amination with ammonium carbonate can provide the amidine.

5

Scheme 9




10 The compounds of Formula I in which ring M is thiazole or
 15 oxazole can be prepared as outlined in Schemes 10-16 wherein,
 R^e and R^f can be Z-A-B or R^{1a} or precursors thereof. There are
 numerous methods by which to prepare and manipulate
 substituted thiazole and oxazole rings (for reviews, see
 20 *Comprehensive Heterocyclic Chemistry*, Katritzky and Rees, eds.
1984, 6, 247 and *Chem. Het. Compds.* **1979**, 34-2, 1). One
 particularly useful method for preparing thiazole and oxazole
 containing compounds of the present invention is the Hantzsch
 method, which involves condensation of α -haloketones with
 25 thioamides, thioureas, amides and ureas.

As shown in Scheme 10, an appropriate ketone can be
 brominated by a variety of electrophilic brominating reagents
 such as pyridinium bromide perbromide, NBS, etc. to afford an
 α -bromoketone. Heating with a wide variety of substituted
 25 thioamides and thioureas, and amides and ureas can afford
 thiazole and oxazole derivatives. Regioisomeric thiazoles and
 oxazoles can be prepared by a similar reaction sequence
 beginning with a similar ketone. The ketones in Scheme 10 are
 readily available by procedures familiar to those skilled in
 30 the art of organic synthesis. The functionality Q can later

be transformed into the group D found in compounds of Formula I.


The thioamides are either commercially available or can be prepared from the corresponding amides using Lawesson's reagent or phosphorous pentasulfide. They can also be prepared and cyclized *in situ* by performing the cyclization reaction with the corresponding amide in the presence of phosphorous pentasulfide. The thioureas are either commercially available or are readily prepared from other commercially available thioureas. The amides and ureas are either commercially available or are readily prepared by procedures known to those skilled in the art.

Scheme 10



In Scheme 11 is shown how α -acylaminoketones can be converted into oxazoles by dehydration with an acid such as sulfuric acid. Treating with phosphorous pentasulfide can afford thiazoles. The starting materials are prepared by standard methods known to those skilled in the art.


Scheme 11



Oxazoles can also be prepared by the cyclization strategy shown in Scheme 12. Ketones can be converted into their oxime derivatives by standard treatment with hydroxylamine.


- 5 Treating these intermediates with acid chlorides can provide the corresponding oxazoles.

Scheme 12



- 2-Unsubstituted oxazoles can be prepared by the cyclization shown in Scheme 13. Treatment of acid chlorides with an isocyanoacetate (wherein R^g can be A-B or a precursor thereof) in the presence of a base such as triethylamine can afford the oxazoles (Suzuki et. al. *Syn. Comm.* **1972**, 2, 237).


Scheme 13



- Other cyclization strategies can afford differently substituted thiazoles and oxazoles. In Scheme 14 is shown how cyclizations can be modified to afford 5-aminooxazoles and 4- and 5-aminothiazoles. Treatment of aldehydes with NaCN and ammonium chloride can afford α -aminonitriles (Steiger *Org. Syn. Coll. Vol. III* **1955**, 84). Acylation with acid chlorides followed by acid-catalyzed dehydration can afford 5-


aminoxazoles. The bromination of nitriles with bromine can afford α -bromonitriles. These can be treated with a variety of thioamides to afford 4-aminothiazoles. The 5-aminothiazoles can be prepared by elaboration of thiazole carboxylic acids. Formation of the acyl azide by standard methods can be followed by heating to effect a Curtius rearrangement to give the isocyanate (South, *J. Het. Chem.* **1991**, 28, 1003). Addition of water can then afford the 5-aminothiazoles.

Scheme 14



In Scheme 15 is shown how thiazoles and oxazoles with halogen substituents can be prepared. The 2-halo- derivatives can be prepared from the corresponding amino derivatives by diazotization with nitrous acid or isoamyl nitrite followed by displacement with an appropriate halide source such as copper bromide or chloride. The 5-halo- derivatives can be prepared by ring bromination with NBS or Br_2 , or chlorination with NCS or Cl_2 . Alternatively, the Hunsdiecker procedure (Ber. **1942**, 75, 291) can be applied to the 5-carboxylic acid derivatives to prepare the bromides. The 4-halo derivatives can be prepared in the same manner from the regioisomer in which the group Q-E occupies position 5 on the ring.


Scheme 15



5


In Scheme 16 is shown how mercapto and sulfonyl derivatives of the thiazoles and oxazoles can be prepared. The 2-mercapto derivatives can be prepared from the corresponding 2-amino heterocycles by diazotization with nitrous acid or isoamyl nitrite followed by reaction with an appropriate thiol. Oxidation of the thiol derivative can afford the sulfonic acid derivatives. The 5-mercapto derivatives can be prepared by thiol displacement of the appropriate 5-bromo derivative. Alternatively, halogen metal exchange of the bromo derivative with *n*-BuLi followed by quenching with sulfur can afford the required 5-mercapto derivatives. The sulfonyl derivatives are available by oxidation of the mercapto derivatives. In some cases direct sulfonation of the thiazole or oxazole ring can be possible. When R' is an activating group such as amino or alkoxy, treatment with chlorosulfonic acid should give the sulfonyl derivative (Mann et. al. *J. Prakt. Chem.* **1978**, 320, 715). The 4-mercapto and sulfonyl derivatives can be prepared in the same manner as shown for the 5-derivatives from the regioisomers in which the group Q-E occupies position 5 on the ring.

Scheme 16




- 5 By the cyclization strategies described in Schemes 10-16, and by other strategies not described but familiar to those skilled in the art of organic synthesis, a wide variety of highly substituted thiazoles and oxazoles can be prepared. Proper manipulation of the starting materials for these
- 10 cyclizations by procedures known to those skilled in the art also allows for the synthesis of oxazoles (Scheme 17, $\text{J} = \text{O}$) and thiazoles (Scheme 17, $\text{J} = \text{S}$), which are regioisomers of the thiazoles and oxazoles of Scheme 10, containing a wide variety of substituents R^e and R^f which by proper manipulation
- 15 described in preceeding and following schemes can be converted into R^{1a} and Z-A-B of compounds of Formula I.

Scheme 17




The present invention also describes compounds of Formula I in which ring M is 1,2,3- and 1,2,5-thiadiazole and 1,2,5-oxadiazole. The following schemes provide methods for synthesizing these heterocycles. In Scheme 18 is shown how 1,2,3-thiadiazoles can be prepared. The ketones from Scheme 10 can be converted by standard procedures into semicarbazones ($R^f = \text{NH}_2$) or acylhydrazones ($R^f = \text{alkyl, alkoxy}$) which can then be treated with thionyl chloride to prepare the 1,2,3-thiadiazoles (*J. Med. Chem.* **1985**, 28, 442). Alternatively, diazo ketones can be prepared by treatment with base and a suitable diazo transfer reagent such as tosyl azide. Treatment of these diazo intermediates with hydrogen sulfide or Lawesson's reagent can afford the 1,2,3-thiadiazoles.

Scheme 18




In Scheme 19 is shown how to prepare the 1,2,5-thiadiazoles contained in compounds of Formula I. The disubstituted alkynes, which are readily available by standard alkyne coupling procedures known to those skilled in the art of organic synthesis, can be treated with sulfur nitride in refluxing toluene to afford the 1,2,5-thiadiazoles (*J. Het. Chem.* **1979**, *16*, 1009).

Scheme 19



In Scheme 20 is shown how 1,2,5-oxadiazole heterocycles can be prepared. Diazotization of ketones followed by treatment with hydroxylamine can afford the bisoximes. Alternatively, diketones can be treated with hydroxylamine to afford the bisoximes. Dehydration of these readily prepared intermediates with acetic acid or thionyl chloride can then afford the 1,2,5-oxadiazoles.

Scheme 20




In the cyclization sequences and strategies described above, in general the substituents Q-E and R^e and R^f can be varied widely. In some cases R^e can be chosen so that it corresponds to Z-A-B in Formula I. In other cases R^b can be chosen so that it is hydrogen, carboxylic ester, amino, alkyl, cyano, alkoxy, hydroxy, thioalkoxy, sulfonyl, etc. which can subsequently be converted into the group Z-A-B of Formula I.

In the following schemes are described some methods by which the various groups Z of Formula I can be prepared from various groups R^e. In these schemes the heterocycle is denoted as ring M and it is understood that the reactions described will generally be applicable to all of the different heterocycles previously described. It is also understood that the reactions described may require some modification of the reaction conditions, change in the reaction order or suitable protecting groups, depending upon the functionality contained in the compound of interest. One skilled in the art of organic synthesis will understand this and be able to modify the reaction sequence to obtain the desired products.

In Scheme 21 is shown how the heterocyclic compounds from above where R^e is a carboxylic ester group can be converted into compounds containing the Z-A-B residue. For the amide linker (Formula I, Z = -CONH-) ring M where R^e = carboalkoxy can be hydrolyzed to the acid. Formation of the acid chloride with thionyl chloride followed by the addition of an appropriate amine H₂N-A-B can afford the amide-linked compounds. Alternatively, the acid can be combined with the amine H₂N-A-B in the presence of a suitable peptide coupling agent, such as BOP-Cl, HBTU or DCC to afford the corresponding amides. In another method the ester can be directly coupled with an aluminum reagent, prepared by the addition of trimethylaluminum to the amine H₂N-A-B, to afford the amide. To form ether- and thioether-linked compounds of Formula I (Z = -CH₂O-, -CH₂S-) the acid can be reduced to the alcohol. Preferred procedures for this transformation are reduction with borane THF complex, or a procedure involving the reduction of the mixed anhydride of the acid with sodium borohydride. Completion of the ether and thioether linked compounds of Formula I can be readily accomplished by the Mitsunobu protocol with an appropriate phenol, thiophenol or hydroxy- or mercaptoheterocycle HZ-A-B (Formula I, A = aryl or heteroaryl). Other ethers or thioethers can be prepared following initial conversion of the alcohol to a suitable leaving group, such as tosylate. Where J = S, thioethers can be further oxidized to prepare the sulfones (Formula I, Z = -

CH₂SO₂-). To prepare the amine-linked compounds of Formula I (Z = -CH₂NH-) the alcohol can be oxidized to the aldehyde by a number of procedures, two preferred methods of which are the Swern oxidation and oxidation with pyridinium chlorochromate (PCC). Reductive amination of aldehyde with an appropriate amine H₂N-A-B and sodium cyanoborohydride can then afford the amine linked compounds. The aldehyde also can be used to prepare the ketone-linked compounds of Formula I (Z = -COCH₂-). Treatment of the aldehyde with an organometallic species can afford the alcohol. The organo metallic species (where M = magnesium or zinc) can be best prepared from the corresponding halide by treatment with metallic magnesium or zinc. These reagents readily react with aldehydes to afford alcohols. Oxidation of the resulting alcohol by any of a number of procedures, such as the Swern oxidation or PCC oxidation, can afford the ketone.

Scheme 21




- 5 Additional compounds of Formula I in which the linking group Z contains a nitrogen atom attached to ring M can be prepared by the procedures described in Scheme 22. The amines can be converted to the sulfonamides (Formula I, $Z = -NHSO_2-$) by treatment with an appropriate sulfonyl chloride $B-A-SO_2Cl$ in the presence of a base such as triethylamine. The amines can be converted into the amides (Formula I, $Z = -NHCO-$) by treatment with an appropriate acid chloride $Cl-CO-A-B$ in the presence of a base or by treatment with an appropriate carboxylic acid $HO-CO-A-B$ in the presence of a suitable
- 10

peptide coupling agent, such as DCC, HBTU or BOP-Cl. The amine can be converted into amines of Formula I ($Z = -NHCH_2-$) by reductive amination with an appropriate aldehyde $OHC-A-B$.

5


Scheme 22



Additional compounds of Formula I in which the linking group Z contains a sulfur atom attached to ring M can be prepared by the procedures described in Scheme 23. Treatment of sulfonyls with phosphorous pentachloride followed by treatment with an appropriate amine H_2N-A-B can afford the sulfonamide-linked compounds (Formula I, $Z = -SO_2NH-$). The thiols can be alkylated with a suitable alkylating reagent in the presence of a base to afford thioethers (Formula I, $Z = -SCH_2-$). These compounds can be further oxidized by a variety of reagents to afford the sulfone-linked compounds (Formula I, $Z = -SO_2CH_2-$).


20

Scheme 23



- 5 Compounds of this invention where B is either a carbocyclic or heterocyclic residue as defined in Formula I are coupled to A as shown generically and by specific example in Scheme 24, either or both of A and B may be substituted with 0-2 R⁴. W is defined as a suitable protected nitrogen, such as NO₂ or NHBOC; a protected sulfur, such as S-tBu or SMOM; or a methyl ester. Halogen-metal exchange of the bromine in bromo-B with n-butyl lithium, quenching with triisopropyl borate and acidic hydrolysis should give the required boronic acid, B'-B(OH)₂. The W-A-Br subunit may be already linked to ring M before the Suzuki coupling reaction. Deprotection can provide the complete subunit.
- 10
- 15

Scheme 24




20

- Scheme 25 describes a typical example of how the A-B subunit can be prepared for attachment to ring M. 4-Bromoaniline can be protected as Boc-derivative and the coupled to 2-(t-butylamino)sulfonylphenylboronic acid under
- 25

Suzuki conditions. 2-(t-Butylamino)sulfonylphenylboronic acid can be prepared by the method described by Rivero (*Bioorg. Med. Chem. Lett.* **1994**, 189). Deprotection with TFA can provide the aminobiphenyl compound. The aminobiphenyl can then be coupled to the core ring structures as described below.

Scheme 25



When B is defined as X-Y, the following description applies. Groups A and B are available either through commercial sources, known in the literature or readily synthesized by the adaptation of standard procedures known to practioners skilled in the art of organic synthesis. The required reactive functional groups appended to analogs of A and B are also available either through commercial sources, known in the literature or readily synthesized by the adaptation of standard procedures known to practioners skilled in the art of organic synthesis. In the tables that follow the chemistry required to effect the coupling of A to B is outlined.

Table A: Preparation of Amide, Ester, Urea, Sulfonamide and Sulfamide linkages between A and B.

Rxn. No.	if A contains :	then the reactive substituent of Y is :	to give the following product A-X-Y :
1	A-NHR ² as a substituent	ClC(O)-Y	A-NR ² -C(O)-Y

2	a secondary NH as part of a ring or chain	ClC(O)-Y	A-C(O)-Y
3	A-OH as a substituent	ClC(O)-Y	A-O-C(O)-Y
4	A-NHR ² as a substituent	ClC(O)-CR ² R ^{2a} -Y	A-NR ² -C(O)-CR ² R ^{2a} -Y
5	a secondary NH as part of a ring or chain	ClC(O)-CR ² R ^{2a} -Y	A-C(O)-CR ² R ^{2a} -Y
6	A-OH as a substituent	ClC(O)-CR ² R ^{2a} -Y	A-O-C(O)-CR ² R ^{2a} -Y
7	A-NHR ³ as a substituent	ClC(O)NR ² -Y	A-NR ² -C(O)NR ² -Y
8	a secondary NH as part of a ring or chain	ClC(O)NR ² -Y	A-C(O)NR ² -Y
9	A-OH as a substituent	ClC(O)NR ² -Y	A-O-C(O)NR ² -Y
10	A-NHR ² as a substituent	ClSO ₂ -Y	A-NR ² -SO ₂ -Y
11	a secondary NH as part of a ring or chain	ClSO ₂ -Y	A-SO ₂ -Y
12	A-NHR ² as a substituent	ClSO ₂ -CR ² R ^{2a} -Y	A-NR ² -SO ₂ -CR ² R ^{2a} -Y
13	a secondary NH as part of a ring or chain	ClSO ₂ -CR ² R ^{2a} -Y	A-SO ₂ -CR ² R ^{2a} -Y
14	A-NHR ² as a substituent	ClSO ₂ -NR ² -Y	A-NR ² -SO ₂ -NR ² -Y
15	a secondary NH as part of a ring or chain	ClSO ₂ -NR ² -Y	A-SO ₂ -NR ² -Y
16	A-C(O)Cl	HO-Y as a substituent	A-C(O)-O-Y

17	A-C(O)Cl	NHR ² -Y as a substituent	A-C(O)-NR ² -Y
18	A-C(O)Cl	a secondary NH as part of a ring or chain	A-C(O)-Y
19	A-CR ² R ^{2a} C(O)Cl	HO-Y as a substituent	A-CR ² R ^{2a} C(O)-O-Y
20	A-CR ² R ^{2a} C(O)Cl	NHR ² -Y as a substituent	A-CR ² R ^{2a} C(O)-NR ² -Y
21	A-CR ² R ^{2a} C(O)Cl	a secondary NH as part of a ring or chain	A-CR ² R ^{2a} C(O)-Y
22	A-SO ₂ Cl	NHR ² -Y as a substituent	A-SO ₂ -NR ² -Y
23	A-SO ₂ Cl	a secondary NH as part of a ring or chain	A-SO ₂ -Y
24	A-CR ² R ^{2a} SO ₂ Cl	NHR ² -Y as a substituent	A-CR ² R ^{2a} SO ₂ -NR ² -Y
25	A-CR ² R ^{2a} SO ₂ Cl	a secondary NH as part of a ring or chain	A-CR ² R ^{2a} SO ₂ -Y

The chemistry of Table A can be carried out in aprotic solvents such as a chlorocarbon, pyridine, benzene or toluene, at temperatures ranging from -20°C to the reflux point of the solvent and with or without a trialkylamine base.

Table B: Preparation of ketone linkages between A and B.

Rxn. No.	if A contains :	then the reactive substituent of Y is :	to give the following product A-X-Y :
1	A-C(O)Cl	BrMg-Y	A-C(O)-Y
2	A-CR ² R ^{2a} C(O)Cl	BrMg-Y	A-CR ² R ^{2a} ₂ C(O)-Y
3	A-C(O)Cl	BrMgCR ² R ^{2a} -Y	A-C(O)CR ² R ^{2a} -Y
4	A-CR ² R ^{2a} C(O)Cl	BrMgCR ² R ^{2a} -Y	A-CR ² R ^{2a} C(O)CR ² R ^{2a} -Y

The coupling chemistry of Table B can be carried out by a variety of methods. The Grignard reagent required for Y is prepared from a halogen analog of Y in dry ether, dimethoxyethane or tetrahydrofuran at 0°C to the reflux point of the solvent. This Grignard reagent can be reacted directly under very controlled conditions, that is low temperature (-20°C or lower) and with a large excess of acid chloride or with catalytic or stoichiometric copper bromide·dimethyl sulfide complex in dimethyl sulfide as a solvent or with a variant thereof. Other methods available include transforming the Grignard reagent to the cadmium reagent and coupling according to the procedure of Carson and Prout (Org. Syn. Col. Vol. 3 (1955) 601) or a coupling mediated by Fe(acac)₃ according to Fiandanese et al (Tetrahedron Lett., (1984) 4805), or a coupling mediated by manganese (II) catalysis (Cahiez and Laboue, Tetrahedron Lett., 33(31), (1992) 4437).

Table C: Preparation of ether and thioether linkages between A and B

Rxn. No.	if A contains :	then the reactive substituent of Y is :	to give the following product A-X-Y :
1	A-OH	Br-Y	A-O-Y
2	A-CR ² R ^{2a} -OH	Br-Y	A-CR ² R ^{2a} O-Y
3	A-OH	Br-CR ² R ^{2a} -Y	A-OCR ² R ^{2a} -Y
4	A-SH	Br-Y	A-S-Y
5	A-CR ² R ^{2a} -SH	Br-Y	A-CR ² R ^{2a} S-Y
6	A-SH	Br-CR ² R ^{2a} -Y	A-SCR ² R ^{2a} -Y

The ether and thioether linkages of Table C can be prepared by reacting the two components in a polar aprotic solvent such as acetone, dimethylformamide or dimethylsulfoxide in the presence of a base such as potassium carbonate, sodium hydride or potassium t-butoxide at temperature ranging from ambient temperature to the reflux point of the solvent used.

Table D: Preparation of -SO- and -SO₂- linkages from thioethers of Table 3.

Rxn. No.	if the starting material is :	and it is oxidized with Alumina (wet)/ Oxone (Greenhalgh, Synlett, (1992) 235) the product is :	and it is oxidized with m-chloroperbenzoic acid (Sato et al., Chem. Lett. (1992) 381), the product is :
1	A-S-Y	A-S(O)-Y	A-SO ₂ -Y
2	A-CR ² R ^{2a} S-Y	A-CR ² R ^{2a} S(O)-Y	A-CR ² R ^{2a} SO ₂ -Y
3	A-SCR ² R ^{2a} -Y	A-S(O)CR ² R ^{2a} -Y	A-SO ₂ CR ² R ^{2a} -Y

The thioethers of Table C serve as a convenient starting material for the preparation of the sulfoxide and sulfone analogs of Table D. A combination of wet alumina and oxone can provide a reliable reagent for the oxidation of the

thioether to the sulfoxide while m-chloroperbenzoic acid oxidation will give the sulfone.

Other features of the invention will become apparent in the course of the following descriptions of exemplary embodiments which are given for illustration for the invention and are not intended to be limiting thereof.

EXAMPLES

10

Example 1

3-(3-amidinophenyl)-4-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-5-(hydroxymethyl)isoxazole, trifluoroacetic acid salt

15 Part A. Preparation of 2-(t-butylaminosulfonyl)phenylboronic acid.

To a solution of 206.5 g (0.968 mol) of benzene-(N-t-butyl)sulfonamide in 2500 mL of THF under N₂ was added 790 mL (1.98 mol) of 2.5M n-butyllithium in hexane over 35 minutes, keeping the temperature between 0-5°C. The reaction mixture was allowed to warm to 10°C, at which time a thick precipitate formed. Triisopropylborate (305 mL, 1.32 mol) was added keeping the temperature below 35°C. After 1 hour, the reaction mixture was cooled, 1N HCl (1570 mL) was added, and the mixture was stirred overnight. The mixture was extracted with 400 mL of ether three times, and the combined organic extracts were extracted with 500 mL of 1N NaOH three times. The aqueous extracts were acidified to pH 1 with 6N HCl, and then extracted with 500 mL ether three times. The combined ether extracts were dried over MgSO₄, and the solvents evaporated in vacuo until the volume was 700 mL. Hexane (150 mL) was added and overnight, a white precipitate formed. The solid was collected and washed with 10% ether/hexane (250 mL), then dried in vacuo to give 216.3 g (87%) of the desired compound as white crystals. m.p. 118-119°C ¹H NMR (CDCl₃) δ: 8.00 (d, 1H); 7.82 (d, 1H); 7.53 (m, 2H); 6.29 (br s, 2H); 5.13 (s, 1H); 1.18 (s, 9H).

Part B. Preparation of N-(4-bromophenyl)-4-(tetrahydropyran-2-yloxymethyl)-2-butyramide.

5 To a solution of 4.98 g (35.5 mmol) of tetrahydro-2-(2-propynyloxy)-2H-pyran in 70 mL of THF under N₂ was added 14.2 mL (35.5 mmol) of 2.5 M n-butyllithium in hexane. After 15 minutes, 7.03 g (35.5 mmol) of 4-bromophenylisocyanate was added and then the reaction was allowed to warm to room
10 temperature. Saturated aqueous ammonium chloride (20 mL) was added and the mixture extracted with 30 mL ethyl acetate three times. The combined organic extracts were dried with MgSO₄, concentrated to an oil *in vacuo* and then chromatographed on silica with 20% EtOAc/hexane to give 7.1 g (59%) of the
15 desired alkyne. ¹H NMR (CDCl₃) δ: 7.53 (br s, 1H); 7.43 (d, 2H); 7.42 (d, 2H); 4.80 (m, 1H); 4.43 (d, 1H); 4.40 (d, 1H); 3.83 (m, 1H); 3.59 (m, 1H); 1.7 (m, 6H).

Part C. Preparation of 3-cyanobenzaldehyde oxime.

20 Hydroxylamine hydrochloride (13.5 g, 194 mmol) was added to a solution of 3-cyanobenzaldehyde (25 g, 191 mmol) in 75 mL of pyridine and 75 mL of ethanol under N₂. This was allowed to stir at room temperature for 14 hours. Water (50 mL) was
25 added with vigorous stirring and an off-white solid precipitated. The solid was filtered through a glass frit and washed with another 50 mL of water. Evaporation of residual water under high vacuum gave 19.2 g (69%) of title compound. ¹H NMR (CDCl₃) δ: 11.61 (s, 1H); 8.21 (s, 1H); 8.00 (s, 1H);
30 7.96 (d, 1H), 7.85 (d, 1H), 7.61 (t, 1H).

Part D. Preparation of N-(4-bromophenyl)-3-(3-cyanophenyl)-5-(tetrahydropyran-2-yloxymethyl)-isoxazo-4-yl-carboxamide and
N-(4-bromophenyl)-3-(3-cyanophenyl)-4-(tetrahydropyran-2-yloxymethyl)-isoxazo-5-yl-carboxamide.
35

To a solution of 2.54 g (17.2 mmol) of 3-cyanobenzaldehyde oxime and 7.10 g (21.0 mmol) of N-(4-

bromophenyl)-4-(tetrahydropyran-2-yloxymethyl)-2-butyramide in 58 mL THF was added 45 mL bleach (0.67M aqueous solution) over a 4-hour period. The solvent was removed in vacuo and the resulting aqueous solution was extracted with 25 mL EtOAc three times. The combined organic extracts were dried with MgSO₄ and the solvent removed in vacuo. Chromatography on silica with 20% EtOAc/hexane gave 1.4 g (17%) of N-(4-bromophenyl)-3-(3-cyanophenyl)-5-(tetrahydropyran-2-yloxymethyl)isoxazo-4-yl-carboxamide and 1.67 g (20%) of N-(4-bromophenyl)-3-(3-cyanophenyl)-4-(tetrahydropyran-2-yloxymethyl)-isoxazo-5-yl-carboxamide. ¹H NMR (CDCl₃) δ: 1st isomer: 9.45 (br s, 1H); 8.11 (s, 1H); 8.04 (d, 1H); 7.77 (d, 1H); 7.58 (t, 1H); 7.50 (m, 4H); 4.98 (dd, 2H); 4.88 (m, 1H); 3.72 (m, 1H); 3.58 (m, 1H); 1.7 (m, 6H). 2nd isomer: 8.66 (br s, 1H); 8.31 (m, 1H); 8.14 (d, 1H); 7.95 (d, 1H); 7.75 (t, 1H), 7.57 (m, 4H); 4.94 (dd, 2H); 4.87 (m, 1H); 3.87 (m, 1H); 3.57 (m, 1H); 1.6 (m, 6H).

Part E. Preparation of 4-(N-[2'-t-butylaminosulfonyl-[1,1']-biphen-4-yl]aminocarbonyl)-3-(3-cyanophenyl)-5-(tetrahydropyran-2-yloxymethyl)-isoxazole.

A mixture of 0.31 g (0.60 mmol) of N-(4-bromophenyl)-3-(3-cyanophenyl)-5-(tetrahydropyran-2-yloxymethyl)isoxazo-4-yl-carboxamide, 0.23 g (0.90 mmol) of 2-(t-butylamino-sulfonyl)phenylboronic acid, 0.052 g (0.045 mmol) of tetrakis(triphenylphosphine) palladium(0), 0.05 mL of 40% aqueous tetrabutylammonium hydroxide, and 0.9 mL of 2M aqueous sodium carbonate were refluxed with 8 mL of toluene under N₂ for 5.5 hours. After cooling, the mixture was separated and the aqueous layer was extracted with 5 mL of ethyl acetate twice. The combined organic extracts were dried with MgSO₄ and concentrated. The resulting solid was chromatographed with 50% EtOAc/hexane to give 0.27 g (73%) of the desired product. ¹H NMR (CDCl₃) δ: 9.57 (br s, 1H); 8.15 (m, 2H); 8.07 (d, 1H); 7.77 (d, 1H); 7.71 (d, 2H); 7.60 (t, 1H); 7.52 (m, 3H); 7.31 (m, 2H); 5.02 (dd, 2H); 4.94 (m, 1H); 3.72 (m, 1H); 3.60 (m, 1H); 1.7 (m, 6H); 1.04 (s, 9H).

Part F. Preparation of 4-[2'-aminosulfonyl-[1,1']-biphen-4-yl]aminocarbonyl-3-(3-cyanophenyl)-5-(hydroxymethyl)-isoxazole.

5

A solution of 0.27 g (0.56 mmol) of 4-(N-[2'-t-butylaminosulfonyl-[1,1']-biphen-4-yl]aminocarbonyl)-3-(3-cyanophenyl)-5-(tetrahydropyran-2-yloxymethyl)isoxazole in 10 mL of trifluoroacetic acid is allowed to stir under N₂ for 16 hours at room temperature. The solvent was removed in vacuo and then chromatographed on silica with 50% EtOAc/hexane to give 0.11 g (51%) of desired product. ¹H NMR (CDCl₃) δ: 9.19 (br s, 1H); 8.12 (d, 1H); 8.05 (m, 1H); 7.99 (d, 1H); 7.81 (d, 1H); 7.64 (t, 1H); 7.58 (m, 3H); 7.50 (m, 1H); 7.42 (d, 2H); 7.31 (d, 1H); 6.77 (m, 1H); 5.03 (d, 2H). HRMS 475.1076 (M+H).

Part G. Preparation of 3-(3-amidinophenyl)-4-[2'-aminosulfonyl-[1,1']-biphen-4-yl]aminocarbonyl-5-(hydroxymethyl)-isoxazole, trifluoroacetic acid salt.

4-(N-[2'-aminosulfonyl-[1,1']-biphen-4-yl]aminocarbonyl)-3-(3-cyanophenyl)-5-(hydroxymethyl)isoxazole (0.11 g, 0.22 mmol) was dissolved in 5 mL of methanol and 10 mL of chloroform. The reaction mixture was cooled in an ice-bath and HCl gas was bubbled-in for 30 minutes to saturate the solution. The mixture was sealed and allowed to stir at room temperature for 14 hours. The solvents were removed in vacuo and the resulting solid was added to 0.5 g (5.2 mmol) of ammonium carbonate and 10 mL of methanol. The mixture was allowed to stir under N₂ for 14 hours. The solvent was removed at reduced pressure. The crude benzamidine was purified by HPLC (C18 reversed phase) eluting with 0.5% TFA in H₂O/CH₃CN to give 0.07 g (53%) of the desired salt. ¹H NMR (DMSO-d₆) δ: 10.60 (s, 1H); 9.43 (br s, 2H); 9.00 (br s, 2H); 8.14 (m, 1H); 7.98 (d, 2H); 7.89 (d, 1H); 7.75 (t, 1H); 7.58 (m, 4H); 7.34 (d, 2H); 7.28 (m, 1H); 4.79 (s, 2H). HRMS 492.1341 (M+H).

Example 2

3-(3-amidinophenyl)-4-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]isoxazole, trifluoroacetic acid salt

5 Part A. Preparation of 3-cyanobenzenehydroximinoyl chloride.

3-Cyanobenzaldehyde oxime (15 g, 103 mmol) was suspended in 90 mL of DMF and then N-chlorosuccinimide (13.7 g, 103 mmol) was added. Approximately 50 mL of gaseous HCl was added
10 via syringe below the liquid surface over a 2 minute period. The reaction was allowed to stir at room temperature for 15 hours and the solution clarified. The solvent was evaporated at 5 torr with a bath temp of 55°C till viscous and cloudy. Water (100 mL) was added with vigorous stirring. An off-white
15 precipitate formed, was filtered through a glass frit and washed with 50 mL water to give 18.1 g (98%) of the desired product after drying under high vacuum. ¹H NMR (CDCl₃) δ: 8.75 (s, 1H), 8.17 (s, 1H), 8.10 (d, 1H), 7.73 (d, 1H), 7.55 (dd, 1H).

20

Part B. Preparation of 2'-t-butylaminosulfonyl-4-amino-[1,1']biphenyl.

A mixture of 3.44 g (20 mmol) of 4-bromoaniline and 5.14
25 g (20 mmol) of 2-(t-butylaminosulfonyl)phenylboronic acid, 1.16 g of tetrakis(triphenylphosphine) palladium(0) (1 mmol), 0.32 g of tetrabutylammonium bromide (1 mmol) and 20 mL of 2M aqueous sodium carbonate were refluxed with 180 mL of benzene under N₂ for 5.5 hours. After cooling, the mixture was diluted
30 with methylene chloride and water. The two phases were separated and the organic phase was washed with water, dried with MgSO₄ and concentrated in vacuo. The resulting thick oil was chromatographed on silica with 30% EtOAc/hexane to afford 2.52 g (41%) of the title compound. ¹H NMR (CDCl₃) δ: 8.14 (d,
35 1H); 7.53 (t, 1H); 7.43 (t, 1H); 7.33 (d, 2H); 7.27 (d, 1H); 6.76 (d, 2H); 3.7 (br s, 1H); 0.99 (s, 9H).

Part C. Preparation of 3-(3-cyanophenyl)-5-carbomethoxy-isoxazole.

Triethylamine (1.01 g, 10 mmol) is added dropwise over 2
5 hours to a solution of 0.72 g (4.0 mmol) of 3-cyanobenzene-
hydroximinoyl chloride and 0.56 g (4.8 mmol) of methyl
methoxyacrylate in 10 mL of CH₂Cl₂ under N₂. The reaction
mixture is diluted with 10 mL of water and the organic layer
separated. The aqueous solution is extracted with 10 mL EtOAc
10 twice and the combined organic extracts are dried with MgSO₄
and concentrated *in vacuo*. The resulting thick oil was
chromatographed on silica with 30% EtOAc/hexane to give 0.90 g
(99%) of the desired product. ¹H NMR (CDCl₃) δ: 9.07 (s, 1H);
8.14 (s, 1H); 8.06 (d, 1H); 7.79 (d, 2H); 7.61 (t, 1H); 3.88
15 (s, 3H).

Part D. Preparation of 3-(3-cyanophenyl)-isoxazole-5-carboxylic acid.

20 A mixture of 0.90 g (3.9 mmol) of 4-carbomethoxy-3-(3-cyanophenyl)isoxazole, 0.25 g (6.0 mmol) of lithium hydroxide monohydrate in 1 mL water and 2 mL methanol is stirred under N₂ for 5 hours. The reaction mixture was acidified to pH 3 with 1N HCl, extracted with 10 mL EtOAc three times, dried with
25 MgSO₄ and concentrated *in vacuo* to give 0.36 g (43%) of the desired acid. ¹H NMR (CDCl₃) δ: 8.94 (s, 1H); 8.01 (s, 1H); 7.94 (d, 1H); 7.60 (d, 2H); 7.43 (t, 1H).

30 Part E. Preparation of 4-(N-[2'-t-butylaminosulfonyl-[1,1']-biphen-4-yl]aminocarbonyl)-3-(3-cyanophenyl)-isoxazole.

Thionyl chloride (10 mL) and 3-(3-cyanophenyl)isoxazole-4-carboxylic acid (0.34 g, 1.6 mmol) are stirred at room temperature under N₂ for 1 hour. The excess thionyl chloride
35 is removed *in vacuo* and the resulting solid is resuspended in 10 mL toluene. The toluene is removed *in vacuo* to remove any residual thionyl chloride. The solid is dissolved in 15 mL CH₂Cl₂ and 0.53 g (1.8 mmol) of 2'-t-butylaminosulfonyl-4-

amino-[1,1']biphenyl and 0.32 g (3.2 mmol) of triethylamine are added. After 2 hours, the reaction mixture is diluted with 10 mL of water and the organic layer separated. The aqueous solution is extracted with 10 mL EtOAc three times and the combined organic extracts are dried with MgSO₄ and concentrated *in vacuo*. The resulting thick oil was chromatographed on silica with 25% EtOAc/hexane to 0.53 g (66%) of the desired product. ¹H NMR (CDCl₃) δ: 9.07 (s, 1H); 8.14 (m, 1H); 8.06 (m, 1H); 7.81 (d, 1H); 7.55 (m, 9H); 1.03 (s, 9H).

Part F. Preparation of 3-(3-amidinophenyl)-4-[2'-aminosulfonyl-[1,1']-biphen-4-yl]aminocarbonyl-5-(hydroxymethyl)-isoxazole, trifluoroacetic acid salt.

4-(N-[2'-aminosulfonyl-[1,1']-biphen-4-yl]aminocarbonyl)-3-(3-cyanophenyl)isoxazole (0.53 g, 1.1 mmol) was dissolved in 10 mL of methanol and 30 mL of chloroform. The reaction mixture was cooled in an ice-bath and HCl gas was bubbled-in for 30 minutes to saturate the solution. The mixture was sealed and allowed to stir at room temperature for 14 hours. The solvents were removed *in vacuo* and the resulting solid was added to 0.5 g (5.2 mmol) of ammonium carbonate and 20 mL of methanol. The mixture was allowed to stir under N₂ for 14 hours. The solvent was removed at reduced pressure. The crude benzamidine was purified by HPLC (C18 reversed phase) eluting with 0.5% TFA in H₂O/CH₃CN to give 0.09 g (15%) of the title salt. ¹H NMR (DMSO-d₆) δ: 10.69 (s, 1H); 9.44 (br s, 2H); 9.06 (br s, 2H); 8.17 (m, 1H); 8.06 (d, 1H); 8.00 (d, 1H); 7.92 (d, 1H); 7.75 (t, 1H); 7.67 (d, 2H); 7.56 (m, 2H); 7.33 (d, 2H); 7.28 (m, 1H). HRMS 462.1252 (M+H).

Example 3

3-(3-amidinophenyl)-4-[(2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]isoxazole, trifluoroacetic acid salt

Part A. Preparation of 2-methylthiophenylboronic acid

2-Bromothioanisole (8.56 g, 42 mmol) was dissolved in 90 mL of dry THF and cooled to -78°C. n-Butyllithium (18.6 mL, 2.5M in hexane, 47 mmol) was added dropwise over 20 minutes, and the resulting solution stirred 90 min. Triisopropylborate (13.7 mL, 59 mmol) was added dropwise over 10 minutes, and the resulting solution stirred at -78°C for 45 minutes before removing the cooling bath. The reaction was stirred overnight at room temperature. HCl (40 mL of a 6M aqueous solution) was added and stirred vigorously 8h. The reaction was diluted with 100 mL water and extracted three times with Et₂O. The organic extracts were combined and extracted twice with 80 mL of 2M NaOH. The basic layers were combined and acidified with 50 mL 6M HCl and 25mL 2M HCl. The resulting cloudy solution was extracted three times with 50 mL of Et₂O, dried over MgSO₄, filtered, and evaporated to yield a white solid (5.22g, 74%). ¹H NMR (CDCl₃) δ: 8.01 (dd, 1H), 7.53 (dd, 1H), 7.43 (td, 1H), 7.34 (td, 1H), 6.22 (s, 2H), 2.50 (s, 3H).

Part B. Preparation of 4-(t-butoxycarbonyl)amino-2'-methylthio-[1,1']biphenyl

2-Methylthiophenylboronic acid (5.2 g, 31 mmol), N-t-butoxycarbonyl-4-bromoaniline (4.0 g, 15 mmol), Na₂CO₃ (31 mL, 2M aqueous), tetrabutylammonium bromide (230 mg, 0.7 mmol), and bis(triphenylphosphine)palladium(II)chloride (515 mg, 0.7 mmol) were combined in 300 mL of benzene, placed briefly under vacuum to degas, and heated at reflux under argon overnight. The reaction was cooled to room temperature and diluted with 100 mL water and 100 mL EtOAc. The organic layer was separated, dried over Na₂SO₄, filtered through celite, and the solvents evaporated. The crude mixture was chromatographed on silica with 10-30% EtOAc/hexane to yield the desired compound (4.17 g, 90%). ¹H NMR (CDCl₃) δ: 7.42 (d, 2H), 7.35 (d, 2H), 7.28 (m, 2H), 7.19 (m, 2H), 6.53 (bs, 1H), 2.36 (s, 3H), 1.53 (s, 9H).

Part C. Preparation of 4-(t-butoxycarbonyl)amino-2'-methylsulfonyl-[1,1']biphenyl

4-(t-Butoxycarbonyl)amino-2'-methylthio-[1,1']biphenyl (4.16 g, 13 mmol) was dissolved in 400 mL of CH₂Cl₂ and cooled to 0°C. MCPBA (11.2 g 57-86%, 37 mmol min.) was added in 4 portions and stirred 25 minutes before removing the cooling bath. The reaction was stirred at room temp for 3 hours. The reaction mixture was then extracted with 50 mL saturated aqueous Na₂SO₃ and then with 50 mL saturated aqueous NaHCO₃. The organic layer was removed, dried over Na₂SO₄, filtered, and evaporated to yield the desired product (4.80 g). ¹H NMR (CDCl₃) δ: 8.22 (dd, 1H), 7.63 (td, 1H), 7.54 (td, 1H), 7.41 (m, 5H), 6.61 (s, 1H), 2.64 (s, 3H), 1.54 (s, 9H).

Part D. Preparation of 4-amino-2'-methylsulfonyl-[1,1']biphenyl

4-(t-butoxycarbonyl)amino-2'-methylsulfonyl-[1,1']biphenyl (4.6 g, 13 mmol), was suspended in 100 mL of 4M HCl in dioxane and stirred 2.5 days. The resulting mixture was filtered and the cake rinsed with Et₂O to yield a tan solid (3.69 g, 98%). ¹H NMR (DMSO-d₆) δ: 8.04 (d, 1H), 7.71 (t, 1H), 7.61 (t, 1H), 7.31 (m, 3H), 7.06 (m, 2H), 2.79 (s, 3H).

Part E. Preparation of 3-(3-cyanophenyl)-4-[(2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]isoxazole

4-Amino-2'-methylsulfonyl-[1,1']biphenyl (0.50 g, 2.2 mmol) was suspended in 10 mL of CH₂Cl₂ and 4.4 mL of a 2M solution of trimethylaluminum in heptane was added slowly via syringe. The reaction was stirred for 30 minutes at room temperature and then 3-(3-cyanophenyl)-5-carbomethoxyisoxazole (0.62 g, 2.2 mmol) was added. The reaction mixture was stirred at room temperature for an additional 14 hours. The aluminum reagent was quenched by careful addition of 1N HCl to pH 2, then extracted with 10 mL of CH₂Cl₂ three times. The combined organic extracts were washed with water then brine, dried over MgSO₄ and the solvent evaporated. The desired product was obtained (0.74 g, 76%) after silica gel

chromatography with 30% EtOAc/hexane. ^1H NMR (CDCl_3) δ : 9.07 (s, 1H); 8.31 (s, 1H); 8.21 (d, 1H); 8.09 (s, 1H); 8.07 (d, 1H); 7.79 (d, 1H); 7.63 (m, 4H); 7.42 (d, 2H); 7.37 (d, 1H); 2.72 (s, 3H).

5

Part F. Preparation of 3-(3-amidinophenyl)-4-[(2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]isoxazole, trifluoroacetic acid salt

10 3-(3-Cyanophenyl)-4-[(2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]isoxazole (0.74 g, 1.7 mmol) was dissolved in 10 mL of methanol and 40 mL of chloroform. The reaction mixture was cooled in an ice-bath and HCl gas was bubbled-in for 1.5 hours to saturate the solution. The mixture was
15 sealed and allowed to stir at room temperature for 14 hours. The solvents were removed in vacuo and the resulting solid was added to 0.66 g (8.5 mmol) of ammonium carbonate and 20 mL of methanol. The mixture was sealed and allowed to stir under Ar for 14 hours. The solvent was removed at reduced pressure.
20 The crude benzamidine was purified by HPLC (C18 reversed phase) eluting with 0.5% TFA in $\text{H}_2\text{O}/\text{CH}_3\text{CN}$ to give 0.33 g (43%) of the desired salt. ^1H NMR ($\text{DMSO}-d_6$) δ : 10.69 (s, 1H); 9.67 (s, 1H); 9.41 (br s, 2H); 9.07 (br s, 2H); 8.18 (t, 1H); 8.06 (dt, 2H); 7.92 (d, 1H); 7.72 (m, 5H); 7.38 (s, 1H); 7.36 (d, 2H); 2.80 (s, 3H). HRMS 461.1284 (M+H).
25

Example 4

3-(3-amidinophenyl)-4-[5-(2-aminosulfonyl)phenylpyrid-2-yl)aminocarbonyl]-5-(methoxymethyl)isoxazole, trifluoroacetic
30 acid salt

Part A. Preparation of 3-(3-cyanophenyl)-5-methoxymethyl-4-carbomethoxyisoxazole

35 Methyl 4-methoxyacetoacetate (1.6 mL, 12 mmol) was dissolved in 12 mL of 2M sodium methoxide in methanol. 3-Cyanobenzenehydroximinoyl chloride (2.0 g, 11 mmol) was dissolved in 10 mL methanol and added to the basic solution

over a 5-hour period via syringe pump. The reaction was quenched with 20 mL of saturated, aqueous ammonium chloride. The mixture was extracted with 30 mL EtOAc three times and the combined organic extracts washed with 10 mL water three times. 5 The resulting solution was dried with MgSO_4 , concentrated in vacuo and then chromatographed on silica with 10% Et_2O /benzene to 1.14 g (38%) of a white solid. ^1H NMR (CDCl_3) δ : 7.95 (m, 1H); 7.92 (dd, 1H); 7.98 (dd, 1H); 7.60 (t, 1H); 4.91 (s, 2H); 3.83 (s, 3H); 3.54 (s, 3H).

10

Part B. Preparation of 2-amino-5-(2-t-butylaminosulfonyl)phenylpyridine

A mixture of 1.55 g (9.0 mmol) of 2-amino-5-bromopyridine 15 and 2.3 g (9.0 mmol) of 2-(t-butylaminosulfonyl)phenylboronic acid, 0.52 g of tetrakis(triphenylphosphine) palladium(0) (0.45 mmol), 0.15 g of tetrabutylammonium bromide (0.45 mmol) and 9 mL of 2M aqueous sodium carbonate were refluxed with 80 mL of benzene under Ar for 5 hours. After cooling, the 20 mixture was diluted with 25 mL of methylene chloride and 25 mL of water. The two phases were separated and the organic phase was washed with water, dried with MgSO_4 and concentrated in vacuo. The resulting thick oil was chromatographed on silica with 50% EtOAc/hexane to afford 1.34 g (49%) of the aniline. 25 ^1H NMR (CDCl_3) δ : 8.18 (d, 1H); 8.07 (m, 1H); 7.70 (dd, 1H); 7.58 (dt, 1H); 7.48 (dt, 1H); 7.28 (d, 1H); 6.56 (d, 1H); 4.62 (br s, 2H); 3.88 (br s, 1H); 1.06 (s, 9H).

Part C. Preparation of 3-(3-cyanophenyl)-4-[5-(2-t-butylaminosulfonyl)phenylpyrid-2-yl]aminocarbonyl]-5-(methoxymethyl)isoxazole 30

3-(3-cyanophenyl)-5-methoxymethyl-4-carbomethoxyisoxazole (1.12 g, 4.1 mmol) was dissolved in 3 mL of THF and 1 mL 35 water. Lithium hydroxide monohydrate (0.20 g, 4.9 mmol) was added and the reaction stirred at room temperature for 5 hours. The solvent was evaporated in vacuo, 100 mL of water was added and the mixture extracted with 50 mL of EtOAc twice.

The combined organic extracts were dried with MgSO_4 concentrated in vacuo to give 0.8 g (75%) of a white solid. The crude carboxylic acid (0.4g, 1.6 mmol) was dissolved in 1.2 mL of 2.0M oxalyl chloride in CH_2Cl_2 followed by 0.1 mL of

5 DMF. The reaction was allowed to stir at room temperature for 2 hours. The reaction was concentrated under high vacuum for 30 minutes to yield a yellow-orange solid. The crude acid chloride was dissolved in 5 mL CH_2Cl_2 . 2-Amino-5-(2-t-butylaminosulfonyl)phenylpyridine (0.51 g, 1.86 mmol) followed

10 by triethylamine (0.65 mL, 4.65 mmol) was added to the crude acid chloride solution. The reaction mixture was allowed to stir at room temperature for 14 hours. The solution was diluted with 50 mL CH_2Cl_2 , washed with 25 mL saturated, aqueous NaHCO_3 , 25 mL 1M HCl then 25 mL brine. The organic layer was

15 dried with MgSO_4 , concentrated in vacuo, and chromatographed on silica with 20% EtOAc/benzene to give 0.10 g (12%) of the desired product. ^1H NMR (CDCl_3) δ : 10.07 (br s, 1H); 8.40 (d, 1H); 8.30 (d, 1H); 8.18 (dd, 1H); 8.09 (m, 1H); 8.02 (dt, 1H); 7.84 (dd, 1H); 7.79 (dt, 1H); 7.60 (m, 3H); 7.28 (dd, 1H);

20 4.89 (s, 2H); 3.71 (s, 3H); 1.07 (s, 9H).

Part D. Preparation of 3-(3-amidinophenyl)-4-[5-(2-aminosulfonyl)phenylpyrid-2-yl)aminocarbonyl]-5-(methoxymethyl)isoxazole, trifluoroacetic acid salt

25

3-(3-cyanophenyl)-4-[5-(2-t-butylaminosulfonyl)phenylpyrid-2-yl)aminocarbonyl]-5-(methoxymethyl)isoxazole (0.10 g, 0.18 mmol) was dissolved in 1 mL of methanol and 4 mL of chloroform. The reaction mixture

30 was cooled in an ice-bath and HCl gas was bubbled-in for 2.5 hours to saturate the solution. The mixture was sealed and allowed to stir at room temperature for 14 hours. The solvents were removed in vacuo and the resulting solid was added to 0.07 g (0.90 mmol) of ammonium carbonate and 10 mL of

35 methanol. The mixture was sealed and allowed to stir under Ar for 14 hours. The solvent was removed at reduced pressure. The crude benzamidine was purified by HPLC (C18 reversed phase) eluting with 0.5% TFA in $\text{H}_2\text{O}/\text{CH}_3\text{CN}$ to give 0.33 g (43%)

of the desired salt. ^1H NMR ($\text{DMSO}-d_6$) δ : 10.69 (s, 1H); 9.67 (s, 1H); 9.41 (br s, 2H); 9.07 (br s, 2H); 8.18 (t, 1H); 8.06 (dt, 2H); 7.92 (d, 1H); 7.72 (m, 5H); 7.38 (s, 1H); 7.36 (d, 2H); 2.80 (s, 3H). HRMS 507.1458 (M+H).

5

Example 5

3-(3-amidinophenyl)-4-[(2'-trifluoromethyl-[1,1']-biphen-4-yl)aminocarbonyl]isoxazole, trifluoroacetic acid salt

10 Part A. Preparation of 2-(trifluoromethyl)phenylboronic acid

To a solution of 58.8 g (0.261 mol) of 1-bromo-2-(trifluoromethyl)benzene in 250 mL of THF under Ar was added 110 mL (0.275 mol) of 2.5M n-butyllithium in hexane over 35 minutes, keeping the temperature between 0-5°C. The reaction mixture was allowed to warm to 10°C. Triisopropylborate (95 mL, 0.313 mol) was added, keeping the temperature below 35°C. After 1 hour, the reaction mixture was cooled, 1N HCl (425 mL) was added, and the mixture was stirred overnight. The mixture was extracted with 100 mL of ether three times, and the combined organic extracts were extracted with 100 mL of 1N NaOH three times. The aqueous extracts were acidified to pH 1 with 6N HCl, and then extracted with 100 mL ether three times. The combined ether extracts were dried over MgSO_4 , and the solvents evaporated in vacuo to give 46.1 g (93%) of the desired compound as a light yellow oil. ^1H NMR (CDCl_3) δ : 7.77 (d, 1H); 7.72 (d, 1H); 7.56 (m, 2H); 4.87 (br s, 2H).

30 Part B. Preparation of 4-amino-2'-trifluoromethyl-[1,1']biphenyl

A mixture of 3.44 g (20 mmol) of 4-bromoaniline and 3.80 g (20 mmol) of 2-(trifluoromethyl)phenylboronic acid, 1.16 g of tetrakis(triphenylphosphine) palladium(0) (1 mmol), 0.32 g of tetrabutylammonium bromide (1 mmol) and 20 mL of 2M aqueous sodium carbonate were refluxed with 180 mL of benzene under N_2 for 14 hours. After cooling, the mixture was diluted with methylene chloride and water. The two phases were separated

and the organic phase was washed with water, dried with MgSO_4 and concentrated *in vacuo*. The resulting thick oil was chromatographed on silica with 10% EtOAc/hexane to afford 2.09 g (44%) of the aniline. ^1H NMR (CDCl_3) δ : 7.72 (d, 1H); 7.53 (t, 1H); 7.41 (t, 1H); 7.32 (d, 1H); 7.13 (d, 2H); 6.73 (d, 2H); 3.74 (br s, 2H).

Part C. Preparation of 3-(3-cyanophenyl)-4-[(2'-trifluoromethyl-[1,1']-biphen-4-yl)aminocarbonyl]isoxazole

4-Amino-2'-trifluoromethyl-[1,1']biphenyl (0.24 g, 1.0 mmol) was suspended in 5 mL of CH_2Cl_2 and 2.5 mL of a 2M solution of trimethylaluminum in heptane was added slowly via syringe. The reaction was stirred for 30 minutes at room temperature and then 4-carbomethoxy-3-(3-cyanophenyl)isoxazole (0.25 g, 1.0 mmol) was added. The reaction mixture was stirred at room temperature for an additional 14 hours. The aluminum reagent was quenched by careful addition of 1N HCl to pH 2, then extracted with 10 mL of CH_2Cl_2 three times. The combined organic extracts were washed with water then brine, dried over MgSO_4 and the solvent evaporated. The desired product was obtained (0.35 g, 80%) after silica gel chromatography with 20% EtOAc/hexane. ^1H NMR (CDCl_3) δ : 9.02 (s, 1H); 8.17 (s, 1H); 8.08 (d, 1H); 7.83 (d, 1H); 7.76 (d, 1H); 7.67 (t, 1H); 7.56 (m, 1H); 7.48 (m, 3H); 7.34 (m, 3H).

Part D. Preparation of 3-(3-amidinophenyl)-4-[(2'-trifluoromethyl-[1,1']-biphen-4-yl)aminocarbonyl]isoxazole, trifluoroacetic acid salt

3-(3-cyanophenyl)-4-[(2'-trifluoromethyl-[1,1']-biphen-4-yl)aminocarbonyl]isoxazole (0.33 g, 0.76 mmol) was dissolved in 1 mL of methanol and 3 mL of chloroform. The reaction mixture was cooled in an ice-bath and HCl gas was bubbled-in for 0.5 hours to saturate the solution. The mixture was sealed and allowed to stir at room temperature for 14 hours. The solvents were removed *in vacuo* and the resulting solid was added to 0.25 g (2.6 mmol) of ammonium carbonate and 2 mL of

methanol. The mixture was sealed and allowed to stir under Ar for 14 hours. The solvent was removed at reduced pressure. The crude benzamidine was purified by HPLC (C18 reversed phase) eluting with 0.5% TFA in H₂O/CH₃CN to give 0.11 g (32%) of the desired salt. ¹H NMR (DMSO-d₆) δ: 10.68 (s, 1H); 9.68 (s, 1H); 9.43 (br s, 2H); 9.06 (br s, 2H); 8.20 (m, 1H); 8.08 (d, 1H); 7.93 (d, 1H); 7.75 (m, 5H); 7.60 (d, 1H); 7.38 (m, 1H); 7.31 (d, 2H). HRMS 451.1399 (M+H).

10

Example 6

3-(3-amidinophenyl)-4-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-5-(trifluoromethyl)isoxazole, trifluoroacetic acid salt

15 Part A. Preparation of ethyl 3-methoxy-3-(trifluoromethyl)acrylate

Diazald® (14.55, 67.9 mmol) was dissolved in 130 mL Et₂O and 30 mL 95% EtOH. Potassium hydroxide (12.37 g, 220 mmol) is dissolved in 23 mL water and added slowly via additional funnel to the ethanol solution which is heated to 65°C. The ether distillate containing diazomethane is condensed into ethyl 4,4,4-trifluoroacetoacetate (10.0 g, 54.3 mmol). The excess diazomethane was decomposed with the addition of 1 drop of acetic acid. The ethereal solution was evaporated at 450 torr for 30 minutes. The crude enol ether (10.7 g, 100%) was used without purification. ¹H NMR (CDCl₃) δ: 5.78 (s, 1H); 4.22 (q, 2H); 4.04 (s, 3H); 1.32 (t, 3H).

30 Part B. Preparation of 3-(3-cyanophenyl)-5-(trifluoromethyl)-4-carbomethoxyisoxazole

Tributylamine (12.6 g, 67.9 mmol) is added dropwise over 2 hours to a solution of 9.81 g (54.3 mmol) of 3-cyanobenzene-hydroximinoyl chloride and 10.76 g (54.3 mmol) of ethyl 3-methoxy-3-(trifluoromethyl)acrylate in 49 mL of CH₂Cl₂ and 1 mL DMSO under N₂. The reaction mixture is diluted with 100 mL of water and the organic layer separated. The aqueous solution

is extracted with 100 mL EtOAc twice and the combined organic extracts are dried with MgSO_4 and concentrated *in vacuo*. The resulting thick oil was chromatographed on silica with 30% EtOAc/hexane to give 1.60 g (10%) of the desired product. ^1H NMR (CDCl_3) δ : 8.05 (m, 1H); 7.96 (dt, 1H); 7.84 (dt, 1H); 7.63 (t, 1H); 4.37 (q, 2H); 1.33 (t, 3H).

Part C. Preparation of 3-(3-cyanophenyl)-4-[(2'-t-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-5-(trifluoromethyl)isoxazole

2'-t-Butylaminosulfonyl-4-amino-[1,1']biphenyl (0.39 g, 1.3 mmol) was suspended in 4 mL of CH_2Cl_2 and 1.9 mL of a 2M solution of trimethylaluminum in heptane was added slowly via syringe. The reaction was stirred for 30 minutes at room temperature and then 3-(3-cyanophenyl)-5-(trifluoromethyl-4-carbomethoxyisoxazole (0.40 g, 1.3 mmol) was added. The reaction mixture was stirred at room temperature for an additional 14 hours. The aluminum reagent was quenched by careful addition of 1N HCl to pH 2, then extracted with 20 mL of CH_2Cl_2 three times. The combined organic extracts were washed with water then brine, dried over MgSO_4 and the solvent evaporated. The desired product was obtained (0.31 g, 43%) after silica gel chromatography with 20% EtOAc/hexane. ^1H NMR (CDCl_3) δ : 8.74 (br s, 1H); 8.11 (d, 1H); 8.05 (m, 1H); 8.02 (d, 1H); 7.76 (d, 1H); 7.55 (m, 5H); 7.37 (d, 2H); 7.28 (d, 1H); 1.03 (s, 9H).

Part D. Preparation of 3-(3-amidinophenyl)-4-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-5-(trifluoromethyl)isoxazole, trifluoroacetic acid salt

3-(3-cyanophenyl)-4-[(2'-t-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-5-(trifluoromethyl)isoxazole (0.30 g, 0.53 mmol) was dissolved in 1 mL of methanol and 3 mL of chloroform. The reaction mixture was cooled in an ice-bath and HCl gas was bubbled-in for 0.5 hours to saturate the solution. The mixture was sealed and allowed to stir at room

temperature for 14 hours. The solvents were removed in vacuo and the resulting solid was added to 0.25 g (2.6 mmol) of ammonium carbonate and 2 mL of methanol. The mixture was sealed and allowed to stir under Ar for 14 hours. The solvent
5 was removed at reduced pressure. The crude benzamidine was purified by HPLC (C18 reversed phase) eluting with 0.5% TFA in H₂O/CH₃CN to give 0.11 g (36%) of the desired salt. ¹H NMR (DMSO-d₆) δ: 11.17 (s, 1H); 9.46 (br s, 2H); 9.09 (br s, 2H); 8.19 (m, 1H); 7.98 (m, 3H); 7.84 (t, 1H); 7.56 (m, 2H); 7.55
10 (d, 2H); 7.37 (d, 2H); 7.26 (m, 1H). HRMS 586.1743 (M+H).

Examples 7 and 8

2-Acetylamino-4-(3-amidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole, trifluoroacetic acid salt
15 (Example 7) and 2-amino-4-(3-amidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole, trifluoroacetic acid salt (Example 8)

Part A. Preparation of methyl 3-(3-cyanophenyl)-3-hydroxypropionate.
20

To a suspension of activated zinc powder (4.3 g, 65.4 mmol) in 100 mL of tetrahydrofuran was added a few drops of dibromoethane. The resulting mixture was heated to 65° C,
25 stirred for 5 min and then was cooled to 25° C. To this solution was added methyl bromoacetate (5.0 g, 32.7 mmol) and 3-cyanobenzaldehyde (4.3 g, 32.7 mmol). The mixture was heated to 65° C and stirred for 2 h. The reaction was allowed to cool to 25° C and then was quenched with 10% aq HCl and
30 filtered through celite. The mixture was diluted with ethyl acetate and washed with 10% aq HCl and saturated aq NaHCO₃. This wash cycle was repeated until no white precipitate was observed upon addition of saturated aq NaHCO₃. The organics were then washed with brine, dried (MgSO₄) and concentrated in
35 vacuo. The residue was purified by flash chromatography (elution with 1:1 hexanes/ethyl acetate) to afford 4.5 g (67%) of the title compound as an oil. MS (NH₃-DCI) 223.1 (M+H)+.

Part B. Preparation of methyl 3-(3-cyanophenyl)-3-oxopropionate.

To a solution of methyl 3-(3-cyanophenyl)-3-hydroxypropionate (2.22 g, 10.8 mmol) in 30 mL of methylene chloride was added activated manganese dioxide (4.7 g, 54.0 mmol). This mixture was allowed to stir at 25° C for 16 h. The reaction mixture was filtered through a pad of celite and concentrated *in vacuo*. The residue was purified by flash chromatography (elution with 2:1 hexanes/ethyl acetate) to afford 0.9 g (41%) of the title compound along with 0.8 g (36%) of recovered starting material. MS for title compound (H₂O GC-MS) 204 (M+H)+.

Part C. Preparation of methyl 2-bromo-3-(3-cyanophenyl)-3-oxopropionate.

To a solution of methyl 3-(3-cyanophenyl)-3-oxopropionate (0.91 g, 4.48 mmol) in 20 mL of carbon tetrachloride at 0° C was added N-bromosuccinimide (0.80 g, 4.48 mmol). The resulting solution was allowed warm to 25° C and was stirred for 2 h. The insoluble succinimide was filtered off and the solution was concentrated *in vacuo* to afford an oil (1.2 g, 95%) which was sufficiently pure to be used without purification. MS (H₂O GC-MS) 282/284 (M+H)+.

Part D. Preparation of 2-acetylamino-4-(3-cyanophenyl)-5-carbomethoxythiazole.

To a solution of methyl 2-bromo-3-(3-cyanophenyl)-3-oxopropionate (1.25 g, 4.4 mmol) in 20 mL of tetrahydrofuran was added 1-acetylthiourea (0.52 g, 4.4 mmol). The resulting mixture was stirred at 65° C for 3h. The reaction was allowed to cool and the solvent was evaporated *in vacuo*. The residue was taken up in ethyl acetate, washed with 10% aq HCl, saturated aq NaHCO₃ and brine, dried (MgSO₄) and concentrated *in vacuo* to yield a solid. Trituration with hexanes/ethyl

acetate left the title compound as a white solid (0.4 g, 30%).
MS (ESI) 302.2 (M+H)+.

Part E. Preparation of 2-acetylamino-4-(3-cyanophenyl)-5-
5 [(2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole.

To a solution of (2'-tert-butylaminosulfonyl-[1,1']-
biphen-4-yl)amine (0.27 g, 0.88 mmol) in 5 mL of methylene
10 chloride at 25° C was added trimethylaluminum (1.76 mL of a 2.0
M solution in toluene, 3.52 mmol) dropwise. The resulting
solution was allowed to stir until no more gas evolution was
observed (~15 min). To this solution was added 2-acetylamino-
4-(3-cyanophenyl)-5-carbomethoxythiazole (0.12 g, 0.40 mmol)
15 as a solution in methylene chloride. The resulting solution
was stirred at 40° C for 2 h and then was cooled to 25° C and
quenched by the addition of saturated aq NH₄Cl. After diluting
with ethyl acetate, the organic layer was washed with 10% aq
HCl, saturated aq NaHCO₃ and brine, dried (MgSO₄) and
20 concentrated *in vacuo*. The residue was purified by flash
chromatography (elution with 1:1 hexanes/ethyl acetate) to
afford 0.15 g (65%) of the title compound as a solid. MS
(ESI) 574 (M+H)+.

25 Part F. Preparation of 2-acetylamino-4-(3-amidinophenyl)-5-
[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole,
trifluoroacetic acid salt (Example 7) and 2-amino-4-(3-
amidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-
yl)aminocarbonyl]thiazole, trifluoroacetic acid salt (Example
30 8).

Through a solution of 2-acetylamino-4-(3-cyanophenyl)-5-
[(2'-tert-butylaminosulfonyl-[1,1']-biphen-4-
yl)aminocarbonyl]thiazole (0.15 g, 0.26 mmol) in 50 mL of
35 absolute methanol at 0° C was bubbled anhydrous HCl (g) until
the solution was saturated. This solution was tightly
stoppered and allowed to stand at 0° C for 16 h. The solution
was concentrated *in vacuo* and then was taken up in 10 mL of

absolute methanol and then there was added ammonium carbonate (0.15 g, 1.56 mmol). This mixture was allowed to stir at 25° C for 16 h. The reaction mixture was then concentrated *in vacuo* and purified by prep HPLC (C18 reverse phase column, elution with a H₂O/CH₃CN gradient with 0.5% TFA) to afford 2-acetylamino-4-(3-amidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole, trifluoroacetic acid salt (Example 7) as the major product (0.50 g, 30%). MS (ESI) 535 (M+H)⁺. There was also isolated 2-amino-4-(3-amidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole, trifluoroacetic acid salt (Example 8) as a minor product (0.10 g, 6%). MS (ESI) 493 (M+H)⁺.

Example 9

2-Methyl-4-(3-amidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole, trifluoroacetic acid salt

Part A. Preparation of 2-methyl-4-(3-cyanophenyl)-5-carbomethoxythiazole.

To a solution of methyl 2-bromo-3-(3-cyanophenyl)-3-oxopropionate from Example 1, Part C (0.50 g, 1.77 mmol) in 10 mL of tetrahydrofuran was added thioacetamide (0.14 g, 1.77 mmol). The resulting solution was stirred at 65° C for 4h and then was allowed to cool to 25° C. This mixture was diluted with ethyl acetate, washed with 10% aq HCl, saturated aq NaHCO₃ and brine, dried (MgSO₄) and concentrated *in vacuo* to a solid. Trituration with ether left 0.14 g (31%) of the title compound as a solid. MS (NH₃-CI) 259 (M+H)⁺.

Part B. Preparation of 2-methyl-4-(3-cyanophenyl)-5-[(2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole.

Following the procedure of Example 1, Part E, 2-methyl-4-(3-cyanophenyl)-5-carbomethoxythiazole (0.08 g, 0.31 mmol) was converted into 0.085 g (52%) of the title compound as a solid. MS (ESI) 531.3 (M+H)⁺.

Part C. Preparation of 2-methyl-4-(3-amidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole, trifluoroacetic acid salt.

5

A solution of 2-methyl-4-(3-cyanophenyl)-5-[(2'-*tert*-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole (0.085 g, 0.16 mmol) in 5 mL of trifluoroacetic acid was stirred at 70° C until gas evolution was no longer observed (~ 15 min) and then was allowed to cool to room temperature and as concentrated *in vacuo*. The crude residue was dissolved in 40 mL of absolute methanol and cooled to 0° C. Anhydrous HCl gas was bubbled through the solution until saturated (~ 30 min). The flask was then sealed and allowed to stand at 0° C for 16 h. The reaction mixture was concentrated *in vacuo*, dissolved in 10 mL of absolute methanol and then ammonium carbonate (0.09 g, 0.96 mmol) was added and the mixture was allowed to stir at 25° C for 24 h. The reaction mixture was concentrated *in vacuo* and purified by prep HPLC (C18 reverse phase column, elution with a H₂O/CH₃CN gradient with 0.5% TFA) to afford 65 mg (68%) of the title compound as a solid. MS (ESI) 492.3 (N+H)+.

Example 10

25 5-(3-amidinophenyl)-4-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]oxazole

Part A. Preparation of 5-(3-cyanophenyl)-4-carboxymethyloxazole.

30

The title compound was prepared in 50% yield by the condensation of 3-cyanobenzoyl chloride, methylisocyanoacetate and triethylamine in anhydrous THF following the general method of Suzuki et. al. Syn. Comm. 1972, 2, 237. ¹H NMR (CDCl₃) δ: 8.42-8.40 (d, 1H), 7.99 (s, 1H), 9.77-7.75 (d, 1H), 7.63 (t, 1H), 3.99 (s, 3H) ppm; Ammonia mass spectrum analysis m/z (rel intensity) 246(M+NH₄⁺, 100), 229(M+H).

Part B. Preparation of 5-(3-cyanophenyl)-oxazole-4-carboxylic acid.

Standard LiOH hydrolysis in aqueous THF of the product from part A then provided pure oxazole carboxylic acid in quantitative yield. ^1H NMR (CDCl_3) δ : 8.56-8.34 (d, 1H), 8.51 (s, 1H), 8.03 (s, 1H), 7.77-7.76 (d, 1H), 7.67-7.62 (t, 1H) ppm; Ammonia mass spectrum analysis m/z (rel intensity) 232 ($\text{M}+\text{NH}_4^+$, 100) ppm.

10

Part C. Preparation of 5-(3-cyanophenyl)-4-[(2'-t-butyl-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]oxazole.

The acid obtained in part B was then coupled (60% yield) to 2'-tert-butylsulfonamide-biphenylaniline acid chloride as described above. ^1H NMR (CDCl_3) δ : 9.24 (s, 1H), 8.71 (s, 1H), 8.67 (s, 1H), 8.19-8.16 (m, 1H), 8.01 (s, 1H), 7.83-7.80 (d, 2H), 7.76 (m, 1H), 7.67 (d, 1H), 7.35-7.32 (m, 1H), 3.62 (s, 1H), 1.03 (s, 9H) ppm; ESI mass spectrum analysis m/z (rel. intensity) 445 ($\text{M}+\text{H}$, 100).

20

Part D. Preparation of 5-(3-amidinophenyl)-4-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]oxazole.

The compound obtained in part C was subjected to the Pinner-amidine reaction sequence as outlined previously to obtain the title benzamidine compound in 40% yield. Colorless crystals obtained after lyophilization. ^1H NMR ($\text{DMSO}-d_6$) δ : 10.49 (s, 1H), 9.45 (bs, 2H), 9.12 (bs, 2H), 8.85 (s, 1H), 8.61 (s, 1H), 8.54-8.51 (d, 1H), 8.05 (d, 1H), 7.91-7.77 (m, 4H), 7.65-7.54 (m, 2H), 7.40-7.37 (d, 2H), 7.35 (d, 1H), 7.27 (s, 2H) ppm; ESI mass spectrum analysis m/z (rel. intensity) 462 ($\text{M}+\text{H}$, 100); High resolution mass spectrum analysis calc. for $\text{C}_{23}\text{H}_{20}\text{N}_5\text{SO}_4$ 462.123601, found 462.124334.

35

Example 11**3-(3-amidinophenyl)-4-[(2'-t-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]isoxazole, trifluoroacetic acid salt**

5 4-(N-[2'-Aminosulfonyl-[1,1']-biphen-4-yl]aminocarbonyl)-3-(3-cyanophenyl)isoxazole (0.53 g, 1.1 mmol) was dissolved in 10 mL of methanol and 30 mL of chloroform. The reaction mixture was cooled in an ice-bath and HCl gas was bubbled for 30 minutes to saturate the solution. The mixture was sealed
10 and allowed to stir at room temperature for 14 hours. The solvents were removed in vacuo and the resulting solid was used in the next step.

The imidate formed above was added with 0.5 g (5.2 mmol) of ammonium carbonate and 20 mL of methanol. The mixture was
15 allowed to stir under N₂ for 14 hours. The solvent was removed at reduced pressure. The crude benzamidine was purified by HPLC (C18 reversed phase) eluting with 0.5% TFA in H₂O/CH₃CN to give 0.09 g (16%) of the desired salt. ¹H NMR (DMSO-d₆) δ: 10.69 (s, 1H); 9.70 (s, 1H); 9.43 (br s, 2H); 9.05 (br s, 2H); 8.05
20 (d, 1H); 8.00 (d, 1H); 7.92 (d, 1H); 7.74 (t, 1H); 7.67 (d, 2H); 7.59 (t, 1H); 7.52 (t, 1H); 7.34 (d, 2H); 7.26 (m, 1H); 0.98 (s, 9H). HRMS 517.1768 (M+H).

Example 12

25 **3-(3-amidinophenyl)-4-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-5-(methoxymethyl)isoxazole, trifluoroacetic acid salt**

Part A. Preparation of methyl 3-(3-cyanophenyl)-5-(methoxymethyl)isoxazole-4-carboxylate.
30

Methyl 4-methoxyacetoacetate (1.6 mL, 12.2 mmol) was added to a solution of 12.2 mL (24.4 mmol) of 0.5 M NaOMe in methanol. A solution of 3-cyanobenzenehydroximinoyl chloride
35 (2.0 g, 11.1 mmol) in 20 mL methanol was added slowly over 12 hours via a syringe pump. The reaction mixture was diluted with 20 mL of saturated aqueous ammonium chloride and the organic layer separated. The aqueous solution was extracted

with 10 mL EtOAc twice. The combined organic extracts were washed with 10 mL water three times then the organic extract was dried with MgSO_4 and concentrated *in vacuo*. The resulting thick oil (3.1 g) was chromatographed on silica with 10% Et_2O /benzene to give 1.14 g (38%) of the desired product. ^1H NMR (CDCl_3) δ : 7.98 (m, 1H); 7.91 (dd, $J = 8.1, 2.9, 1\text{H}$); 7.79 (dd, $J = 8.1, 2.9, 1\text{H}$); 7.59 (t, $J = 8.1, 1\text{H}$); 4.91 (s, 2H); 3.83 (s, 3H); 3.53 (s, 3H) MS ($\text{NH}_3\text{-CI}$) m/z 273.0 (M+H).

10 Part B. Preparation of 3-(3-cyanophenyl)-5-(methoxymethyl)-isoxazole-4-carboxylic acid.

Methyl 3-(3-cyanophenyl)-5-(methoxymethyl)isoxazole-4-carboxylate (1.12 g, 4.1 mmol) was dissolved in 3 mL THF and 1 mL water. Lithium hydroxide monohydrate (0.20 g, 4.9 mmol) was added and the reaction was allowed to stir for 24 hours under N_2 . The solvent was removed *in vacuo* and redissolved in 100 mL of water. The resulting solution was extracted with 30 mL EtOAc twice then acidified with 1N HCl to pH 3. The acidic solution was extracted three times with 30 mL of EtOAc. The combined organic were dried with MgSO_4 and concentrated *in vacuo* to give the desired white solid (0.80 g, 75%). ^1H NMR (CDCl_3) δ : 7.98 (m, 1H); 7.91 (dd, $J = 8.1, 2.9, 1\text{H}$); 7.79 (dd, $J = 8.1, 2.9, 1\text{H}$); 7.59 (t, $J = 8.1, 1\text{H}$); 4.89 (s, 2H); 3.53 (s, 3H).

Part C. Preparation of 3-(3-cyanophenyl)-4-[(2'-t-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-5-(methoxymethyl)isoxazole.

30 Oxalyl chloride (1.15 mL of 2.0 M solution in CH_2Cl_2 , 2.3 mmol) and 3-(3-cyanophenyl)-5-(methoxymethyl)isoxazole-4-carboxylic acid (0.40 g, 1.55 mmol) were stirred at room temperature under N_2 for 1 hour. The excess oxalyl chloride was removed *in vacuo*. The solid was dissolved in 15 mL CH_2Cl_2 and 0.58 g (1.9 mmol) of 2'-t-butylaminosulfonyl-4-amino-[1,1']-biphenyl and 0.65 mL (4.7 mmol) of triethylamine were added. After 12 hours, the reaction mixture was diluted with

10 mL of water and the organic layer separated. The aqueous solution was extracted with 10 mL CH₂Cl₂ three times and the combined organic extracts were washed with 10 mL of each of the following: saturated aqueous NaHCO₃, 1M aqueous HCl and
5 saturate aqueous brine. The organic extract was dried with MgSO₄ and concentrated *in vacuo* to give 0.63 g of an orange-brown solid. The solid was chromatographed on silica with 10% EtOAc/benzene to 0.63 g (74%) of the desired product. ¹H NMR (CDCl₃) δ: 9.50 (s, 1H); 8.16 (d, J = 8.1, 1H); 8.11 (s, 1H);
10 8.04 (d, J = 8.1, 1H); 7.78 (d, J = 8.1, 1H); 7.66 (d, J = 8.8, 2H); 7.60 (m, 1H); 7.56 (m, 2H); 7.50 (d, J = 8.8, 2H); 7.29 (d, J = 7.3, 1H); 4.89 (s, 2H); 3.67 (s, 3H); 1.03 (s, 9H).

15 Part D. Preparation of 3-(3-amidinophenyl)-4-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-5-(methoxymethyl)isoxazole, trifluoroacetic acid salt.

3-(3-Cyanophenyl)-4-[(2'-t-butylaminosulfonyl-[1,1']-
20 biphen-4-yl)aminocarbonyl]-5-(methoxymethyl)isoxazole (0.62 g, 1.14 mmol) was dissolved in 7 mL of methanol and 20 mL of chloroform. The reaction mixture was cooled in an ice-bath and HCl gas was bubbled-in for 1.5 hours to saturate the solution. The mixture was sealed and allowed to stir at room
25 temperature for 14 hours. The solvents were removed *in vacuo* and the resulting solid was used in the next step.

The imidate formed above was added with 0.44 g (5.7 mmol) of ammonium carbonate and 20 mL of methanol. The mixture was allowed to stir under N₂ for 14 hours. The solvent was removed
30 at reduced pressure. The crude benzamidine was purified by HPLC (C18 reversed phase) eluting with 0.5% TFA in H₂O/CH₃CN to give 0.37 g (65%) of the desired salt. ¹H NMR (DMSO-d₆) δ: 10.63 (s, 1H); 9.42 (s, 1H); 9.03 (br s, 2H); 9.05 (br s, 2H); 8.16 (s, 1H); 8.00 (m, 1H); 7.98 (m, 1H); 7.90 (d, J = 7.3, 1H);
35 7.76 (t, J = 8.1, 1H); 7.57 (m, 4H); 7.34 (d, J = 8.1, 2H); 7.28 (m, 1H); 4.77 (s, 2H); 3.34 (s, 3H). HRMS 506.1487 (M+H).

Example 13

2-Methyl-4-(3-amidinophenyl)-5-[(2'-trifluoromethyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole, trifluoroacetic acid salt

- 5 Part A. Preparation of 2-methyl-4-(3-cyanophenyl)-5-[4-bromophenyl)aminocarbonyl]thiazole.

To a solution of 4-bromoaniline (0.21 g, 1.2 mmol) in 25 mL of methylene chloride at room temperature was added
10 trimethylaluminum (1.02 mL of a 2.0M solution in toluene, 2.04 mmol) dropwise. The reaction was stirred until gas evolution ceased and then 2-methyl-4-(3-cyanophenyl)-5-carbomethoxythiazole from Example 3, Part A (0.26 g, 1.02 mmol) was added in 10 mL of methylene chloride. The resulting
15 solution was stirred at 40° C for 16h and then was allowed to cool to 25° C. This mixture was quenched with saturated aq NH₄Cl, diluted with ethyl acetate, washed with water and brine, dried (MgSO₄) and concentrated *in vacuo*. The residue was purified by flash chromatography (elution with 4:1
20 hexanes/ethyl acetate) to afford 0.18 g (44%) of the title compound as a solid. MS (ESI) 398.0/400.0 (M+H)+.

Part B. Preparation of 2-methyl-4-(3-cyanophenyl)-5-[(2'-trifluoromethyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole.

25

To a solution of 2-methyl-4-(3-cyanophenyl)-5-[4-bromophenyl)aminocarbonyl]thiazole (0.175 g, 0.44 mmol) in 25 mL of benzene was added 2-(trifluoromethyl)phenylboronic acid (0.118 g, 0.62 mmol); tetrabutylammonium bromide (0.006 g, 0.02 mmol); sodium carbonate (0.14 g, 1.3 mmol) and 1.2 mL of
30 H₂O. This mixture was degassed with a stream of nitrogen and then tetrakis(triphenylphosphine)palladium (0.02 g, 0.02 mmol) was added and the reaction mixture was stirred at 80° C for 16 h. The mixture was allowed to cool to room temperature and
35 then was diluted with ethyl acetate, washed with H₂O and brine, dried over MgSO₄ and was concentrated *in vacuo* to afford 0.166 g (83%) of the title compound, which was sufficiently pure to be used without purification. MS (ESI) 464.2 (M+H)+.

Part C. Preparation of 2-methyl-4-(3-amidinophenyl)-5-[(2'-trifluoromethyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole, trifluoroacetic acid salt.

5

Following the procedure described in Example 7, Part F, 2-methyl-4-(3-cyanophenyl)-5-[(2'-trifluoromethyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole (0.166 g, 0.36 mmol) was converted into 45 mg (21%) of the title compound as a white solid following HPLC purification. MS (ESI) 481.3 (M+H)+.

10

Example 14

2-Phenyl-4-(3-amidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole, trifluoroacetic acid salt

15

Part A. Preparation of 2-phenyl-4-(3-cyanophenyl)-5-carbomethoxythiazole.

To a solution of methyl 2-bromo-3-(3-cyanophenyl)-3-oxopropionate from Example 7, Part C (0.51 g, 1.8 mmol) in 20 mL of absolute ethanol was added thiobenzamide (0.25 g, 1.8 mmol). The resulting mixture was stirred at 80° C for 24h. The reaction was allowed to cool and then was filtered. The solid was washed with ethanol and dried in vacuo to yield 0.53 g (91%) of the title compound. MS (ESI) 321.1 (M+H)+.

25

Part B. Preparation of 2-phenyl-4-(3-cyanophenyl)-5-[(2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole.

30

Following the procedure of Example 7, Part E, 2-phenyl-4-(3-cyanophenyl)-5-carbomethoxythiazole (0.30 g, 0.94 mmol) was converted into 0.53 g (95%) of the title compound as a solid. MS (ESI) 593.3 (M+H)+.

35

Part C. Preparation of 2-phenyl-4-(3-amidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole, trifluoroacetic acid salt.

Following the procedure of Example 9, Part C, 2-phenyl-4-(3-cyanophenyl)-5-[(2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole (0.53 g, 0.90 mmol) was converted into 61 mg of the title compound (10%) as a white powder following HPLC purification. ¹H NMR (DMSO-d₆) δ: 10.77 (s, 1H), 9.40 (broad s, 2H), 8.98 (broad s, 2H), 8.28 (broad s, 1H), 8.08 (m, 3H), 7.99 (d, 1H, J=8 Hz), 7.81 (d, 1H, J=8 Hz), 7.71 (t, 1H, J=8 Hz), 7.61-7.52 (m, 7H), 7.38-7.22 (m, 5H). MS (ESI) 554.3 (M+H)+.

Example 15

3-(3-amidinophenyl)-4-[(3-fluoro-2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]isoxazole, trifluoroacetic acid salt

Part A. Preparation of 3-fluoro-2'-methylthio-[1,1']-biphenylamine.

A benzene solution (100 mL) of 2-methylthiophenylboronic acid (2.07 g, 12.3 mmol); 4-bromo-2-fluoro aniline (1.06 g, 5.6 mmol); aq. Na₂CO₃ (12.5 mL, 2 M, 25 mmol); and tetra n-butyl ammonium bromide (90 mg, 0.3 mmol) was purged with vacuum and Ar. Bis(triphenylphosphine)palladium (II) chloride (195 mg, 0.3 mmol) was added, and the reaction was refluxed 10 h. After cooling, the reaction was diluted with EtOAc and H₂O, the layers were separated, the organic was dried over Na₂SO₄, filtered, and evaporated. The crude product was chromatographed on silica gel (10% EtOAc/hexanes) to yield the desired product (1.05 g, 81%). ¹H NMR (CDCl₃) δ: 7.29 (m, 2H); 7.18 (m, 2H); 7.09 (dd, 1H, J = 11.7, J' = 1.8); 7.01 (dd, 1H, J = 8.0, J' = 1.4); 6.82 (t, 1H, J = 9.2); 3.79 (bs, 2H); 2.38 (s, 3H).

Part B. Preparation of 3-(3-cyanophenyl)-4-[(3-fluoro-2'-methylthio-[1,1']-biphen-4-yl)aminocarbonyl]isoxazole.

Trimethyl aluminum (1.3 mL, 2.0 M in heptane, 2.6 mmol) was added to the 4-amino-3-fluoro-2'-methylthio-[1,1']-

biphenyl (309 mg, 1.3 mmol) in CH_2Cl_2 (7 mL) and stirred at room temp 25 min. Then methyl 3-(3-cyanophenyl)isoxazole-4-carboxylate isoxazole (300 mg, 1.3 mmol) was added and stirred 48 h. More trimethyl aluminum (1.3 mL, 2.6 mmol) and CH_2Cl_2 (5 mL) were added. After an additional 3 days, the reaction was quenched carefully with 1N HCl and extracted into CH_2Cl_2 . The organic layer was further washed with H_2O and brine, dried over Na_2SO_4 , filtered, and evaporated. The crude product was chromatographed on silica gel (20-30% EtOAc/hexanes followed by 2% MeOH/ CHCl_3) to yield the desired product (0.50 g, 89%). ^1H NMR (CDCl_3 , 400 MHz) δ : 9.07 (s, 1H); 8.34 (bt, 1H, $J = 8.3$); 8.13 (t, 1H, $J = 1.2$); 8.04 (dt, 1H, $J = 7.8$, $J' = 1.2$); 7.84 (dt, 1H, $J = 7.8$, $J' = 1.4$); 7.67 (t, 1H, $J = 8.1$); 7.57 (bs, 1H); 7.35 (m, 1H); 7.28 (m, 1H); 7.19 (m, 4H); 2.38 (s, 1H).

Part C. Preparation of 3-(3-cyanophenyl)-4-[(3-fluoro-2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]isoxazole.

To a chloroform solution (50 mL) of 3-(3-cyanophenyl)-4-[(3-fluoro-2'-methylthio-[1,1']-biphen-4-yl)aminocarbonyl]isoxazole (0.47 g, 1.1 mmol); m-CPBA (351 mg, 57-86%, max 1.7 mmol) was added. The resulting mixture was stirred at room temp under Ar 22 h. Additional m-CPBA (94mg of 50-60% and 348mg of 57-86%, max 2.1 mmol) was added and stirred 4 h. The reaction was extracted with sat. aq. Na_2SO_3 and sat. NaHCO_3 , dried over Na_2SO_4 , filtered, and evaporated. The crude product was chromatographed on silica gel (30-50% EtOAc/hexanes) to yield the desired sulfone (447 mg, 88%). ^1H NMR (CDCl_3) δ : 9.10 (s, 1H); 8.41 (t, 1H, $J = 8.4$); 8.22 (dd, 1H, $J = 7.7$, $J' = 1.5$); 8.12 (d, 1H, $J = 1.5$); 8.05 (dt, 1H, $J = 8.1$, $J' = 1.5$); 7.86 (dt, 1H, $J = 7.7$, $J' = 1.5$); 7.64 (m, 4H); 7.28 (m, 3H); 2.73 (s, 3H).

Part C. Preparation of 3-(3-amidinophenyl)-4-[(3-fluoro-2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]isoxazole, trifluoroacetic acid salt.

Solid 3-(3-cyanophenyl)-4-[(3-fluoro-2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]isoxazole (423 mg, 0.92 mmol) was suspended in methanol (15 mL) and cooled to 0°C. HCl (g) was generated by the addition of conc. H₂SO₄ (90 mL) into solid NaCl (360 g) over 1 h. The HCl (g) continued bubbling through the reaction an additional 90 min, at which time the generator and ice bath were removed. The reaction stirred under argon 20 h and was evaporated. After a few hours under high vacuum, the reaction was redissolved in methanol (15 mL); and ammonium carbonate (440 mg, 4.6 mmol) was added. The reaction was stirred 22 h and evaporated. The crude product was purified by prep HPLC on a C-18 reverse phase column (20-80% MeCN/H₂O/0.05% TFA) to yield a white solid (0.14 g, 26%). ¹H NMR (DMSO-d₆) δ: 10.47 (s, 1H); 9.68 (s, 1H); 9.40 (s, 1.5H); 9.01 (s, 1.5 H); 8.16 (s, 1H); 8.06 (m, 2H); 7.91 (d, 1H, J = 8.1); 7.72 (m, 4H); 7.37 (m, 2H); 7.21 (d, 1H, J = 8.1); 2.90 (s, 3H). HRMS calc. for C₂₄H₂₀FN₄O₄S, 479.1189; found, 479.1169.

Example 16

3-(3-amidinophenyl)-4-[(2'-trifluoromethylthio-[1,1']-biphen-4-yl)aminocarbonyl]isoxazole, trifluoroacetic acid salt

Part A. Preparation of 3-(3-cyanophenyl)-4-[(2'-trifluoromethylthio-[1,1']-biphen-4-yl)aminocarbonyl]isoxazole.

Trimethyl aluminum (2.6 mL, 2.0 M in heptane, 5.2 mmol) was added to the 4-amino-2'-trifluoromethylthio-[1,1']-biphenyl (361 mg, 1.3 mmol); in CH₂Cl₂ (5 mL) and stirred at room temp 8 min. A CH₂Cl₂ solution (5 mL) of methyl 3-(3-cyanophenyl)isoxazole-4-carboxylate (300 mg, 1.3 mmol) was added and stirred 2 days. No further reaction was observed after adding more trimethyl aluminum (650 mL) and stirring an additional 20 hr. The reaction was quenched carefully with 1 M HCl and extracted into CH₂Cl₂. The organic was extracted again with water and brine, dried over Na₂SO₄, filtered, and evaporated. The crude mixture was chromatographed on silica gel (30% EtOAc/hexanes) to yield a yellow solid (565 mg, 92%).

¹H NMR (CDCl₃) δ: 9.02 (s, 1H); 8.17 (s, 1H); 8.09 (d, 1H, J = 8.1); 7.82 (t, 2H, J = 8.1); 7.66 (t, 1H, J = 7.7); 7.53 (m, 4H); 7.42 (t, 2H, J = 7.7); 7.32 (d, 2H, J = 8.4).

- 5 Part B. Preparation of 3-(3-amidinophenyl)-4-[(2'-trifluoromethylthio-[1,1']-biphen-4-yl)aminocarbonyl]isoxazole.

A methanol solution (10 mL) of 3-(3-cyanophenyl)-4-[(2'-trifluoromethylthio-[1,1']-biphen-4-yl)aminocarbonyl]isoxazole
10 (137 mg, 0.29 mmol) was cooled to 0° C. HCl was generated by the slow addition of conc. H₂SO₄ (60 mL) to solid NaCl (240 g) over 1 h. The HCl thus generated was bubbled into the reaction mixture over 2 h. The generator and ice bath were removed, and the reaction stirred under Ar 16 h. The reaction
15 was evaporated, placed briefly under high vacuum, and redissolved in methanol (10 mL). Ammonium carbonate (138 mg, 1.4 mmol) was added. After stirring 19 h, the reaction was evaporated and purified by prep HPLC on a C-18 reverse phase column (20-80% MeCN/H₂O/0.05% TFA) to yield a white powder (84
20 mg, 48%). ¹H NMR (DMSO-d₆) δ: 10.66 (s, 1H); 9.66 (s, 1H); 9.41 (s, 2H); 8.98 (s, 2H); 8.18 (s, 1H); 8.06 (d, 1H, J = 7.6); 7.91 (d, 1H, J = 8.5); 7.82 (d, 1H, J = 6.9); 7.68 (m, 4H); 7.50 (m, 2H); 7.35 (d, 2H, J = 8.8). ¹⁹F NMR (DMSO-d₆) -42.45, -73.86. HRMS calc. for C₂₄H₁₈F₃N₄O₂S, 483.1103; found, 483.1101.

25

Example 17

3-(3-amidinophenyl)-5-amino-4-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]isoxazole, trifluoroacetic acid salt

- 30 Part A. Preparation of methyl N-(2'-t-butylaminosulfonyl-[1,1']-biphen-4-yl)cyanoacetamide.

Added to the 4-amino-2'-t-butylaminosulfonyl-[1,1']-biphenyl (2.0 g, 6.6 mmol); in CH₂Cl₂ (10 mL) and stirred at
35 room temp 30 min. A CH₂Cl₂ solution (5 mL) of methyl cyanoacetate (0.58 mL, 6.6 mmol) was added and stirred 1 day. The reaction was quenched carefully with 1 M HCl and extracted into CH₂Cl₂. The organic was extracted again with water and

brine, dried over MgSO_4 , filtered, and evaporated. The crude mixture was chromatographed on silica gel (50% EtOAc/hexanes) to yield a yellow solid (0.81 g, 33%). ^1H NMR (CDCl_3) δ : 8.74 (s, 1H); 8.16 (d, $J = 7.7$, 1H); 7.64 (d, $J = 8.8$, 2H); 7.58 (d, $J = 7.3$, 1H); 7.66 (d, $J = 8.1$, 1H); 7.46 (d, $J = 8.4$, 2H); 7.31 (d, $J = 7.7$, 1H); 3.89 (s, 1H); 3.63 (s, 2H); 1.04 (s, 9H). MS ($\text{NH}_3\text{-CI}$) m/z 389 ($\text{M}+\text{NH}_3$).

Part B. Preparation of 3-(3-cyanophenyl)-5-amino-4-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]isoxazole, trifluoroacetate salt.

Methyl N-(2'-t-butylaminosulfonyl-[1,1']-biphen-4-yl)cyanacetamide (0.81 g, 2.18 mmol) was added to a solution of triethylamine (0.33 g, 3.27 mmol) in 25 mL of ethanol and 5 mL of CH_2Cl_2 . A solution of 3-cyanobenzenehydroximinoyl chloride (0.39 g, 2.18 mmol) in 10 mL ethanol was added slowly over 12 hours via a syringe pump. The reaction mixture was diluted with 50 mL of ether and washed three times with 10 mL of water and twice with 10 mL saturated NaHCO_3 , then dried with MgSO_4 and concentrated *in vacuo*. The resulting off-white solid was purified by prep HPLC on a C-18 reverse phase column (30-100% MeCN/ H_2O /0.05% TFA) to yield a white powder (0.10 g, 8.3%). ^1H NMR (CDCl_3) δ : 9.09 (s, 1H); 8.03 (s, 1H); 7.99 (d, $J = 6.2$, 1H); 7.92 (t, $J = 7.7$, 2H); 7.72 (br s, 2H); 7.59 (t, $J = 7.7$, 1H); 7.53 (m, 2H); 7.45 (d, $J = 8.4$, 2H); 7.26 (d, $J = 8.4$, 2H); 7.25 (m, 1H); 6.51 (s, 1H), 0.97 (s, 9H). MS ($\text{NH}_3\text{-CI}$) m/z 273.0 ($\text{M}+\text{H}$).

Part C. Preparation of 3-(3-amidinophenyl)-5-amino-4-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]isoxazole, trifluoroacetate salt.

3-(3-cyanophenyl)-5-amino-4-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]isoxazole (0.10 g, 0.19 mmol) was dissolved in 1 mL of methanol and 4 mL of chloroform. The reaction mixture was cooled in an ice-bath and HCl gas was bubbled-in for 10 minutes to saturate the solution. The

mixture was sealed and allowed to stir at room temperature for 14 hours. The solvents were removed in vacuo and the resulting solid was used in the next step.

The imidate formed above was added with 0.07 g (0.95 mmol) of ammonium carbonate and 10 mL of methanol. The mixture was allowed to stir under N₂ for 14 hours. The solvent was removed at reduced pressure. The crude benzamidine was purified by HPLC (C18 reversed phase) eluting with 0.5% TFA in H₂O/CH₃CN to give 0.0064 g (7%) of the desired salt. ¹H NMR (DMSO-d₆) δ: 9.39 (br s, 2H); 9.12 (s, 1H); 8.96 (br s, 2H); 8.06 (s, 1H); 7.98 (d, J = 7.7, 1H); 7.96 (d, J = 7.8, 1H); 7.88 (d, J = 7.7, 1H); 7.74 (br s, 2H); 7.70 (t, J = 7.8, 1H); 7.54 (m, 3H); 7.42 (d, J = 8.8, 2H); 7.26 (d, J = 8.8, 2H); 7.25 (m, 1H); 7.21 (br s, 2H); 6.51 (s, 1H). HRMS m/z 477.1338 (M+H).

Example 18

2-(Phenylamino)-4-(3-amidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole, trifluoroacetic acid salt.

Part A. Preparation of ethyl 3-(3-cyanophenyl)-3-oxopropionate.

To a suspension of sodium hydride (1.2 g of 60% suspension in mineral oil, hexane-washed, 30.3 mmol) in 40 mL of tetrahydrofuran was added diethyl carbonate (3.7 mL, 30.3 mmol) and 3-acetyl benzonitrile (2.2 g, 15.2 mmol). The resulting suspension was stirred at 65° C for 1 h and then was cooled to room temperature. There was added 40 mL of 10% aqueous HCl and the reaction mixture was diluted with ethyl acetate and the layers were separated. The organic layer was washed with brine, dried (MgSO₄) and concentrated in vacuo to afford 3.2 g (96%) of the title compound, which was sufficiently pure to be used without purification. MS (NH₃-Cl) 218.3 (M+H)+.

Part B. Preparation of ethyl 2-bromo-3-(3-cyanophenyl)-3-oxopropionate.

According to the procedure of Example 7, Part C, ethyl 3-(3-cyanophenyl)-3-oxopropionate (3.2 g, 14.7 mmol) was converted into the crude bromide, which was purified by flash chromatography (elution with 4:1 hexanes/ethyl acetate) to afford 2.1 g (48%) of the title compound. MS (H₂O, GC/MS) 296/298 (M+H)+.

Part C. Preparation of 2-(phenylamino)-4-(3-cyanophenyl)-5-carboethoxythiazole.

To a solution of ethyl 2-bromo-3-(3-cyanophenyl)-3-oxopropionate (0.60 g, 2.03 mmol) in 20 mL of absolute ethanol was added N-phenylthiourea (0.31 g, 2.03 mmol). The resulting mixture was stirred at 80° C for 3h. The reaction was allowed to cool and the solvent was evaporated *in vacuo*. The residue was taken up in ethyl acetate, washed with saturated aq NaHCO₃ and brine, dried (MgSO₄) and concentrated *in vacuo* to yield a solid. Trituration with hexanes/ethyl ether left the title compound as an off-white solid (0.35 g, 49%). MS (NH₃-CI) 350 (M+H)+.

Part D. Preparation of 2-(phenylamino)-4-(3-cyanophenyl)-5-[(2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole.

Following the procedure of Example 7, Part E, 2-(phenylamino)-4-(3-cyanophenyl)-5-carboethoxythiazole (0.29 g, 0.83 mmol) was converted into 0.37 g (74%) of the title compound as a solid. MS (ESI) 608.3 (M+H)+.

Part E. Preparation of 2-(phenylamino)-4-(3-amidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole, trifluoroacetic acid salt.

Following the procedure of Example 9, Part C, 2-(phenylamino)-4-(3-cyanophenyl)-5-[(2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole (350 mg, 0.58 mmol) was converted into 50 mg of the title compound (12%) as an off-white powder following HPLC purification. ¹H NMR (DMSO-d₆) δ: 10.70 (s, 1H), 10.25 (s, 1H), 9.41 (broad s, 2H), 9.02 (broad s, 2H), 8.19 (m, 1H), 8.08 (d, 1H, J=7.7 Hz), 8.03 (d, 1H, J=8.0 Hz), 7.73-7.54 (m, 8H), 7.41-7.27 (m, 8H). MS (ESI) 569.0 (M+H)+.

Example 19

2-(Benzylamino)-4-(3-amidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole, trifluoroacetic acid salt.

Part A. Preparation of 2-(benzylamino)-4-(3-cyanophenyl)-5-carboethoxythiazole.

To a solution of ethyl 2-bromo-3-(3-cyanophenyl)-3-oxopropionate (0.60 g, 2.03 mmol) in 20 mL of absolute ethanol was added N-benzylthiourea (0.34 g, 2.03 mmol). The resulting mixture was stirred at 80° C for 3h. The reaction was allowed to cool and the solvent was evaporated *in vacuo*. The residue was taken up in ethyl acetate, washed with saturated aq NaHCO₃ and brine, dried (MgSO₄) and concentrated *in vacuo* to yield a solid. Trituration with hexanes/ethyl ether left the title compound as an off-white solid (0.36 g, 49%). MS (ESI) 364.1 (M+H)+.

Part B. Preparation of 2-(benzylamino)-4-(3-cyanophenyl)-5-[(2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole.

Following the procedure of Example 7, Part E, 2-(benzylamino)-4-(3-cyanophenyl)-5-carboethoxythiazole (0.27 g, 0.74 mmol) was converted into 0.30 g (65%) of the title compound as a yellowwash solid. MS (ESI) 622.3 (M+H)+.

Part C. Preparation of 2-(benzylamino)-4-(3-amidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole, trifluoroacetic acid salt.

5 Following the procedure of Example 9, Part C, 2-(benzylamino)-4-(3-cyanophenyl)-5-[(2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole (260 mg, 0.42 mmol) was converted into 95 mg of the title compound (33%) as an off-white powder following HPLC
10 purification. ¹H NMR (DMSO-d₆) δ: 9.97 (s, 1H), 9.36 (broad s, 2H), 8.98 (broad s, 2H), 8.78 (t, 1H, J=5.9 Hz), 8.09 (broad s, 1H), 8.02 (dd, 1H, J=7.8, 1.6 Hz), 7.98 (d, 1H, J=7.8 Hz), 7.76 (d, 1H, J=8.0 Hz), 7.66-7.25 (m, 15H), 4.58 (d, 2H, J=5.9 Hz). MS (ESI) 583.0 (M+H)+.

15 **Examples 20 and 21**

2-(methylamino)-4-(3-amidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole, trifluoroacetic acid salt (Example 20) and 2-(methylamino)-4-(3-carboxamidophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole (Example 21)

Part A. Preparation of 2-(methylamino)-4-(3-cyanophenyl)-5-carboethoxythiazole.

25 To a solution of ethyl 2-bromo-3-(3-cyanophenyl)-3-oxopropionate (0.65 g, 2.2 mmol) in 20 mL of tetrahydrofuran was added N-methylthiourea (0.20 g, 2.2 mmol). The resulting mixture was stirred at 65° C for 16h. The reaction was allowed
30 to cool and the solvent was evaporated *in vacuo*. The residue was taken up in ethyl acetate, washed with saturated aq NaHCO₃ and brine, dried (MgSO₄) and concentrated *in vacuo* to yield a solid. Trituration with hexanes/ethyl ether left the title compound as an off-white solid (0.46 g, 73%). MS (ESI) 288.3
35 (M+H)+.

Part B. Preparation of 2-(methylamino)-4-(3-cyanophenyl)-5-[(2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole.

5 Following the procedure of Example 7, Part E, 2-(methylamino)-4-(3-cyanophenyl)-5-carboethoxythiazole (0.46 g, 1.6 mmol) was converted into 0.68 g (78%) of the title compound as a yellowish solid. MS (ESI) 546.7 (M+H)+.

10 Part C. Preparation of 2-(methylamino)-4-(3-amidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole, trifluoroacetic acid salt (Example 6) and 2-(methylamino)-4-(3-carboxamidophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole (Example 20).

15 Following the procedure of Example 9, Part C, 2-(methylamino)-4-(3-cyanophenyl)-5-[(2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole (500 mg, 1.0 mmol) was converted into 85 mg of the title
20 compound (13%), 2-(methylamino)-4-(3-amidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole, trifluoroacetic acid salt (Example 20) as a white powder following HPLC purification. ¹H NMR (DMSO-d₆) δ: 9.89 (s, 1H), 9.33 (broad s, 2H), 8.96 (broad s, 2H), 8.05 (broad s, 1H),
25 7.95 (m, 2H), 7.72 (d, 1H, J=8.0 Hz), 7.62-7.44 (m, 6H), 7.29-7.21 (m, 5H), 2.89 (d, 3H). MS (ESI) 507.2 (M+H)+. There was also isolated 50 mg (8%) of 2-(methylamino)-4-(3-carboxamidophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole (Example 21). MS (ESI) 508.1 (M+H)+.

30

Examples 22 and 23

2-methyl-4-(3-amidinophenyl)-5-[[5-(2'-aminosulfonylphenyl-1-yl)pyridin-2-yl]aminocarbonyl]thiazole, trifluoroacetic acid salt (Example 22) and 2-methyl-4-(3-carboxamidophenyl)-5-[[5-(2'-aminosulfonylphenyl-1-yl)pyridin-2-yl]aminocarbonyl]thiazole (Example 23)

35

Part A. Preparation of 2-methyl-4-(3-cyanophenyl)-5-carboxythiazole.

To a solution of 2-methyl-4-(3-cyanophenyl)-5-carbomethoxythiazole from Example 9, Part A (0.96 g, 3.7 mmol) in 20 mL of tetrahydrofuran and 10 mL of water was added lithium hydroxide monohydrate (0.31 g, 7.4 mmol). The resulting mixture was stirred at room temperature for 16 h. The mixture was concentrated *in vacuo*, diluted with H₂O and saturated aqueous NaHCO₃ and extracted with hexane. The organic layer was discarded and the aqueous layer was acidified and extracted twice with ethyl acetate. The combined ethyl acetate extracts were washed with brine, dried over MgSO₄ and concentrated *in vacuo* to afford 0.90 g (99%) of the title compound, which was sufficiently pure to be used without purification. MS (NH₃-CI) 245 (M+H)+.

Part B. Preparation of 2-methyl-4-(3-cyanophenyl)-5-[[5-(2'-tert-butylaminosulfonylphenyl-1-yl)pyridin-2-yl]aminocarbonyl]thiazole.

To a solution of 2-methyl-4-(3-cyanophenyl)-5-carboxythiazole (0.22 g, 0.89 mmol) in 10 mL of acetonitrile was added thionyl chloride (0.60 g, 5.0 mmol) and 2 drops of dimethylformamide. The resulting solution was allowed to stir at 50°C for 10 min and then at room temperature for 1 h. The solution was concentrated *in vacuo*, the residue was dissolved in 20 mL of methylene chloride and then [[5-(2'-tert-butylaminosulfonylphenyl-1-yl)pyridin-2-yl]amine (0.30 g, 0.98 mmol) and triethylamine (1.3 mL, 8.9 mmol) were added. The reaction mixture was allowed to stir at 25° C for 16 h. The reaction was diluted with ethyl acetate, washed with 10% aq HCl, saturated aq NaHCO₃ and brine, dried (MgSO₄); filtered through a pad of silica gel and concentrated *in vacuo*. The residue was purified by flash chromatography (elution with 3:1 hexanes/ethyl acetate) to afford 0.07 g (15%) of the title compound. MS (ESI) 532.2 (M+H)+.

Part C. Preparation of 2-methyl-4-(3-amidinophenyl)-5-[[5-(2'-aminosulfonylphenyl-1-yl)pyridin-2-yl]aminocarbonyl]thiazole,

trifluoroacetic acid salt (Example 22) and 2-methyl-4-(3-carboxamidophenyl)-5-[[5-(2'-aminosulfonylphenyl-1-yl)pyridin-2-yl]aminocarbonyl]thiazole (Example 23).

5 Following the procedure described in Example 9, Part C, 2-methyl-4-(3-cyanophenyl)-5-[[5-(2'-tert-butylaminosulfonylphenyl-1-yl)pyridin-2-yl]aminocarbonyl]thiazole (0.07 g, 0.14 mmol) was converted into 10 mg (16%) of 2-methyl-4-(3-amidinophenyl)-5-[[5-(2'-aminosulfonylphenyl-1-yl)pyridin-2-yl]aminocarbonyl]thiazole, 10 trifluoroacetic acid salt (Example 22) following HPLC purification. MS (ESI) 493.1 (M+H)+. There was also isolated 25 mg (29%) of 2-methyl-4-(3-carboxamidophenyl)-5-[[5-(2'-aminosulfonylphenyl-1-yl)pyridin-2-yl]aminocarbonyl]thiazole (Example 23). MS (ESI) 494.1 (M+H)+.

15

Examples 24 and 25

2-(3-pyridyl)-4-(3-amidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole, trifluoroacetic acid salt (Example 24) and 2-(3-pyridyl)-4-(3-carboxamidophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole (Example 25)

20

Part A. Preparation of 2-(3-pyridyl)-4-(3-cyanophenyl)-5-carboethoxythiazole.

25

To a solution of ethyl 2-bromo-3-(3-cyanophenyl)-3-oxopropionate (1.0 g, 3.4 mmol) in 20 mL of tetrahydrofuran was added thionicotinamide (0.46 g, 3.4 mmol). The resulting mixture was stirred at 65° C for 16h. The reaction was allowed 30 to cool and the solvent was evaporated *in vacuo*. The residue was triturated with chloroform, taken up in ethyl acetate, washed with saturated aq Na₂CO₃ and brine, dried (MgSO₄) and concentrated *in vacuo* to yield a solid. Trituration with ethyl acetate left the title compound as an off-white solid 35 (0.26 g, 23%). MS (ESI) 336.1 (M+H)+.

Part B. Preparation of 2-(3-pyridyl)-4-(3-cyanophenyl)-5-[(2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole.

5 4-(3-cyanophenyl)-5-carboethoxythiazole (0.26 g, 0.77 mmol) was converted into 0.24 g (52%) of the title compound as a yellowish solid. MS (ESI) 594.1 (M+H)+.

10 Part C. Preparation of 2-(3-pyridyl)-4-(3-amidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole, trifluoroacetic acid salt (Example 24) and 2-(3-pyridyl)-4-(3-carboxamidophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole (Example 25).

15 Following the procedure of Example 9, Part C, 2-(3-pyridyl)-4-(3-cyanophenyl)-5-[(2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole (0.24 g, 0.45 mmol) was converted into 80 mg of the title compound (27%) of 2-(3-pyridyl)-4-(3-amidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole, trifluoroacetic acid salt (Example 24) as a white powder following HPLC purification. ¹H NMR (DMSO-d₆) δ: 10.82 (s, 1H), 9.41 (broad s, 2H), 9.30 (broad s, 1H), 9.02 (broad s, 2H), 8.75 (d, 1H, J=5.5 Hz), 8.43 (d, 1H, J=8 Hz), 8.30 (broad s, 1H), 8.10 (d, 1H, J=8 Hz), 7.98 (d, 1H, J=8 Hz), 7.82 (d, 1H, J=8 Hz), 7.70 (t, 1H, J=8 Hz), 7.62-7.50 (m, 5H), 7.38-7.22 (m, 5H). MS (ESI) 555.0 (M+H)+. There was also isolated 30 mg (10%) of 2-(3-pyridyl)-4-(3-carboxamidophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole (Example 25). ¹H NMR (DMSO-d₆) δ: 10.72 (s, 1H), 9.28 (broad s, 1H), 8.75 (broad s, 1H), 8.42 (m, 2H), 8.09 (broad s, 1H), 7.98 (d, 1H, J=8 Hz), 7.90 (m, 2H), 7.62-7.50 (m, 6H), 7.42 (broad s, 1H), 7.37-7.26 (m, 3H), 7.21 (broad s, 2H). MS (ESI) 578.0 (M+Na)+.

35

Examples 26 and 27

2-chloro-4-(3-amidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole, trifluoroacetic acid salt (Example 26) and 2-chloro-4-(3-carboxamidophenyl)-5-[(2'-

aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole
(Example 27)

Part A. Preparation of 2-amino-4-(3-cyanophenyl)-5-carboethoxythiazole.

To a solution of ethyl 2-bromo-3-(3-cyanophenyl)-3-oxopropionate (4.0 g, 13.5 mmol) in 100 mL of tetrahydrofuran was added thiourea (1.03 g, 13.5 mmol). The resulting mixture was stirred at 65° C for 16h. The reaction was allowed to cool and the solvent was evaporated *in vacuo*. The residue was triturated with ether, taken up in ethyl acetate, washed with saturated aq Na₂CO₃ and brine, dried (MgSO₄) and concentrated *in vacuo* to yield a solid. Trituration with ethyl ether left the title compound as an off-white solid (3.79 g, 98%).

Part B. Preparation of 2-chloro-4-(3-cyanophenyl)-5-carboethoxythiazole.

To a suspension of anhydrous copper (II) chloride (1.86 g, 13.9 mmol) in 180 mL of acetonitrile was added *tert*-butyl nitrite (1.43 g, 1.39 mmol). The solution was warmed to reflux and then 2-amino-4-(3-cyanophenyl)-5-carboethoxythiazole (3.79 g, 13.9 mmol) in 50 mL of acetonitrile was added via addition funnel over 5 min. The mixture was stirred at reflux until gas evolution ceased (about 30 min). The reaction was cooled to room temperature, poured into 10% aq HCl and extracted with ether. The ether layer was washed with saturated aq NaHCO₃ and brine, dried (MgSO₄) and concentrated to afford 3.44 g (82%) of the title compound which was used without purification. MS (NH₃-CI) 293.2 (M+H)+.

Part C. Preparation of 2-chloro-4-(3-cyanophenyl)-5-[(2'-*tert*-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole.

Cyanophenyl)-5-carboethoxythiazole (0.084 g, 0.3 mmol) was converted into 0.056 g (34%) of the title compound as a solid. MS (ESI) 551.0 (M+H)+.

- 5 Part D. Preparation of 2-chloro-4-(3-amidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole, trifluoroacetic acid salt (Example 26) and 2-chloro-4-(3-carboxamidophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole (Example 27).

10

- Following the procedure of Example 9, Part C, 2-chloro-4-(3-cyanophenyl)-5-[(2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole (0.056 g, 0.11 mmol) was converted into 15 mg of the title compound (23%) of 2-chloro-4-(3-amidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole, trifluoroacetic acid salt (Example 26) as a white powder following HPLC purification. ¹H NMR (DMSO-d₆) δ: 10.77 (s, 1H), 9.40 (broad s, 2H), 8.96 (broad s, 2H), 8.15 (broad s, 1H), 7.98 (m, 2H), 7.80 (d, 1H, J=8 Hz), 7.69 (d, 1H, J=8 Hz), 7.61-7.52 (m, 4H), 7.37-7.23 (m, 5H). MS (ESI) 512.0 (M+H)+. There was also isolated 20 mg (31%) of 2-chloro-4-(3-carboxamidophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole (Example 27). ¹H NMR (DMSO-d₆) δ: 10.68 (s, 1H), 8.28 (broad s, 1H), 8.05 (broad s, 1H), 7.98 (d, 1H, J=8 Hz), 7.89 (d, 1H, J=8 Hz), 7.78 (d, 1H, J=8 Hz), 7.61-7.52 (m, 5H), 7.40 (broad s, 1H), 7.37-7.23 (m, 3H), 7.21 (broad s, 2H). MS (ESI) 534.9 (M+Na)+.

Examples 28, 29 and 30

- 30 2-chloro-4-(3-amidinophenyl)-5-[[5-(2'-aminosulfonylphenyl-1-yl)pyridin-2-yl]aminocarbonyl]thiazole, trifluoroacetic acid salt (Example 28), 2-chloro-4-(3-carboxamidophenyl)-5-[[5-(2'-aminosulfonylphenyl-1-yl)pyridin-2-yl]aminocarbonyl]thiazole (Example 29), and 2-hydroxy-4-(3-amidinophenyl)-5-[[5-(2'-aminosulfonylphenyl-1-yl)pyridin-2-yl]aminocarbonyl]thiazole, trifluoroacetic acid salt (Example 30)

Part A. Preparation of 2-chloro-4-(3-cyanophenyl)-5-carboxythiazole.

To a solution of 2-chloro-4-(3-cyanophenyl)-5-carboethoxythiazole (0.44 g, 1.45 mmol) in 25 mL of methanol and 25 mL of water was added potassium hydroxide (0.09 g, 1.6 mmol). The resulting solution was stirred at reflux for 2h and then was cooled to room temperature. The methanol was removed in vacuo and the aqueous layer was diluted with water and washed with hexanes. The hexane layer was discarded. The aqueous layer was acidified and extracted with ethyl acetate. The ethyl acetate layer was washed with brine, dried (MgSO₄) and concentrated to yield 0.33g (97%) of the title compound which was used without further purification. MS (ESI)- 262.9 (M-H)-.

Part B. Preparation of 2-chloro-4-(3-cyanophenyl)-5-[[5-(2'-tert-butylaminosulfonylphenyl-1-yl)pyridin-2-yl]aminocarbonyl]thiazole.

Following the procedure described in Example 22, Part B, 2-chloro-4-(3-cyanophenyl)-5-carboxythiazole (0.35 g, 1.33 mmol) was converted into 0.20 g (27%) of the title compound. MS (ESI) 552.0 (M+H)+.

Part C. Preparation of 2-chloro-4-(3-amidinophenyl)-5-[[5-(2'-aminosulfonylphenyl-1-yl)pyridin-2-yl]aminocarbonyl]thiazole, trifluoroacetic acid salt (Example 28) and 2-chloro-4-(3-carboxamidophenyl)-5-[[5-(2'-aminosulfonylphenyl-1-yl)pyridin-2-yl]aminocarbonyl]thiazole, trifluoroacetic acid salt (Example 29) and 2-hydroxy-4-(3-amidinophenyl)-5-[[5-(2'-aminosulfonylphenyl-1-yl)pyridin-2-yl]aminocarbonyl]thiazole, trifluoroacetic acid salt (Example 30).

Following the procedure described in Example 3, Part C, 2-chloro-4-(3-cyanophenyl)-5-[[5-(2'-tert-butylaminosulfonylphenyl-1-yl)pyridin-2-

yl]aminocarbonyl]thiazole (0.20 g, 0.4 mmol) was converted into 75 mg (32%) of 2-chloro-4-(3-amidinophenyl)-5-[[5-(2'-aminosulfonylphenyl-1-yl)pyridin-2-yl]aminocarbonyl]thiazole, trifluoroacetic acid salt (Example 28) following HPLC purification. ¹H NMR (DMSO-d₆) δ: 9.38 (broad s, 2H), 8.98 (broad s, 2H), 8.28 (d, 1H, J=1.5 Hz), 8.15 (broad s, 1H), 8.03-7.96 (m, 3H), 7.81 (m, 2H), 7.70-7.57 (m, 4H), 7.40 (broad s, 1H), 7.36 (m, 2H). MS (ESI) 513.0 (M+H)⁺. There was also isolated 2-chloro-4-(3-carboxamidophenyl)-5-[[5-(2'-aminosulfonylphenyl-1-yl)pyridin-2-yl]aminocarbonyl]thiazole (Example 29). MS (ESI) 513.9 (M+H)⁺. There was also isolated 2-hydroxy-4-(3-amidinophenyl)-5-[[5-(2'-aminosulfonylphenyl-1-yl)pyridin-2-yl]aminocarbonyl]thiazole, trifluoroacetic acid salt (Example 30). MS (ESI) 495.0 (M+H)⁺.

Example 31

2-Chloro-4-(3-aminophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole, trifluoroacetic acid salt

Part A. Preparation of ethyl 3-(3-nitrophenyl)-3-oxopropionate.

To a suspension of anhydrous tin (II) chloride (2.5 g, 13.2 mmol) in 150 mL of methylene chloride was added ethyl diazoacetate (8.3 g, 72.8 mmol). Then 3-nitrobenzaldehyde (10.0 g, 66.2 mmol) was added as a solid in small portions over 30 min. The resulting suspension was stirred at room temperature for 24 h. Additional tin (II) chloride (2.5 g) was added and the reaction was stirred an additional 24 h. The reaction was concentrated in vacuo, diluted with ethyl acetate, washed with water (2 times) and brine, dried (MgSO₄) and concentrated. The residue was purified by flash chromatography (elution with 4:1 hexanes/ethyl acetate) to afford 5 g (32%) of the title compound. ¹H NMR (CDCl₃) (approximately 12:1 mixture of enol and keto tautomers, data for enol only) δ: 12.6 (s, 1H); 8.60 (t, 1H, J=1.8 Hz); 8.3 (m, 1H); 8.1 (m, 1H); 7.62 (t, 1H, J=7.9 Hz) 5.77 (s, 1H); 4.30 (q, 2H, J=7.2 Hz); 1.35 (t, 3H, J=7.2 Hz).

Part B. Preparation of 2-amino-4-(3-nitrophenyl)-5-carboethoxythiazole.

5 To a solution of ethyl 3-(3-nitrophenyl)-3-oxopropionate (3.45 g, 14.5 mmol) in 100 mL of acetonitrile was added hydroxy(tosyloxy)iodobenzene (6.3 g, 16.0 mmol). The resulting suspension was stirred at 65° C for 1h at which time the reaction was a homogeneous solution. Thiourea (1.22 g, 10 16.0 mmol) was added and stirring was continued at 65° C for 2 h. The mixture was cooled and concentrated, and the residue was taken up in ethyl acetate, washed with saturated aq Na₂CO₃ and brine, dried (MgSO₄) and concentrated. The residue was triturated with ethyl ether to afford 3.0 g (71%) of the title 15 compound as a yellow solid. ¹H NMR (DMSO-d₆) δ: 8.47 (t, 1H, J=1.9 Hz); 8.24 (m, 1H); 8.21 (m, 1H); 7.97 (broad s, 2H); 7.65 (t, 1H, J=8.1 Hz); 4.08 (q, 2H, J=7.1 Hz); 1.11 (t, 3H, J=7.1 Hz).

20 Part C. Preparation of 2-amino-4-(3-nitrophenyl)-5-[(2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole.

Following the procedure of Example 7, Part E, 2-amino-4-(3-nitrophenyl)-5-carboethoxythiazole (0.30 g, 1.02 mmol) was 25 converted into 0.22 g (39%) of the title compound as a solid. MS (ESI) 574.0 (M+Na)+.

Part D. Preparation of 2-chloro-4-(3-nitrophenyl)-5-[(2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole. 30

Following the procedure described in Example 26, Part B, 2-amino-4-(3-nitrophenyl)-5-[(2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole (155 mg, 0.28 mmol) 35 was converted into 150 mg (94%) of the title compound which was used without purification. MS (NH₃-CI) 588 (M+NH₄)⁺.

Part E. Preparation of 2-chloro-4-(3-aminophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole, trifluoroacetic acid salt.

5 To a solution of 2-chloro-4-(3-nitrophenyl)-5-[(2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole (100 mg, 0.18 mmol) in ethyl acetate was added tin (II) chloride dihydrate (0.32 g, 1.4 mmol). The resulting suspension was stirred at reflux for 2 h and then was cooled and quenched with saturated aq NaHCO₃. The reaction was
10 diluted with ethyl acetate, washed with brine, dried (MgSO₄) and concentrated to yield 90 mg (93%) of an amine which was used without purification. The residue was taken up in 5 mL of trifluoroacetic acid and stirred at reflux for 15 min. The
15 reaction was concentrated and the residue was purified by prep HPLC to afford 40 mg (37%) of the title compound as a white powder. ¹H NMR (DMSO-d₆) δ: 10.67 (s, 1H), 7.98 (d, 1H, J=8 Hz), 7.60-7.50 (m, 4H), 7.34-7.19 (m, 7H), 7.11 (broad m, 1H), 6.84 (broad m, 1H). MS (ESI) 484.9 (M+H)+.

20

Example 32

2-amino-4-[(3-amino-4-chloro)phenyl]-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole, trifluoroacetic acid salt

25

Part A. Preparation of ethyl 3-[(3-nitro-4-chloro)phenyl]-3-oxopropionate.

Following the procedure described in Example 31, Part A,
30 4-chloro-3-nitrobenzaldehyde (10.0 g, 53.9 mmol) was converted into 4.8 g (33%) of the title compound as a yellow solid. ¹H NMR (CDCl₃) (approximately 15:1 mixture of enol and keto tautomers, data for enol only) δ: 12.6 (s, 1H); 8.25 (d, 1H); 7.9 (dd, 1H); 7.6 (d, 1H) 5.7 (s, 1H); 4.27 (q, 2H); 1.35 (t, 3H).
35

Part B. Preparation of 2-amino-4-[(3-nitro-4-chloro)phenyl]-5-carboethoxythiazole.

Following the procedure described in Example 31, Part B, ethyl 3-[(3-nitro-4-chloro)phenyl]-3-oxopropionate (1.6 g, 5.9 mmol) was converted into the title compound as a yellow solid.
5 ¹H NMR (DMSO-d₆) δ: 8.32 (d, 1H); 7.98 (s, 2H); 7.95 (d, 1H); 7.75 (d, 1H); 4.08 (q, 2H); 1.13 (t, 3H).

Part C. Preparation of 2-amino-4-[(3-nitro-4-chloro)phenyl]-5-[(2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole.
10

Following the procedure of Example 7, Part E, 2-amino-4-[(3-nitro-4-chloro)phenyl]-5-carboethoxythiazole (0.49 g, 1.5 mmol) was converted into 0.79 g (89%) of the title compound as
15 a solid. MS (ESI) 586.0 (M+H)+.

Part D. Preparation of 2-amino-4-[(3-amino-4-chloro)phenyl]-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole, trifluoroacetic acid salt.
20

Following the procedure described in Example 31, Part E, 2-amino-4-[(3-nitro-4-chloro)phenyl]-5-[(2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole (100 mg, 0.17 mmol) was converted into 35 mg (41%) of the
25 title compound which was a white powder following HPLC purification. ¹H NMR (DMSO-d₆) δ: 9.68 (s, 1H), 7.98 (d, 1H, J=8 Hz), 7.60-7.43 (m, 5H), 7.30-7.10 (m, 8H), 6.75 (dd, 1H, J=8, 2 Hz). MS (ESI) 499.9 (M+H)+.

30

Example 33

2-chloro-4-[(3-amino-4-chloro)phenyl]-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole, trifluoroacetic acid salt

35 Part A. Preparation of 2-chloro-4-[(3-nitro-4-chloro)phenyl]-5-[(2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole.

Following the procedure described in Example 26, Part B, 2-amino-4-[(3-nitro-4-chloro)phenyl]-5-[(2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole (199 mg, 0.34 mmol) was converted into 150 mg (71%) of the title compound. MS (ESI) 626.9 (M+Na)+.

Part B. Preparation of 2-chloro-4-[(3-amino-4-chloro)phenyl]-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole, trifluoroacetic acid salt.

Following the procedure described in Example 31, Part E, 2-chloro-4-[(3-nitro-4-chloro)phenyl]-5-[(2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole (107 mg, 0.18 mmol) was converted into 10 mg (11%) of the title compound which was a white powder following HPLC purification. ¹H NMR (DMSO-d₆) δ: 10.66 (s, 1H), 7.99 (d, 1H, J=8.0 Hz), 7.60-7.50 (m, 4H), 7.37-7.20 (m, 7H), 6.81 (dd, 1H, J=8.0, 2 Hz). MS (ESI) 518.9 (M+H)+.

Example 34

2-amino-4-[(3-aminomethyl)phenyl]-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole, trifluoroacetic acid salt

Part A. Preparation of 2-amino-4-(3-cyanophenyl)-5-carboethoxythiazole.

To a solution of ethyl 2-bromo-3-(3-cyanophenyl)-3-oxopropionate (2.0 g, 6.75 mmol) in 100 mL of absolute ethanol was added thiourea (0.51 g, 6.75 mmol). The resulting mixture was stirred at 80° C for 3h. The reaction was allowed to cool and the solvent was evaporated in vacuo. The residue was taken up in ethyl acetate, washed with saturated aq NaHCO₃ and brine, dried (MgSO₄) and concentrated in vacuo to yield a solid. Trituration with hexanes/ethyl ether left the title compound as an off-white solid (1.55 g, 65%). ¹H NMR (DMSO-d₆) δ: 8.03 (s, 1H); 7.93 (s, 2H); 7.91 (d, 1H); 7.82 (d, 1H); 7.58 (t, 1H); 4.05 (q, 2H); 1.10 (t, 3H).



Part B. Preparation of 2-amino-4-(3-cyanophenyl)-5-[(2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole.

Following the procedure of Example 7, Part E, 2-amino-4-(3-cyanophenyl)-5-carboethoxythiazole (0.49 g, 1.8 mmol) was converted into 0.31 g (32%) of the title compound as a solid. MS (ESI) 532.3 (M+H)+.

Part C. Preparation of 2-amino-4-[(3-aminomethyl)phenyl]-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole, trifluoroacetic acid salt.

To a solution of lithium aluminum hydride (0.63 mL of a 1.0 M solution in tetrahydrofuran, 0.63 mmol) in 10 mL of tetrahydrofuran at 0°C was added concentrated H₂SO₄ (0.020 mL, 0.32 mmol). This solution was stirred for 30 min and then 2-amino-4-(3-cyanophenyl)-5-[(2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole (112 mg, 0.21 mmol) was added as a solution in tetrahydrofuran. The resulting mixture was allowed to warm to room temperature and then was stirred for 16 h. The reaction was cooled to 0°C and quenched by dropwise addition of water. Dilute aqueous NaOH was added and the mixture was extracted with ethyl acetate. The organic layer was washed with brine, dried (MgSO₄) and concentrated. The residue was taken up in 5 mL of trifluoroacetic acid and stirred at reflux for 30 min. This mixture was cooled and concentrated in vacuo. The residue was purified by prep HPLC to afford the title compound as a white powder. ¹H NMR (DMSO-d₆) δ: 9.77 (s, 1H), 8.13 (broad s, 3H), 7.97 (d, 1H, J=8 Hz), 7.71 (s, 1H), 7.60-7.40 (m, 8H), 7.29-7.20 (m, 6H), 3.98 (broad q, 2H). MS (ESI) 480.0 (M+H)+.

Table 1




Ex.	Ring M	D	R	R ^{1a}	R ^{4a}	A'	MS
1	isoxazole	C(=NH)NH ₂	H	CH ₂ OH	SO ₂ NH ₂	CH	492.1
2	isoxazole	C(=NH)NH ₂	H	H	SO ₂ NH ₂	CH	462.1
3	isoxazole	C(=NH)NH ₂	H	H	SO ₂ Me	CH	461.1
4	isoxazole	C(=NH)NH ₂	H	CH ₂ OMe	SO ₂ NH ₂	N	506.1
5	isoxazole	C(=NH)NH ₂	H	H	CF ₃	CH	451.1
6	isoxazole	C(=NH)NH ₂	H	CF ₃	SO ₂ NH ₂	CH	530.1
7	thiazole	C(=NH)NH ₂	H	NHAc	SO ₂ NH ₂	CH	535.0
8	thiazole	C(=NH)NH ₂	H	NH ₂	SO ₂ NH ₂	CH	493.0
9	thiazole	C(=NH)NH ₂	H	CH ₃	SO ₂ NH ₂	CH	492.3
10	oxazole	C(=NH)NH ₂	H	H	SO ₂ NH ₂	CH	462.1
11	isoxazole	C(=NH)NH ₂	H	H	SO ₂ NHtBu	CH	518.2
12	isoxazole	C(=NH)NH ₂	H	CH ₂ OMe	SO ₂ NH ₂	CH	506.1
13	thiazole	C(=NH)NH ₂	H	Me	CF ₃	CH	481.3
14	thiazole	C(=NH)NH ₂	H	Ph	SO ₂ NH ₂	CH	554.3
15	isoxazole	C(=NH)NH ₂	H	H	SO ₂ Me	CF	479.1
16	isoxazole	C(=NH)NH ₂	H	H	SCF ₃	CH	483.1
17	isoxazole	C(=NH)NH ₂	H	NH ₂	SO ₂ NH ₂	CH	477.1
18	thiazole	C(=NH)NH ₂	H	NHPh	SO ₂ NH ₂	CH	569.0
19	thiazole	C(=NH)NH ₂	H	NHCH ₂ Ph	SO ₂ NH ₂	CH	583.0
20	thiazole	C(=NH)NH ₂	H	NHMe	SO ₂ NH ₂	CH	507.2
21	thiazole	CONH ₂	H	NHMe	SO ₂ NH ₂	CH	508.1
22	thiazole	C(=NH)NH ₂	H	Me	SO ₂ NH ₂	N	493.1
23	thiazole	CONH ₂	H	Me	SO ₂ NH ₂	N	494.1
24	thiazole	C(=NH)NH ₂	H	3-pyridyl	SO ₂ NH ₂	CH	554.3
25	thiazole	CONH ₂	H	3-pyridyl	SO ₂ NH ₂	CH	578.0 (M+Na) ⁺

26	thiazole	C(=NH)NH ₂	H	Cl	SO ₂ NH ₂	CH	512.0
27	thiazole	CONH ₂	H	Cl	SO ₂ NH ₂	CH	534.9 (M+Na) ⁺
28	thiazole	C(=NH)NH ₂	H	Cl	SO ₂ NH ₂	N	513.0
29	thiazole	CONH ₂	H	Cl	SO ₂ NH ₂	N	513.9
30	thiazole	C(=NH)NH ₂	H	OH	SO ₂ NH ₂	N	495.0
31	thiazole	NH ₂	H	Cl	SO ₂ NH ₂	CH	484.9
32	thiazole	NH ₂	Cl	NH ₂	SO ₂ NH ₂	CH	499.9
33	thiazole	NH ₂	Cl	Cl	SO ₂ NH ₂	CH	518.9
34	thiazole	CH ₂ NH ₂	H	NH ₂	SO ₂ NH ₂	CH	480.0


The following tables contain representative examples of the present invention. Each entry in each table is intended to be paired with each formulae at the start of the table.

- 5 For example, example 1 in Table 2 is intended to be paired with each of formulae A-BB and example 1 in Table 3 is intended to be paired with each of fomulae a-dd.


- 10 The following groups are intended for group A in the tables.




2-pyridyl




3-pyridyl




2-pyrimidyl




5-pyrimidyl



2-Cl-phenyl





2-F-phenyl



2,6-diF-phenyl

Table 2





Ex #	R ^{1a}	A	B
1	CH ₃	phenyl	2-(aminosulfonyl)phenyl
2	CH ₃	phenyl	2-(methylaminosulfonyl)phenyl
3	CH ₃	phenyl	1-pyrrolidinocarbonyl
4	CH ₃	phenyl	2-(methylsulfonyl)phenyl
5	CH ₃	phenyl	4-morpholino
6	CH ₃	phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
7	CH ₃	phenyl	4-morpholinocarbonyl
8	CH ₃	phenyl	2-methyl-1-imidazolyl
9	CH ₃	phenyl	5-methyl-1-imidazolyl
10	CH ₃	phenyl	2-methylsulfonyl-1-imidazolyl
11	CH ₃	2-pyridyl	2-(aminosulfonyl)phenyl
12	CH ₃	2-pyridyl	2-(methylaminosulfonyl)phenyl
13	CH ₃	2-pyridyl	1-pyrrolidinocarbonyl
14	CH ₃	2-pyridyl	2-(methylsulfonyl)phenyl
15	CH ₃	2-pyridyl	4-morpholino
16	CH ₃	2-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
17	CH ₃	2-pyridyl	4-morpholinocarbonyl
18	CH ₃	2-pyridyl	2-methyl-1-imidazolyl
19	CH ₃	2-pyridyl	5-methyl-1-imidazolyl
20	CH ₃	2-pyridyl	2-methylsulfonyl-1-imidazolyl
21	CH ₃	3-pyridyl	2-(aminosulfonyl)phenyl
22	CH ₃	3-pyridyl	2-(methylaminosulfonyl)phenyl
23	CH ₃	3-pyridyl	1-pyrrolidinocarbonyl
24	CH ₃	3-pyridyl	2-(methylsulfonyl)phenyl
25	CH ₃	3-pyridyl	4-morpholino
26	CH ₃	3-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
27	CH ₃	3-pyridyl	4-morpholinocarbonyl
28	CH ₃	3-pyridyl	2-methyl-1-imidazolyl
29	CH ₃	3-pyridyl	5-methyl-1-imidazolyl
30	CH ₃	3-pyridyl	2-methylsulfonyl-1-imidazolyl
31	CH ₃	2-pyrimidyl	2-(aminosulfonyl)phenyl
32	CH ₃	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
33	CH ₃	2-pyrimidyl	1-pyrrolidinocarbonyl
34	CH ₃	2-pyrimidyl	2-(methylsulfonyl)phenyl
35	CH ₃	2-pyrimidyl	4-morpholino
36	CH ₃	2-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
37	CH ₃	2-pyrimidyl	4-morpholinocarbonyl
38	CH ₃	2-pyrimidyl	2-methyl-1-imidazolyl
39	CH ₃	2-pyrimidyl	5-methyl-1-imidazolyl
40	CH ₃	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
41	CH ₃	5-pyrimidyl	2-(aminosulfonyl)phenyl
42	CH ₃	5-pyrimidyl	2-(methylaminosulfonyl)phenyl
43	CH ₃	5-pyrimidyl	1-pyrrolidinocarbonyl
44	CH ₃	5-pyrimidyl	2-(methylsulfonyl)phenyl
45	CH ₃	5-pyrimidyl	4-morpholino

46	CH ₃	5-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
47	CH ₃	5-pyrimidyl	4-morpholinocarbonyl
48	CH ₃	5-pyrimidyl	2-methyl-1-imidazolyl
49	CH ₃	5-pyrimidyl	5-methyl-1-imidazolyl
50	CH ₃	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
51	CH ₃	2-Cl-phenyl	2-(aminosulfonyl)phenyl
52	CH ₃	2-Cl-phenyl	2-(methylaminosulfonyl)phenyl
53	CH ₃	2-Cl-phenyl	1-pyrrolidinocarbonyl
54	CH ₃	2-Cl-phenyl	2-(methylsulfonyl)phenyl
55	CH ₃	2-Cl-phenyl	4-morpholino
56	CH ₃	2-Cl-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
57	CH ₃	2-Cl-phenyl	4-morpholinocarbonyl
58	CH ₃	2-Cl-phenyl	2-methyl-1-imidazolyl
59	CH ₃	2-Cl-phenyl	5-methyl-1-imidazolyl
60	CH ₃	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
61	CH ₃	2-F-phenyl	2-(aminosulfonyl)phenyl
62	CH ₃	2-F-phenyl	2-(methylaminosulfonyl)phenyl
63	CH ₃	2-F-phenyl	1-pyrrolidinocarbonyl
64	CH ₃	2-F-phenyl	2-(methylsulfonyl)phenyl
65	CH ₃	2-F-phenyl	4-morpholino
66	CH ₃	2-F-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
67	CH ₃	2-F-phenyl	4-morpholinocarbonyl
68	CH ₃	2-F-phenyl	2-methyl-1-imidazolyl
69	CH ₃	2-F-phenyl	5-methyl-1-imidazolyl
70	CH ₃	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
71	CH ₃	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
72	CH ₃	2,6-diF-phenyl	2-(methylaminosulfonyl)phenyl
73	CH ₃	2,6-diF-phenyl	1-pyrrolidinocarbonyl
74	CH ₃	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
75	CH ₃	2,6-diF-phenyl	4-morpholino
76	CH ₃	2,6-diF-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
77	CH ₃	2,6-diF-phenyl	4-morpholinocarbonyl
78	CH ₃	2,6-diF-phenyl	2-methyl-1-imidazolyl
79	CH ₃	2,6-diF-phenyl	5-methyl-1-imidazolyl
80	CH ₃	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
81	CH ₂ CH ₃	phenyl	2-(aminosulfonyl)phenyl
82	CH ₂ CH ₃	phenyl	2-(methylaminosulfonyl)phenyl
83	CH ₂ CH ₃	phenyl	1-pyrrolidinocarbonyl
84	CH ₂ CH ₃	phenyl	2-(methylsulfonyl)phenyl
85	CH ₂ CH ₃	phenyl	4-morpholino
86	CH ₂ CH ₃	phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
87	CH ₂ CH ₃	phenyl	4-morpholinocarbonyl
88	CH ₂ CH ₃	phenyl	2-methyl-1-imidazolyl
89	CH ₂ CH ₃	phenyl	5-methyl-1-imidazolyl
90	CH ₂ CH ₃	phenyl	2-methylsulfonyl-1-imidazolyl
91	CH ₂ CH ₃	2-pyridyl	2-(aminosulfonyl)phenyl
92	CH ₂ CH ₃	2-pyridyl	2-(methylaminosulfonyl)phenyl
93	CH ₂ CH ₃	2-pyridyl	1-pyrrolidinocarbonyl

94	CH ₂ CH ₃	2-pyridyl	2-(methylsulfonyl)phenyl
95	CH ₂ CH ₃	2-pyridyl	4-morpholino
96	CH ₂ CH ₃	2-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
97	CH ₂ CH ₃	2-pyridyl	4-morpholinocarbonyl
98	CH ₂ CH ₃	2-pyridyl	2-methyl-1-imidazolyl
99	CH ₂ CH ₃	2-pyridyl	5-methyl-1-imidazolyl
100	CH ₂ CH ₃	2-pyridyl	2-methylsulfonyl-1-imidazolyl
101	CH ₂ CH ₃	3-pyridyl	2-(aminosulfonyl)phenyl
102	CH ₂ CH ₃	3-pyridyl	2-(methylaminosulfonyl)phenyl
103	CH ₂ CH ₃	3-pyridyl	1-pyrrolidinocarbonyl
104	CH ₂ CH ₃	3-pyridyl	2-(methylsulfonyl)phenyl
105	CH ₂ CH ₃	3-pyridyl	4-morpholino
106	CH ₂ CH ₃	3-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
107	CH ₂ CH ₃	3-pyridyl	4-morpholinocarbonyl
108	CH ₂ CH ₃	3-pyridyl	2-methyl-1-imidazolyl
109	CH ₂ CH ₃	3-pyridyl	5-methyl-1-imidazolyl
110	CH ₂ CH ₃	3-pyridyl	2-methylsulfonyl-1-imidazolyl
111	CH ₂ CH ₃	2-pyrimidyl	2-(aminosulfonyl)phenyl
112	CH ₂ CH ₃	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
113	CH ₂ CH ₃	2-pyrimidyl	1-pyrrolidinocarbonyl
114	CH ₂ CH ₃	2-pyrimidyl	2-(methylsulfonyl)phenyl
115	CH ₂ CH ₃	2-pyrimidyl	4-morpholino
116	CH ₂ CH ₃	2-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
117	CH ₂ CH ₃	2-pyrimidyl	4-morpholinocarbonyl
118	CH ₂ CH ₃	2-pyrimidyl	2-methyl-1-imidazolyl
119	CH ₂ CH ₃	2-pyrimidyl	5-methyl-1-imidazolyl
120	CH ₂ CH ₃	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
121	CH ₂ CH ₃	5-pyrimidyl	2-(aminosulfonyl)phenyl
122	CH ₂ CH ₃	5-pyrimidyl	2-(methylaminosulfonyl)phenyl
123	CH ₂ CH ₃	5-pyrimidyl	1-pyrrolidinocarbonyl
124	CH ₂ CH ₃	5-pyrimidyl	2-(methylsulfonyl)phenyl
125	CH ₂ CH ₃	5-pyrimidyl	4-morpholino
126	CH ₂ CH ₃	5-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
127	CH ₂ CH ₃	5-pyrimidyl	4-morpholinocarbonyl
128	CH ₂ CH ₃	5-pyrimidyl	2-methyl-1-imidazolyl
129	CH ₂ CH ₃	5-pyrimidyl	5-methyl-1-imidazolyl
130	CH ₂ CH ₃	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
131	CH ₂ CH ₃	2-Cl-phenyl	2-(aminosulfonyl)phenyl
132	CH ₂ CH ₃	2-Cl-phenyl	2-(methylaminosulfonyl)phenyl
133	CH ₂ CH ₃	2-Cl-phenyl	1-pyrrolidinocarbonyl
134	CH ₂ CH ₃	2-Cl-phenyl	2-(methylsulfonyl)phenyl
135	CH ₂ CH ₃	2-Cl-phenyl	4-morpholino
136	CH ₂ CH ₃	2-Cl-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
137	CH ₂ CH ₃	2-Cl-phenyl	4-morpholinocarbonyl
138	CH ₂ CH ₃	2-Cl-phenyl	2-methyl-1-imidazolyl
139	CH ₂ CH ₃	2-Cl-phenyl	5-methyl-1-imidazolyl
140	CH ₂ CH ₃	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
141	CH ₂ CH ₃	2-F-phenyl	2-(aminosulfonyl)phenyl

142	CH ₂ CH ₃	2-F-phenyl	2-(methylaminosulfonyl)phenyl
143	CH ₂ CH ₃	2-F-phenyl	1-pyrrolidinocarbonyl
144	CH ₂ CH ₃	2-F-phenyl	2-(methylsulfonyl)phenyl
145	CH ₂ CH ₃	2-F-phenyl	4-morpholino
146	CH ₂ CH ₃	2-F-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
147	CH ₂ CH ₃	2-F-phenyl	4-morpholinocarbonyl
148	CH ₂ CH ₃	2-F-phenyl	2-methyl-1-imidazolyl
149	CH ₂ CH ₃	2-F-phenyl	5-methyl-1-imidazolyl
150	CH ₂ CH ₃	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
151	CH ₂ CH ₃	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
152	CH ₂ CH ₃	2,6-diF-phenyl	2-(methylaminosulfonyl)phenyl
153	CH ₂ CH ₃	2,6-diF-phenyl	1-pyrrolidinocarbonyl
154	CH ₂ CH ₃	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
155	CH ₂ CH ₃	2,6-diF-phenyl	4-morpholino
156	CH ₂ CH ₃	2,6-diF-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
157	CH ₂ CH ₃	2,6-diF-phenyl	4-morpholinocarbonyl
158	CH ₂ CH ₃	2,6-diF-phenyl	2-methyl-1-imidazolyl
159	CH ₂ CH ₃	2,6-diF-phenyl	5-methyl-1-imidazolyl
160	CH ₂ CH ₃	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
161	CF ₃	phenyl	2-(aminosulfonyl)phenyl
162	CF ₃	phenyl	2-(methylaminosulfonyl)phenyl
163	CF ₃	phenyl	1-pyrrolidinocarbonyl
164	CF ₃	phenyl	2-(methylsulfonyl)phenyl
165	CF ₃	phenyl	4-morpholino
166	CF ₃	phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
167	CF ₃	phenyl	4-morpholinocarbonyl
168	CF ₃	phenyl	2-methyl-1-imidazolyl
169	CF ₃	phenyl	5-methyl-1-imidazolyl
170	CF ₃	phenyl	2-methylsulfonyl-1-imidazolyl
171	CF ₃	2-pyridyl	2-(aminosulfonyl)phenyl
172	CF ₃	2-pyridyl	2-(methylaminosulfonyl)phenyl
173	CF ₃	2-pyridyl	1-pyrrolidinocarbonyl
174	CF ₃	2-pyridyl	2-(methylsulfonyl)phenyl
175	CF ₃	2-pyridyl	4-morpholino
176	CF ₃	2-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
177	CF ₃	2-pyridyl	4-morpholinocarbonyl
178	CF ₃	2-pyridyl	2-methyl-1-imidazolyl
179	CF ₃	2-pyridyl	5-methyl-1-imidazolyl
180	CF ₃	2-pyridyl	2-methylsulfonyl-1-imidazolyl
181	CF ₃	3-pyridyl	2-(aminosulfonyl)phenyl
182	CF ₃	3-pyridyl	2-(methylaminosulfonyl)phenyl
183	CF ₃	3-pyridyl	1-pyrrolidinocarbonyl
184	CF ₃	3-pyridyl	2-(methylsulfonyl)phenyl
185	CF ₃	3-pyridyl	4-morpholino
186	CF ₃	3-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
187	CF ₃	3-pyridyl	4-morpholinocarbonyl
188	CF ₃	3-pyridyl	2-methyl-1-imidazolyl
189	CF ₃	3-pyridyl	5-methyl-1-imidazolyl

190	CF ₃	3-pyridyl	2-methylsulfonyl-1-imidazolyl
191	CF ₃	2-pyrimidyl	2-(aminosulfonyl)phenyl
192	CF ₃	2-pyrimidyl	2-(methylaninosulfonyl)phenyl
193	CF ₃	2-pyrimidyl	1-pyrrolidinocarbonyl
194	CF ₃	2-pyrimidyl	2-(methylsulfonyl)phenyl
195	CF ₃	2-pyrimidyl	4-morpholino
196	CF ₃	2-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
197	CF ₃	2-pyrimidyl	4-morpholinocarbonyl
198	CF ₃	2-pyrimidyl	2-methyl-1-imidazolyl
199	CF ₃	2-pyrimidyl	5-methyl-1-imidazolyl
200	CF ₃	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
201	CF ₃	5-pyrimidyl	2-(aminosulfonyl)phenyl
202	CF ₃	5-pyrimidyl	2-(methylaninosulfonyl)phenyl
203	CF ₃	5-pyrimidyl	1-pyrrolidinocarbonyl
204	CF ₃	5-pyrimidyl	2-(methylsulfonyl)phenyl
205	CF ₃	5-pyrimidyl	4-morpholino
206	CF ₃	5-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
207	CF ₃	5-pyrimidyl	4-morpholinocarbonyl
208	CF ₃	5-pyrimidyl	2-methyl-1-imidazolyl
209	CF ₃	5-pyrimidyl	5-methyl-1-imidazolyl
210	CF ₃	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
211	CF ₃	2-Cl-phenyl	2-(aminosulfonyl)phenyl
212	CF ₃	2-Cl-phenyl	2-(methylaninosulfonyl)phenyl
213	CF ₃	2-Cl-phenyl	1-pyrrolidinocarbonyl
214	CF ₃	2-Cl-phenyl	2-(methylsulfonyl)phenyl
215	CF ₃	2-Cl-phenyl	4-morpholino
216	CF ₃	2-Cl-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
217	CF ₃	2-Cl-phenyl	4-morpholinocarbonyl
218	CF ₃	2-Cl-phenyl	2-methyl-1-imidazolyl
219	CF ₃	2-Cl-phenyl	5-methyl-1-imidazolyl
220	CF ₃	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
221	CF ₃	2-F-phenyl	2-(aminosulfonyl)phenyl
222	CF ₃	2-F-phenyl	2-(methylaninosulfonyl)phenyl
223	CF ₃	2-F-phenyl	1-pyrrolidinocarbonyl
224	CF ₃	2-F-phenyl	2-(methylsulfonyl)phenyl
225	CF ₃	2-F-phenyl	4-morpholino
226	CF ₃	2-F-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
227	CF ₃	2-F-phenyl	4-morpholinocarbonyl
228	CF ₃	2-F-phenyl	2-methyl-1-imidazolyl
229	CF ₃	2-F-phenyl	5-methyl-1-imidazolyl
230	CF ₃	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
231	CF ₃	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
232	CF ₃	2,6-diF-phenyl	2-(methylaninosulfonyl)phenyl
233	CF ₃	2,6-diF-phenyl	1-pyrrolidinocarbonyl
234	CF ₃	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
235	CF ₃	2,6-diF-phenyl	4-morpholino
236	CF ₃	2,6-diF-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
237	CF ₃	2,6-diF-phenyl	4-morpholinocarbonyl

238	CF ₃	2,6-diF-phenyl	2-methyl-1-imidazolyl
239	CF ₃	2,6-diF-phenyl	5-methyl-1-imidazolyl
240	CF ₃	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
241	SCH ₃	phenyl	2-(aminosulfonyl)phenyl
242	SCH ₃	phenyl	2-(methylaminosulfonyl)phenyl
243	SCH ₃	phenyl	1-pyrrolidinocarbonyl
244	SCH ₃	phenyl	2-(methylsulfonyl)phenyl
245	SCH ₃	phenyl	4-morpholino
246	SCH ₃	phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
247	SCH ₃	phenyl	4-morpholinocarbonyl
248	SCH ₃	phenyl	2-methyl-1-imidazolyl
249	SCH ₃	phenyl	5-methyl-1-imidazolyl
250	SCH ₃	phenyl	2-methylsulfonyl-1-imidazolyl
251	SCH ₃	2-pyridyl	2-(aminosulfonyl)phenyl
252	SCH ₃	2-pyridyl	2-(methylaminosulfonyl)phenyl
253	SCH ₃	2-pyridyl	1-pyrrolidinocarbonyl
254	SCH ₃	2-pyridyl	2-(methylsulfonyl)phenyl
255	SCH ₃	2-pyridyl	4-morpholino
256	SCH ₃	2-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
257	SCH ₃	2-pyridyl	4-morpholinocarbonyl
258	SCH ₃	2-pyridyl	2-methyl-1-imidazolyl
259	SCH ₃	2-pyridyl	5-methyl-1-imidazolyl
260	SCH ₃	2-pyridyl	2-methylsulfonyl-1-imidazolyl
261	SCH ₃	3-pyridyl	2-(aminosulfonyl)phenyl
262	SCH ₃	3-pyridyl	2-(methylaminosulfonyl)phenyl
263	SCH ₃	3-pyridyl	1-pyrrolidinocarbonyl
264	SCH ₃	3-pyridyl	2-(methylsulfonyl)phenyl
265	SCH ₃	3-pyridyl	4-morpholino
266	SCH ₃	3-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
267	SCH ₃	3-pyridyl	4-morpholinocarbonyl
268	SCH ₃	3-pyridyl	2-methyl-1-imidazolyl
269	SCH ₃	3-pyridyl	5-methyl-1-imidazolyl
270	SCH ₃	3-pyridyl	2-methylsulfonyl-1-imidazolyl
271	SCH ₃	2-pyrimidyl	2-(aminosulfonyl)phenyl
272	SCH ₃	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
273	SCH ₃	2-pyrimidyl	1-pyrrolidinocarbonyl
274	SCH ₃	2-pyrimidyl	2-(methylsulfonyl)phenyl
275	SCH ₃	2-pyrimidyl	4-morpholino
276	SCH ₃	2-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
277	SCH ₃	2-pyrimidyl	4-morpholinocarbonyl
278	SCH ₃	2-pyrimidyl	2-methyl-1-imidazolyl
279	SCH ₃	2-pyrimidyl	5-methyl-1-imidazolyl
280	SCH ₃	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
281	SCH ₃	5-pyrimidyl	2-(aminosulfonyl)phenyl
282	SCH ₃	5-pyrimidyl	2-(methylaminosulfonyl)phenyl
283	SCH ₃	5-pyrimidyl	1-pyrrolidinocarbonyl
284	SCH ₃	5-pyrimidyl	2-(methylsulfonyl)phenyl
285	SCH ₃	5-pyrimidyl	4-morpholino

286	SCH ₃	5-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
287	SCH ₃	5-pyrimidyl	4-morpholinocarbonyl
288	SCH ₃	5-pyrimidyl	2-methyl-1-imidazolyl
289	SCH ₃	5-pyrimidyl	5-methyl-1-imidazolyl
290	SCH ₃	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
291	SCH ₃	2-Cl-phenyl	2-(aminosulfonyl)phenyl
292	SCH ₃	2-Cl-phenyl	2-(methylaminosulfonyl)phenyl
293	SCH ₃	2-Cl-phenyl	1-pyrrolidinocarbonyl
294	SCH ₃	2-Cl-phenyl	2-(methylsulfonyl)phenyl
295	SCH ₃	2-Cl-phenyl	4-morpholino
296	SCH ₃	2-Cl-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
297	SCH ₃	2-Cl-phenyl	4-morpholinocarbonyl
298	SCH ₃	2-Cl-phenyl	2-methyl-1-imidazolyl
299	SCH ₃	2-Cl-phenyl	5-methyl-1-imidazolyl
300	SCH ₃	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
301	SCH ₃	2-F-phenyl	2-(aminosulfonyl)phenyl
302	SCH ₃	2-F-phenyl	2-(methylaminosulfonyl)phenyl
303	SCH ₃	2-F-phenyl	1-pyrrolidinocarbonyl
304	SCH ₃	2-F-phenyl	2-(methylsulfonyl)phenyl
305	SCH ₃	2-F-phenyl	4-morpholino
306	SCH ₃	2-F-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
307	SCH ₃	2-F-phenyl	4-morpholinocarbonyl
308	SCH ₃	2-F-phenyl	2-methyl-1-imidazolyl
309	SCH ₃	2-F-phenyl	5-methyl-1-imidazolyl
310	SCH ₃	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
311	SCH ₃	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
312	SCH ₃	2,6-diF-phenyl	2-(methylaminosulfonyl)phenyl
313	SCH ₃	2,6-diF-phenyl	1-pyrrolidinocarbonyl
314	SCH ₃	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
315	SCH ₃	2,6-diF-phenyl	4-morpholino
316	SCH ₃	2,6-diF-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
317	SCH ₃	2,6-diF-phenyl	4-morpholinocarbonyl
318	SCH ₃	2,6-diF-phenyl	2-methyl-1-imidazolyl
319	SCH ₃	2,6-diF-phenyl	5-methyl-1-imidazolyl
320	SCH ₃	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
321	SOCH ₃	phenyl	2-(aminosulfonyl)phenyl
322	SOCH ₃	phenyl	2-(methylaminosulfonyl)phenyl
323	SOCH ₃	phenyl	1-pyrrolidinocarbonyl
324	SOCH ₃	phenyl	2-(methylsulfonyl)phenyl
325	SOCH ₃	phenyl	4-morpholino
326	SOCH ₃	phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
327	SOCH ₃	phenyl	4-morpholinocarbonyl
328	SOCH ₃	phenyl	2-methyl-1-imidazolyl
329	SOCH ₃	phenyl	5-methyl-1-imidazolyl
330	SOCH ₃	phenyl	2-methylsulfonyl-1-imidazolyl
331	SOCH ₃	2-pyridyl	2-(aminosulfonyl)phenyl
332	SOCH ₃	2-pyridyl	2-(methylaminosulfonyl)phenyl
333	SOCH ₃	2-pyridyl	1-pyrrolidinocarbonyl

334	SOCH ₃	2-pyridyl	2-(methylsulfonyl)phenyl
335	SOCH ₃	2-pyridyl	4-morpholino
336	SOCH ₃	2-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
337	SOCH ₃	2-pyridyl	4-morpholinocarbonyl
338	SOCH ₃	2-pyridyl	2-methyl-1-imidazolyl
339	SOCH ₃	2-pyridyl	5-methyl-1-imidazolyl
340	SOCH ₃	2-pyridyl	2-methylsulfonyl-1-imidazolyl
341	SOCH ₃	3-pyridyl	2-(aminosulfonyl)phenyl
342	SOCH ₃	3-pyridyl	2-(methylaminosulfonyl)phenyl
343	SOCH ₃	3-pyridyl	1-pyrrolidinocarbonyl
344	SOCH ₃	3-pyridyl	2-(methylsulfonyl)phenyl
345	SOCH ₃	3-pyridyl	4-morpholino
346	SOCH ₃	3-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
347	SOCH ₃	3-pyridyl	4-morpholinocarbonyl
348	SOCH ₃	3-pyridyl	2-methyl-1-imidazolyl
349	SOCH ₃	3-pyridyl	5-methyl-1-imidazolyl
350	SOCH ₃	3-pyridyl	2-methylsulfonyl-1-imidazolyl
351	SOCH ₃	2-pyrimidyl	2-(aminosulfonyl)phenyl
352	SOCH ₃	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
353	SOCH ₃	2-pyrimidyl	1-pyrrolidinocarbonyl
354	SOCH ₃	2-pyrimidyl	2-(methylsulfonyl)phenyl
355	SOCH ₃	2-pyrimidyl	4-morpholino
356	SOCH ₃	2-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
357	SOCH ₃	2-pyrimidyl	4-morpholinocarbonyl
358	SOCH ₃	2-pyrimidyl	2-methyl-1-imidazolyl
359	SOCH ₃	2-pyrimidyl	5-methyl-1-imidazolyl
360	SOCH ₃	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
361	SOCH ₃	5-pyrimidyl	2-(aminosulfonyl)phenyl
362	SOCH ₃	5-pyrimidyl	2-(methylaminosulfonyl)phenyl
363	SOCH ₃	5-pyrimidyl	1-pyrrolidinocarbonyl
364	SOCH ₃	5-pyrimidyl	2-(methylsulfonyl)phenyl
365	SOCH ₃	5-pyrimidyl	4-morpholino
366	SOCH ₃	5-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
367	SOCH ₃	5-pyrimidyl	4-morpholinocarbonyl
368	SOCH ₃	5-pyrimidyl	2-methyl-1-imidazolyl
369	SOCH ₃	5-pyrimidyl	5-methyl-1-imidazolyl
370	SOCH ₃	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
371	SOCH ₃	2-Cl-phenyl	2-(aminosulfonyl)phenyl
372	SOCH ₃	2-Cl-phenyl	2-(methylaminosulfonyl)phenyl
373	SOCH ₃	2-Cl-phenyl	1-pyrrolidinocarbonyl
374	SOCH ₃	2-Cl-phenyl	2-(methylsulfonyl)phenyl
375	SOCH ₃	2-Cl-phenyl	4-morpholino
376	SOCH ₃	2-Cl-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
377	SOCH ₃	2-Cl-phenyl	4-morpholinocarbonyl
378	SOCH ₃	2-Cl-phenyl	2-methyl-1-imidazolyl
379	SOCH ₃	2-Cl-phenyl	5-methyl-1-imidazolyl
380	SOCH ₃	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
381	SOCH ₃	2-F-phenyl	2-(aminosulfonyl)phenyl

382	SOCH ₃	2-F-phenyl	2-(methylaminosulfonyl)phenyl
383	SOCH ₃	2-F-phenyl	1-pyrrolidinocarbonyl
384	SOCH ₃	2-F-phenyl	2-(methylsulfonyl)phenyl
385	SOCH ₃	2-F-phenyl	4-morpholino
386	SOCH ₃	2-F-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
387	SOCH ₃	2-F-phenyl	4-morpholinocarbonyl
388	SOCH ₃	2-F-phenyl	2-methyl-1-imidazolyl
389	SOCH ₃	2-F-phenyl	5-methyl-1-imidazolyl
390	SOCH ₃	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
391	SOCH ₃	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
392	SOCH ₃	2,6-diF-phenyl	2-(methylaminosulfonyl)phenyl
393	SOCH ₃	2,6-diF-phenyl	1-pyrrolidinocarbonyl
394	SOCH ₃	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
395	SOCH ₃	2,6-diF-phenyl	4-morpholino
396	SOCH ₃	2,6-diF-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
397	SOCH ₃	2,6-diF-phenyl	4-morpholinocarbonyl
398	SOCH ₃	2,6-diF-phenyl	2-methyl-1-imidazolyl
399	SOCH ₃	2,6-diF-phenyl	5-methyl-1-imidazolyl
400	SOCH ₃	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
401	SO ₂ CH ₃	phenyl	2-(aminosulfonyl)phenyl
402	SO ₂ CH ₃	phenyl	2-(methylaminosulfonyl)phenyl
403	SO ₂ CH ₃	phenyl	1-pyrrolidinocarbonyl
404	SO ₂ CH ₃	phenyl	2-(methylsulfonyl)phenyl
405	SO ₂ CH ₃	phenyl	4-morpholino
406	SO ₂ CH ₃	phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
407	SO ₂ CH ₃	phenyl	4-morpholinocarbonyl
408	SO ₂ CH ₃	phenyl	2-methyl-1-imidazolyl
409	SO ₂ CH ₃	phenyl	5-methyl-1-imidazolyl
410	SO ₂ CH ₃	phenyl	2-methylsulfonyl-1-imidazolyl
411	SO ₂ CH ₃	2-pyridyl	2-(aminosulfonyl)phenyl
412	SO ₂ CH ₃	2-pyridyl	2-(methylaminosulfonyl)phenyl
413	SO ₂ CH ₃	2-pyridyl	1-pyrrolidinocarbonyl
414	SO ₂ CH ₃	2-pyridyl	2-(methylsulfonyl)phenyl
415	SO ₂ CH ₃	2-pyridyl	4-morpholino
416	SO ₂ CH ₃	2-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
417	SO ₂ CH ₃	2-pyridyl	4-morpholinocarbonyl
418	SO ₂ CH ₃	2-pyridyl	2-methyl-1-imidazolyl
419	SO ₂ CH ₃	2-pyridyl	5-methyl-1-imidazolyl
420	SO ₂ CH ₃	2-pyridyl	2-methylsulfonyl-1-imidazolyl
421	SO ₂ CH ₃	3-pyridyl	2-(aminosulfonyl)phenyl
422	SO ₂ CH ₃	3-pyridyl	2-(methylaminosulfonyl)phenyl
423	SO ₂ CH ₃	3-pyridyl	1-pyrrolidinocarbonyl
424	SO ₂ CH ₃	3-pyridyl	2-(methylsulfonyl)phenyl
425	SO ₂ CH ₃	3-pyridyl	4-morpholino
426	SO ₂ CH ₃	3-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
427	SO ₂ CH ₃	3-pyridyl	4-morpholinocarbonyl
428	SO ₂ CH ₃	3-pyridyl	2-methyl-1-imidazolyl
429	SO ₂ CH ₃	3-pyridyl	5-methyl-1-imidazolyl

430	SO ₂ CH ₃	3-pyridyl	2-methylsulfonyl-1-imidazolyl
431	SO ₂ CH ₃	2-pyrimidyl	2-(aminosulfonyl)phenyl
432	SO ₂ CH ₃	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
433	SO ₂ CH ₃	2-pyrimidyl	1-pyrrolidinocarbonyl
434	SO ₂ CH ₃	2-pyrimidyl	2-(methylsulfonyl)phenyl
435	SO ₂ CH ₃	2-pyrimidyl	4-morpholino
436	SO ₂ CH ₃	2-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
437	SO ₂ CH ₃	2-pyrimidyl	4-morpholinocarbonyl
438	SO ₂ CH ₃	2-pyrimidyl	2-methyl-1-imidazolyl
439	SO ₂ CH ₃	2-pyrimidyl	5-methyl-1-imidazolyl
440	SO ₂ CH ₃	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
441	SO ₂ CH ₃	5-pyrimidyl	2-(aminosulfonyl)phenyl
442	SO ₂ CH ₃	5-pyrimidyl	2-(methylaminosulfonyl)phenyl
443	SO ₂ CH ₃	5-pyrimidyl	1-pyrrolidinocarbonyl
444	SO ₂ CH ₃	5-pyrimidyl	2-(methylsulfonyl)phenyl
445	SO ₂ CH ₃	5-pyrimidyl	4-morpholino
446	SO ₂ CH ₃	5-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
447	SO ₂ CH ₃	5-pyrimidyl	4-morpholinocarbonyl
448	SO ₂ CH ₃	5-pyrimidyl	2-methyl-1-imidazolyl
449	SO ₂ CH ₃	5-pyrimidyl	5-methyl-1-imidazolyl
450	SO ₂ CH ₃	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
451	SO ₂ CH ₃	2-Cl-phenyl	2-(aminosulfonyl)phenyl
452	SO ₂ CH ₃	2-Cl-phenyl	2-(methylaminosulfonyl)phenyl
453	SO ₂ CH ₃	2-Cl-phenyl	1-pyrrolidinocarbonyl
454	SO ₂ CH ₃	2-Cl-phenyl	2-(methylsulfonyl)phenyl
455	SO ₂ CH ₃	2-Cl-phenyl	4-morpholino
456	SO ₂ CH ₃	2-Cl-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
457	SO ₂ CH ₃	2-Cl-phenyl	4-morpholinocarbonyl
458	SO ₂ CH ₃	2-Cl-phenyl	2-methyl-1-imidazolyl
459	SO ₂ CH ₃	2-Cl-phenyl	5-methyl-1-imidazolyl
460	SO ₂ CH ₃	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
461	SO ₂ CH ₃	2-F-phenyl	2-(aminosulfonyl)phenyl
462	SO ₂ CH ₃	2-F-phenyl	2-(methylaminosulfonyl)phenyl
463	SO ₂ CH ₃	2-F-phenyl	1-pyrrolidinocarbonyl
464	SO ₂ CH ₃	2-F-phenyl	2-(methylsulfonyl)phenyl
465	SO ₂ CH ₃	2-F-phenyl	4-morpholino
466	SO ₂ CH ₃	2-F-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
467	SO ₂ CH ₃	2-F-phenyl	4-morpholinocarbonyl
468	SO ₂ CH ₃	2-F-phenyl	2-methyl-1-imidazolyl
469	SO ₂ CH ₃	2-F-phenyl	5-methyl-1-imidazolyl
470	SO ₂ CH ₃	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
471	SO ₂ CH ₃	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
472	SO ₂ CH ₃	2,6-diF-phenyl	2-(methylaminosulfonyl)phenyl
473	SO ₂ CH ₃	2,6-diF-phenyl	1-pyrrolidinocarbonyl
474	SO ₂ CH ₃	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
475	SO ₂ CH ₃	2,6-diF-phenyl	4-morpholino
476	SO ₂ CH ₃	2,6-diF-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
477	SO ₂ CH ₃	2,6-diF-phenyl	4-morpholinocarbonyl

478	SO ₂ CH ₃	2,6-diF-phenyl	2-methyl-1-imidazolyl
479	SO ₂ CH ₃	2,6-diF-phenyl	5-methyl-1-imidazolyl
480	SO ₂ CH ₃	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
481	CH ₂ NH- SO ₂ CH ₃	phenyl	2-(aminosulfonyl)phenyl
482	CH ₂ NH- SO ₂ CH ₃	phenyl	2-(methylaminosulfonyl)phenyl
483	CH ₂ NH- SO ₂ CH ₃	phenyl	1-pyrrolidinocarbonyl
484	CH ₂ NH- SO ₂ CH ₃	phenyl	2-(methylsulfonyl)phenyl
485	CH ₂ NH- SO ₂ CH ₃	phenyl	4-morpholino
486	CH ₂ NH- SO ₂ CH ₃	phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
487	CH ₂ NH- SO ₂ CH ₃	phenyl	4-morpholinocarbonyl
488	CH ₂ NH- SO ₂ CH ₃	phenyl	2-methyl-1-imidazolyl
489	CH ₂ NH- SO ₂ CH ₃	phenyl	5-methyl-1-imidazolyl
490	CH ₂ NH- SO ₂ CH ₃	phenyl	2-methylsulfonyl-1-imidazolyl
491	CH ₂ NH- SO ₂ CH ₃	2-pyridyl	2-(aminosulfonyl)phenyl
492	CH ₂ NH- SO ₂ CH ₃	2-pyridyl	2-(methylaminosulfonyl)phenyl
493	CH ₂ NH- SO ₂ CH ₃	2-pyridyl	1-pyrrolidinocarbonyl
494	CH ₂ NH- SO ₂ CH ₃	2-pyridyl	2-(methylsulfonyl)phenyl
495	CH ₂ NH- SO ₂ CH ₃	2-pyridyl	4-morpholino
496	CH ₂ NH- SO ₂ CH ₃	2-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
497	CH ₂ NH- SO ₂ CH ₃	2-pyridyl	4-morpholinocarbonyl
498	CH ₂ NH- SO ₂ CH ₃	2-pyridyl	2-methyl-1-imidazolyl
499	CH ₂ NH- SO ₂ CH ₃	2-pyridyl	5-methyl-1-imidazolyl
500	CH ₂ NH- SO ₂ CH ₃	2-pyridyl	2-methylsulfonyl-1-imidazolyl
501	CH ₂ NH- SO ₂ CH ₃	3-pyridyl	2-(aminosulfonyl)phenyl
502	CH ₂ NH- SO ₂ CH ₃	3-pyridyl	2-(methylaminosulfonyl)phenyl

503	CH ₂ NH-SO ₂ CH ₃	3-pyridyl	1-pyrrolidinocarbonyl
504	CH ₂ NH-SO ₂ CH ₃	3-pyridyl	2-(methylsulfonyl)phenyl
505	CH ₂ NH-SO ₂ CH ₃	3-pyridyl	4-morpholino
506	CH ₂ NH-SO ₂ CH ₃	3-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
507	CH ₂ NH-SO ₂ CH ₃	3-pyridyl	4-morpholinocarbonyl
508	CH ₂ NH-SO ₂ CH ₃	3-pyridyl	2-methyl-1-imidazolyl
509	CH ₂ NH-SO ₂ CH ₃	3-pyridyl	5-methyl-1-imidazolyl
510	CH ₂ NH-SO ₂ CH ₃	3-pyridyl	2-methylsulfonyl-1-imidazolyl
511	CH ₂ NH-SO ₂ CH ₃	2-pyrimidyl	2-(aminosulfonyl)phenyl
512	CH ₂ NH-SO ₂ CH ₃	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
513	CH ₂ NH-SO ₂ CH ₃	2-pyrimidyl	1-pyrrolidinocarbonyl
514	CH ₂ NH-SO ₂ CH ₃	2-pyrimidyl	2-(methylsulfonyl)phenyl
515	CH ₂ NH-SO ₂ CH ₃	2-pyrimidyl	4-morpholino
516	CH ₂ NH-SO ₂ CH ₃	2-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
517	CH ₂ NH-SO ₂ CH ₃	2-pyrimidyl	4-morpholinocarbonyl
518	CH ₂ NH-SO ₂ CH ₃	2-pyrimidyl	2-methyl-1-imidazolyl
519	CH ₂ NH-SO ₂ CH ₃	2-pyrimidyl	5-methyl-1-imidazolyl
520	CH ₂ NH-SO ₂ CH ₃	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
521	CH ₂ NH-SO ₂ CH ₃	5-pyrimidyl	2-(aminosulfonyl)phenyl
522	CH ₂ NH-SO ₂ CH ₃	5-pyrimidyl	2-(methylaminosulfonyl)phenyl
523	CH ₂ NH-SO ₂ CH ₃	5-pyrimidyl	1-pyrrolidinocarbonyl
524	CH ₂ NH-SO ₂ CH ₃	5-pyrimidyl	2-(methylsulfonyl)phenyl
525	CH ₂ NH-SO ₂ CH ₃	5-pyrimidyl	4-morpholino
526	CH ₂ NH-SO ₂ CH ₃	5-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl

527	CH ₂ NH-SO ₂ CH ₃	5-pyrimidyl	4-morpholinocarbonyl
528	CH ₂ NH-SO ₂ CH ₃	5-pyrimidyl	2-methyl-1-imidazolyl
529	CH ₂ NH-SO ₂ CH ₃	5-pyrimidyl	5-methyl-1-imidazolyl
530	CH ₂ NH-SO ₂ CH ₃	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
531	CH ₂ NH-SO ₂ CH ₃	2-Cl-phenyl	2-(aminosulfonyl)phenyl
532	CH ₂ NH-SO ₂ CH ₃	2-Cl-phenyl	2-(methylaminosulfonyl)phenyl
533	CH ₂ NH-SO ₂ CH ₃	2-Cl-phenyl	1-pyrrolidinocarbonyl
534	CH ₂ NH-SO ₂ CH ₃	2-Cl-phenyl	2-(methylsulfonyl)phenyl
535	CH ₂ NH-SO ₂ CH ₃	2-Cl-phenyl	4-morpholino
536	CH ₂ NH-SO ₂ CH ₃	2-Cl-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
537	CH ₂ NH-SO ₂ CH ₃	2-Cl-phenyl	4-morpholinocarbonyl
538	CH ₂ NH-SO ₂ CH ₃	2-Cl-phenyl	2-methyl-1-imidazolyl
539	CH ₂ NH-SO ₂ CH ₃	2-Cl-phenyl	5-methyl-1-imidazolyl
540	CH ₂ NH-SO ₂ CH ₃	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
541	CH ₂ NH-SO ₂ CH ₃	2-F-phenyl	2-(aminosulfonyl)phenyl
542	CH ₂ NH-SO ₂ CH ₃	2-F-phenyl	2-(methylaminosulfonyl)phenyl
543	CH ₂ NH-SO ₂ CH ₃	2-F-phenyl	1-pyrrolidinocarbonyl
544	CH ₂ NH-SO ₂ CH ₃	2-F-phenyl	2-(methylsulfonyl)phenyl
545	CH ₂ NH-SO ₂ CH ₃	2-F-phenyl	4-morpholino
546	CH ₂ NH-SO ₂ CH ₃	2-F-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
547	CH ₂ NH-SO ₂ CH ₃	2-F-phenyl	4-morpholinocarbonyl
548	CH ₂ NH-SO ₂ CH ₃	2-F-phenyl	2-methyl-1-imidazolyl
549	CH ₂ NH-SO ₂ CH ₃	2-F-phenyl	5-methyl-1-imidazolyl
550	CH ₂ NH-SO ₂ CH ₃	2-F-phenyl	2-methylsulfonyl-1-imidazolyl

551	CH ₂ NH- SO ₂ CH ₃	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
552	CH ₂ NH- SO ₂ CH ₃	2,6-diF-phenyl	2-(methylaminosulfonyl)phenyl
553	CH ₂ NH- SO ₂ CH ₃	2,6-diF-phenyl	1-pyrrolidinocarbonyl
554	CH ₂ NH- SO ₂ CH ₃	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
555	CH ₂ NH- SO ₂ CH ₃	2,6-diF-phenyl	4-morpholino
556	CH ₂ NH- SO ₂ CH ₃	2,6-diF-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
557	CH ₂ NH- SO ₂ CH ₃	2,6-diF-phenyl	4-morpholinocarbonyl
558	CH ₂ NH- SO ₂ CH ₃	2,6-diF-phenyl	2-methyl-1-imidazolyl
559	CH ₂ NH- SO ₂ CH ₃	2,6-diF-phenyl	5-methyl-1-imidazolyl
560	CH ₂ NH- SO ₂ CH ₃	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
561	Cl	phenyl	2-(aminosulfonyl)phenyl
562	Cl	phenyl	2-(methylaminosulfonyl)phenyl
563	Cl	phenyl	1-pyrrolidinocarbonyl
564	Cl	phenyl	2-(methylsulfonyl)phenyl
565	Cl	phenyl	4-morpholino
566	Cl	phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
567	Cl	phenyl	4-morpholinocarbonyl
568	Cl	phenyl	2-methyl-1-imidazolyl
569	Cl	phenyl	5-methyl-1-imidazolyl
570	Cl	phenyl	2-methylsulfonyl-1-imidazolyl
571	Cl	2-pyridyl	2-(aminosulfonyl)phenyl
572	Cl	2-pyridyl	2-(methylaminosulfonyl)phenyl
573	Cl	2-pyridyl	1-pyrrolidinocarbonyl
574	Cl	2-pyridyl	2-(methylsulfonyl)phenyl
575	Cl	2-pyridyl	4-morpholino
576	Cl	2-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
577	Cl	2-pyridyl	4-morpholinocarbonyl
578	Cl	2-pyridyl	2-methyl-1-imidazolyl
579	Cl	2-pyridyl	5-methyl-1-imidazolyl
580	Cl	2-pyridyl	2-methylsulfonyl-1-imidazolyl
581	Cl	3-pyridyl	2-(aminosulfonyl)phenyl
582	Cl	3-pyridyl	2-(methylaminosulfonyl)phenyl
583	Cl	3-pyridyl	1-pyrrolidinocarbonyl
584	Cl	3-pyridyl	2-(methylsulfonyl)phenyl
585	Cl	3-pyridyl	4-morpholino
586	Cl	3-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
587	Cl	3-pyridyl	4-morpholinocarbonyl
588	Cl	3-pyridyl	2-methyl-1-imidazolyl
589	Cl	3-pyridyl	5-methyl-1-imidazolyl
590	Cl	3-pyridyl	2-methylsulfonyl-1-imidazolyl
591	Cl	2-pyrimidyl	2-(aminosulfonyl)phenyl
592	Cl	2-pyrimidyl	2-(methylaminosulfonyl)phenyl

593	Cl	2-pyrimidyl	1-pyrrolidinocarbonyl
594	Cl	2-pyrimidyl	2-(methylsulfonyl)phenyl
595	Cl	2-pyrimidyl	4-morpholino
596	Cl	2-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
597	Cl	2-pyrimidyl	4-morpholinocarbonyl
598	Cl	2-pyrimidyl	2-methyl-1-imidazolyl
599	Cl	2-pyrimidyl	5-methyl-1-imidazolyl
600	Cl	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
601	Cl	5-pyrimidyl	2-(aminosulfonyl)phenyl
602	Cl	5-pyrimidyl	2-(methylaminosulfonyl)phenyl
603	Cl	5-pyrimidyl	1-pyrrolidinocarbonyl
604	Cl	5-pyrimidyl	2-(methylsulfonyl)phenyl
605	Cl	5-pyrimidyl	4-morpholino
606	Cl	5-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
607	Cl	5-pyrimidyl	4-morpholinocarbonyl
608	Cl	5-pyrimidyl	2-methyl-1-imidazolyl
609	Cl	5-pyrimidyl	5-methyl-1-imidazolyl
610	Cl	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
611	Cl	2-Cl-phenyl	2-(aminosulfonyl)phenyl
612	Cl	2-Cl-phenyl	2-(methylaminosulfonyl)phenyl
613	Cl	2-Cl-phenyl	1-pyrrolidinocarbonyl
614	Cl	2-Cl-phenyl	2-(methylsulfonyl)phenyl
615	Cl	2-Cl-phenyl	4-morpholino
616	Cl	2-Cl-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
617	Cl	2-Cl-phenyl	4-morpholinocarbonyl
618	Cl	2-Cl-phenyl	2-methyl-1-imidazolyl
619	Cl	2-Cl-phenyl	5-methyl-1-imidazolyl
620	Cl	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
621	Cl	2-F-phenyl	2-(aminosulfonyl)phenyl
622	Cl	2-F-phenyl	2-(methylaminosulfonyl)phenyl
623	Cl	2-F-phenyl	1-pyrrolidinocarbonyl
624	Cl	2-F-phenyl	2-(methylsulfonyl)phenyl
625	Cl	2-F-phenyl	4-morpholino
626	Cl	2-F-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
627	Cl	2-F-phenyl	4-morpholinocarbonyl
628	Cl	2-F-phenyl	2-methyl-1-imidazolyl
629	Cl	2-F-phenyl	5-methyl-1-imidazolyl
630	Cl	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
631	Cl	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
632	Cl	2,6-diF-phenyl	2-(methylaminosulfonyl)phenyl
633	Cl	2,6-diF-phenyl	1-pyrrolidinocarbonyl
634	Cl	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
635	Cl	2,6-diF-phenyl	4-morpholino
636	Cl	2,6-diF-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
637	Cl	2,6-diF-phenyl	4-morpholinocarbonyl
638	Cl	2,6-diF-phenyl	2-methyl-1-imidazolyl
639	Cl	2,6-diF-phenyl	5-methyl-1-imidazolyl
640	Cl	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
641	F	phenyl	2-(aminosulfonyl)phenyl
642	F	phenyl	2-(methylaminosulfonyl)phenyl
643	F	phenyl	1-pyrrolidinocarbonyl
644	F	phenyl	2-(methylsulfonyl)phenyl
645	F	phenyl	4-morpholino
646	F	phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
647	F	phenyl	4-morpholinocarbonyl

648	F	phenyl	2-methyl-1-imidazolyl
649	F	phenyl	5-methyl-1-imidazolyl
650	F	phenyl	2-methylsulfonyl-1-imidazolyl
651	F	2-pyridyl	2-(aminosulfonyl)phenyl
652	F	2-pyridyl	2-(methylaminosulfonyl)phenyl
653	F	2-pyridyl	1-pyrrolidinocarbonyl
654	F	2-pyridyl	2-(methylsulfonyl)phenyl
655	F	2-pyridyl	4-morpholino
656	F	2-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
657	F	2-pyridyl	4-morpholinocarbonyl
658	F	2-pyridyl	2-methyl-1-imidazolyl
659	F	2-pyridyl	5-methyl-1-imidazolyl
660	F	2-pyridyl	2-methylsulfonyl-1-imidazolyl
661	F	3-pyridyl	2-(aminosulfonyl)phenyl
662	F	3-pyridyl	2-(methylaminosulfonyl)phenyl
663	F	3-pyridyl	1-pyrrolidinocarbonyl
664	F	3-pyridyl	2-(methylsulfonyl)phenyl
665	F	3-pyridyl	4-morpholino
666	F	3-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
667	F	3-pyridyl	4-morpholinocarbonyl
668	F	3-pyridyl	2-methyl-1-imidazolyl
669	F	3-pyridyl	5-methyl-1-imidazolyl
670	F	3-pyridyl	2-methylsulfonyl-1-imidazolyl
671	F	2-pyrimidyl	2-(aminosulfonyl)phenyl
672	F	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
673	F	2-pyrimidyl	1-pyrrolidinocarbonyl
674	F	2-pyrimidyl	2-(methylsulfonyl)phenyl
675	F	2-pyrimidyl	4-morpholino
676	F	2-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
677	F	2-pyrimidyl	4-morpholinocarbonyl
678	F	2-pyrimidyl	2-methyl-1-imidazolyl
679	F	2-pyrimidyl	5-methyl-1-imidazolyl
680	F	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
681	F	5-pyrimidyl	2-(aminosulfonyl)phenyl
682	F	5-pyrimidyl	2-(methylaminosulfonyl)phenyl
683	F	5-pyrimidyl	1-pyrrolidinocarbonyl
684	F	5-pyrimidyl	2-(methylsulfonyl)phenyl
685	F	5-pyrimidyl	4-morpholino
686	F	5-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
687	F	5-pyrimidyl	4-morpholinocarbonyl
688	F	5-pyrimidyl	2-methyl-1-imidazolyl
689	F	5-pyrimidyl	5-methyl-1-imidazolyl
690	F	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
691	F	2-Cl-phenyl	2-(aminosulfonyl)phenyl
692	F	2-Cl-phenyl	2-(methylaminosulfonyl)phenyl
693	F	2-Cl-phenyl	1-pyrrolidinocarbonyl
694	F	2-Cl-phenyl	2-(methylsulfonyl)phenyl
695	F	2-Cl-phenyl	4-morpholino
696	F	2-Cl-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
697	F	2-Cl-phenyl	4-morpholinocarbonyl
698	F	2-Cl-phenyl	2-methyl-1-imidazolyl
699	F	2-Cl-phenyl	5-methyl-1-imidazolyl
700	F	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
701	F	2-F-phenyl	2-(aminosulfonyl)phenyl
702	F	2-F-phenyl	2-(methylaminosulfonyl)phenyl

703	F	2-F-phenyl	1-pyrrolidinocarbonyl
704	F	2-F-phenyl	2-(methylsulfonyl)phenyl
705	F	2-F-phenyl	4-morpholino
706	F	2-F-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
707	F	2-F-phenyl	4-morpholinocarbonyl
708	F	2-F-phenyl	2-methyl-1-imidazolyl
709	F	2-F-phenyl	5-methyl-1-imidazolyl
710	F	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
711	F	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
712	F	2,6-diF-phenyl	2-(methylaminosulfonyl)phenyl
713	F	2,6-diF-phenyl	1-pyrrolidinocarbonyl
714	F	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
715	F	2,6-diF-phenyl	4-morpholino
716	F	2,6-diF-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
717	F	2,6-diF-phenyl	4-morpholinocarbonyl
718	F	2,6-diF-phenyl	2-methyl-1-imidazolyl
719	F	2,6-diF-phenyl	5-methyl-1-imidazolyl
720	F	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
721	CO ₂ CH ₃	phenyl	2-(aminosulfonyl)phenyl
722	CO ₂ CH ₃	phenyl	2-(methylaminosulfonyl)phenyl
723	CO ₂ CH ₃	phenyl	1-pyrrolidinocarbonyl
724	CO ₂ CH ₃	phenyl	2-(methylsulfonyl)phenyl
725	CO ₂ CH ₃	phenyl	4-morpholino
726	CO ₂ CH ₃	phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
727	CO ₂ CH ₃	phenyl	4-morpholinocarbonyl
728	CO ₂ CH ₃	phenyl	2-methyl-1-imidazolyl
729	CO ₂ CH ₃	phenyl	5-methyl-1-imidazolyl
730	CO ₂ CH ₃	phenyl	2-methylsulfonyl-1-imidazolyl
731	CO ₂ CH ₃	2-pyridyl	2-(aminosulfonyl)phenyl
732	CO ₂ CH ₃	2-pyridyl	2-(methylaminosulfonyl)phenyl
733	CO ₂ CH ₃	2-pyridyl	1-pyrrolidinocarbonyl
734	CO ₂ CH ₃	2-pyridyl	2-(methylsulfonyl)phenyl
735	CO ₂ CH ₃	2-pyridyl	4-morpholino
736	CO ₂ CH ₃	2-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
737	CO ₂ CH ₃	2-pyridyl	4-morpholinocarbonyl
738	CO ₂ CH ₃	2-pyridyl	2-methyl-1-imidazolyl
739	CO ₂ CH ₃	2-pyridyl	5-methyl-1-imidazolyl
740	CO ₂ CH ₃	2-pyridyl	2-methylsulfonyl-1-imidazolyl
741	CO ₂ CH ₃	3-pyridyl	2-(aminosulfonyl)phenyl
742	CO ₂ CH ₃	3-pyridyl	2-(methylaminosulfonyl)phenyl
743	CO ₂ CH ₃	3-pyridyl	1-pyrrolidinocarbonyl
744	CO ₂ CH ₃	3-pyridyl	2-(methylsulfonyl)phenyl
745	CO ₂ CH ₃	3-pyridyl	4-morpholino
746	CO ₂ CH ₃	3-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
747	CO ₂ CH ₃	3-pyridyl	4-morpholinocarbonyl
748	CO ₂ CH ₃	3-pyridyl	2-methyl-1-imidazolyl
749	CO ₂ CH ₃	3-pyridyl	5-methyl-1-imidazolyl
750	CO ₂ CH ₃	3-pyridyl	2-methylsulfonyl-1-imidazolyl
751	CO ₂ CH ₃	2-pyrimidyl	2-(aminosulfonyl)phenyl
752	CO ₂ CH ₃	2-pyrimidyl	2-(methylaminosulfonyl)phenyl

753	CO ₂ CH ₃	2-pyrimidyl	1-pyrrolidinocarbonyl
754	CO ₂ CH ₃	2-pyrimidyl	2-(methylsulfonyl)phenyl
755	CO ₂ CH ₃	2-pyrimidyl	4-morpholino
756	CO ₂ CH ₃	2-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
757	CO ₂ CH ₃	2-pyrimidyl	4-morpholinocarbonyl
758	CO ₂ CH ₃	2-pyrimidyl	2-methyl-1-imidazolyl
759	CO ₂ CH ₃	2-pyrimidyl	5-methyl-1-imidazolyl
760	CO ₂ CH ₃	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
761	CO ₂ CH ₃	5-pyrimidyl	2-(aminosulfonyl)phenyl
762	CO ₂ CH ₃	5-pyrimidyl	2-(methylaminosulfonyl)phenyl
763	CO ₂ CH ₃	5-pyrimidyl	1-pyrrolidinocarbonyl
764	CO ₂ CH ₃	5-pyrimidyl	2-(methylsulfonyl)phenyl
765	CO ₂ CH ₃	5-pyrimidyl	4-morpholino
766	CO ₂ CH ₃	5-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
767	CO ₂ CH ₃	5-pyrimidyl	4-morpholinocarbonyl
768	CO ₂ CH ₃	5-pyrimidyl	2-methyl-1-imidazolyl
769	CO ₂ CH ₃	5-pyrimidyl	5-methyl-1-imidazolyl
770	CO ₂ CH ₃	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
771	CO ₂ CH ₃	2-Cl-phenyl	2-(aminosulfonyl)phenyl
772	CO ₂ CH ₃	2-Cl-phenyl	2-(methylaminosulfonyl)phenyl
773	CO ₂ CH ₃	2-Cl-phenyl	1-pyrrolidinocarbonyl
774	CO ₂ CH ₃	2-Cl-phenyl	2-(methylsulfonyl)phenyl
775	CO ₂ CH ₃	2-Cl-phenyl	4-morpholino
776	CO ₂ CH ₃	2-Cl-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
777	CO ₂ CH ₃	2-Cl-phenyl	4-morpholinocarbonyl
778	CO ₂ CH ₃	2-Cl-phenyl	2-methyl-1-imidazolyl
779	CO ₂ CH ₃	2-Cl-phenyl	5-methyl-1-imidazolyl
780	CO ₂ CH ₃	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
781	CO ₂ CH ₃	2-F-phenyl	2-(aminosulfonyl)phenyl
782	CO ₂ CH ₃	2-F-phenyl	2-(methylaminosulfonyl)phenyl
783	CO ₂ CH ₃	2-F-phenyl	1-pyrrolidinocarbonyl
784	CO ₂ CH ₃	2-F-phenyl	2-(methylsulfonyl)phenyl
785	CO ₂ CH ₃	2-F-phenyl	4-morpholino
786	CO ₂ CH ₃	2-F-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
787	CO ₂ CH ₃	2-F-phenyl	4-morpholinocarbonyl
788	CO ₂ CH ₃	2-F-phenyl	2-methyl-1-imidazolyl
789	CO ₂ CH ₃	2-F-phenyl	5-methyl-1-imidazolyl
790	CO ₂ CH ₃	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
791	CO ₂ CH ₃	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
792	CO ₂ CH ₃	2,6-diF-phenyl	2-(methylaminosulfonyl)phenyl
793	CO ₂ CH ₃	2,6-diF-phenyl	1-pyrrolidinocarbonyl
794	CO ₂ CH ₃	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
795	CO ₂ CH ₃	2,6-diF-phenyl	4-morpholino
796	CO ₂ CH ₃	2,6-diF-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
797	CO ₂ CH ₃	2,6-diF-phenyl	4-morpholinocarbonyl
798	CO ₂ CH ₃	2,6-diF-phenyl	2-methyl-1-imidazolyl
799	CO ₂ CH ₃	2,6-diF-phenyl	5-methyl-1-imidazolyl
800	CO ₂ CH ₃	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl


801	CH ₂ OCH ₃	phenyl	2-(aminosulfonyl)phenyl
802	CH ₂ OCH ₃	phenyl	2-(methylaminosulfonyl)phenyl
803	CH ₂ OCH ₃	phenyl	1-pyrrolidinocarbonyl
804	CH ₂ OCH ₃	phenyl	2-(methysulfonyl)phenyl
805	CH ₂ OCH ₃	phenyl	4-morpholino
806	CH ₂ OCH ₃	phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
807	CH ₂ OCH ₃	phenyl	4-morpholinocarbonyl
808	CH ₂ OCH ₃	phenyl	2-methyl-1-imidazolyl
809	CH ₂ OCH ₃	phenyl	5-methyl-1-imidazolyl
810	CH ₂ OCH ₃	phenyl	2-methylsulfonyl-1-imidazolyl
811	CH ₂ OCH ₃	2-pyridyl	2-(aminosulfonyl)phenyl
812	CH ₂ OCH ₃	2-pyridyl	2-(methylaminosulfonyl)phenyl
813	CH ₂ OCH ₃	2-pyridyl	1-pyrrolidinocarbonyl
814	CH ₂ OCH ₃	2-pyridyl	2-(methysulfonyl)phenyl
815	CH ₂ OCH ₃	2-pyridyl	4-morpholino
816	CH ₂ OCH ₃	2-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
817	CH ₂ OCH ₃	2-pyridyl	4-morpholinocarbonyl
818	CH ₂ OCH ₃	2-pyridyl	2-methyl-1-imidazolyl
819	CH ₂ OCH ₃	2-pyridyl	5-methyl-1-imidazolyl
820	CH ₂ OCH ₃	2-pyridyl	2-methylsulfonyl-1-imidazolyl
821	CH ₂ OCH ₃	3-pyridyl	2-(aminosulfonyl)phenyl
822	CH ₂ OCH ₃	3-pyridyl	2-(methylaminosulfonyl)phenyl
823	CH ₂ OCH ₃	3-pyridyl	1-pyrrolidinocarbonyl
824	CH ₂ OCH ₃	3-pyridyl	2-(methysulfonyl)phenyl
825	CH ₂ OCH ₃	3-pyridyl	4-morpholino
826	CH ₂ OCH ₃	3-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
827	CH ₂ OCH ₃	3-pyridyl	4-morpholinocarbonyl
828	CH ₂ OCH ₃	3-pyridyl	2-methyl-1-imidazolyl
829	CH ₂ OCH ₃	3-pyridyl	5-methyl-1-imidazolyl
830	CH ₂ OCH ₃	3-pyridyl	2-methylsulfonyl-1-imidazolyl
831	CH ₂ OCH ₃	2-pyrimidyl	2-(aminosulfonyl)phenyl
832	CH ₂ OCH ₃	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
833	CH ₂ OCH ₃	2-pyrimidyl	1-pyrrolidinocarbonyl
834	CH ₂ OCH ₃	2-pyrimidyl	2-(methysulfonyl)phenyl
835	CH ₂ OCH ₃	2-pyrimidyl	4-morpholino
836	CH ₂ OCH ₃	2-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
837	CH ₂ OCH ₃	2-pyrimidyl	4-morpholinocarbonyl
838	CH ₂ OCH ₃	2-pyrimidyl	2-methyl-1-imidazolyl
839	CH ₂ OCH ₃	2-pyrimidyl	5-methyl-1-imidazolyl
840	CH ₂ OCH ₃	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
841	CH ₂ OCH ₃	5-pyrimidyl	2-(aminosulfonyl)phenyl
842	CH ₂ OCH ₃	5-pyrimidyl	2-(methylaminosulfonyl)phenyl
843	CH ₂ OCH ₃	5-pyrimidyl	1-pyrrolidinocarbonyl
844	CH ₂ OCH ₃	5-pyrimidyl	2-(methysulfonyl)phenyl
845	CH ₂ OCH ₃	5-pyrimidyl	4-morpholino
846	CH ₂ OCH ₃	5-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
847	CH ₂ OCH ₃	5-pyrimidyl	4-morpholinocarbonyl
848	CH ₂ OCH ₃	5-pyrimidyl	2-methyl-1-imidazolyl

849	CH ₂ OCH ₃	5-pyrimidyl	5-methyl-1-imidazolyl
850	CH ₂ OCH ₃	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
851	CH ₂ OCH ₃	2-Cl-phenyl	2-(aminosulfonyl)phenyl
852	CH ₂ OCH ₃	2-Cl-phenyl	2-(methylaninosulfonyl)phenyl
853	CH ₂ OCH ₃	2-Cl-phenyl	1-pyrrolidinocarbonyl
854	CH ₂ OCH ₃	2-Cl-phenyl	2-(methylsulfonyl)phenyl
855	CH ₂ OCH ₃	2-Cl-phenyl	4-morpholino
856	CH ₂ OCH ₃	2-Cl-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
857	CH ₂ OCH ₃	2-Cl-phenyl	4-morpholinocarbonyl
858	CH ₂ OCH ₃	2-Cl-phenyl	2-methyl-1-imidazolyl
859	CH ₂ OCH ₃	2-Cl-phenyl	5-methyl-1-imidazolyl
860	CH ₂ OCH ₃	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
861	CH ₂ OCH ₃	2-F-phenyl	2-(aminosulfonyl)phenyl
862	CH ₂ OCH ₃	2-F-phenyl	2-(methylaninosulfonyl)phenyl
863	CH ₂ OCH ₃	2-F-phenyl	1-pyrrolidinocarbonyl
864	CH ₂ OCH ₃	2-F-phenyl	2-(methylsulfonyl)phenyl
865	CH ₂ OCH ₃	2-F-phenyl	4-morpholino
866	CH ₂ OCH ₃	2-F-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
867	CH ₂ OCH ₃	2-F-phenyl	4-morpholinocarbonyl
868	CH ₂ OCH ₃	2-F-phenyl	2-methyl-1-imidazolyl
869	CH ₂ OCH ₃	2-F-phenyl	5-methyl-1-imidazolyl
870	CH ₂ OCH ₃	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
871	CH ₂ OCH ₃	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
872	CH ₂ OCH ₃	2,6-diF-phenyl	2-(methylaninosulfonyl)phenyl
873	CH ₂ OCH ₃	2,6-diF-phenyl	1-pyrrolidinocarbonyl
874	CH ₂ OCH ₃	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
875	CH ₂ OCH ₃	2,6-diF-phenyl	4-morpholino
876	CH ₂ OCH ₃	2,6-diF-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
877	CH ₂ OCH ₃	2,6-diF-phenyl	4-morpholinocarbonyl
878	CH ₂ OCH ₃	2,6-diF-phenyl	2-methyl-1-imidazolyl
879	CH ₂ OCH ₃	2,6-diF-phenyl	5-methyl-1-imidazolyl
880	CH ₂ OCH ₃	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
881	CONH ₂	phenyl	2-(aminosulfonyl)phenyl
882	CONH ₂	phenyl	2-(methylaninosulfonyl)phenyl
883	CONH ₂	phenyl	1-pyrrolidinocarbonyl
884	CONH ₂	phenyl	2-(methylsulfonyl)phenyl
885	CONH ₂	phenyl	4-morpholino
886	CONH ₂	phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
887	CONH ₂	phenyl	4-morpholinocarbonyl
888	CONH ₂	phenyl	2-methyl-1-imidazolyl
889	CONH ₂	phenyl	5-methyl-1-imidazolyl
890	CONH ₂	phenyl	2-methylsulfonyl-1-imidazolyl
891	CONH ₂	2-pyridyl	2-(aminosulfonyl)phenyl
892	CONH ₂	2-pyridyl	2-(methylaninosulfonyl)phenyl
893	CONH ₂	2-pyridyl	1-pyrrolidinocarbonyl
894	CONH ₂	2-pyridyl	2-(methylsulfonyl)phenyl
895	CONH ₂	2-pyridyl	4-morpholino
896	CONH ₂	2-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl

897	CONH ₂	2-pyridyl	4-morpholinocarbonyl
898	CONH ₂	2-pyridyl	2-methyl-1-imidazolyl
899	CONH ₂	2-pyridyl	5-methyl-1-imidazolyl
900	CONH ₂	2-pyridyl	2-methylsulfonyl-1-imidazolyl
901	CONH ₂	3-pyridyl	2-(aminosulfonyl)phenyl
902	CONH ₂	3-pyridyl	2-(methylaminosulfonyl)phenyl
903	CONH ₂	3-pyridyl	1-pyrrolidinocarbonyl
904	CONH ₂	3-pyridyl	2-(methylsulfonyl)phenyl
905	CONH ₂	3-pyridyl	4-morpholino
906	CONH ₂	3-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
907	CONH ₂	3-pyridyl	4-morpholinocarbonyl
908	CONH ₂	3-pyridyl	2-methyl-1-imidazolyl
909	CONH ₂	3-pyridyl	5-methyl-1-imidazolyl
910	CONH ₂	3-pyridyl	2-methylsulfonyl-1-imidazolyl
911	CONH ₂	2-pyrimidyl	2-(aminosulfonyl)phenyl
912	CONH ₂	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
913	CONH ₂	2-pyrimidyl	1-pyrrolidinocarbonyl
914	CONH ₂	2-pyrimidyl	2-(methylsulfonyl)phenyl
915	CONH ₂	2-pyrimidyl	4-morpholino
916	CONH ₂	2-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
917	CONH ₂	2-pyrimidyl	4-morpholinocarbonyl
918	CONH ₂	2-pyrimidyl	2-methyl-1-imidazolyl
919	CONH ₂	2-pyrimidyl	5-methyl-1-imidazolyl
920	CONH ₂	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
921	CONH ₂	5-pyrimidyl	2-(aminosulfonyl)phenyl
922	CONH ₂	5-pyrimidyl	2-(methylaminosulfonyl)phenyl
923	CONH ₂	5-pyrimidyl	1-pyrrolidinocarbonyl
924	CONH ₂	5-pyrimidyl	2-(methylsulfonyl)phenyl
925	CONH ₂	5-pyrimidyl	4-morpholino
926	CONH ₂	5-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
927	CONH ₂	5-pyrimidyl	4-morpholinocarbonyl
928	CONH ₂	5-pyrimidyl	2-methyl-1-imidazolyl
929	CONH ₂	5-pyrimidyl	5-methyl-1-imidazolyl
930	CONH ₂	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
931	CONH ₂	2-Cl-phenyl	2-(aminosulfonyl)phenyl
932	CONH ₂	2-Cl-phenyl	2-(methylaminosulfonyl)phenyl
933	CONH ₂	2-Cl-phenyl	1-pyrrolidinocarbonyl
934	CONH ₂	2-Cl-phenyl	2-(methylsulfonyl)phenyl
935	CONH ₂	2-Cl-phenyl	4-morpholino
936	CONH ₂	2-Cl-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
937	CONH ₂	2-Cl-phenyl	4-morpholinocarbonyl
938	CONH ₂	2-Cl-phenyl	2-methyl-1-imidazolyl
939	CONH ₂	2-Cl-phenyl	5-methyl-1-imidazolyl
940	CONH ₂	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
941	CONH ₂	2-F-phenyl	2-(aminosulfonyl)phenyl
942	CONH ₂	2-F-phenyl	2-(methylaminosulfonyl)phenyl
943	CONH ₂	2-F-phenyl	1-pyrrolidinocarbonyl
944	CONH ₂	2-F-phenyl	2-(methylsulfonyl)phenyl

945	CONH ₂	2-F-phenyl	4-morpholino
946	CONH ₂	2-F-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
947	CONH ₂	2-F-phenyl	4-morpholinocarbonyl
948	CONH ₂	2-F-phenyl	2-methyl-1-imidazolyl
949	CONH ₂	2-F-phenyl	5-methyl-1-imidazolyl
950	CONH ₂	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
951	CONH ₂	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
952	CONH ₂	2,6-diF-phenyl	2-(methylaminosulfonyl)phenyl
953	CONH ₂	2,6-diF-phenyl	1-pyrrolidinocarbonyl
954	CONH ₂	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
955	CONH ₂	2,6-diF-phenyl	4-morpholino
956	CONH ₂	2,6-diF-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
957	CONH ₂	2,6-diF-phenyl	4-morpholinocarbonyl
958	CONH ₂	2,6-diF-phenyl	2-methyl-1-imidazolyl
959	CONH ₂	2,6-diF-phenyl	5-methyl-1-imidazolyl
960	CONH ₂	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl


Table 3



Ex #	A	B
1	phenyl	2-(aminosulfonyl)phenyl
2	phenyl	2-(methylaminosulfonyl)phenyl
3	phenyl	1-pyrrolidinocarbonyl
4	phenyl	2-(methylsulfonyl)phenyl
5	phenyl	4-morpholino
6	phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
7	phenyl	4-morpholinocarbonyl
8	phenyl	2-methyl-1-imidazolyl
9	phenyl	5-methyl-1-imidazolyl
10	phenyl	2-methylsulfonyl-1-imidazolyl
11	2-pyridyl	2-(aminosulfonyl)phenyl
12	2-pyridyl	2-(methylaminosulfonyl)phenyl
13	2-pyridyl	1-pyrrolidinocarbonyl
14	2-pyridyl	2-(methylsulfonyl)phenyl
15	2-pyridyl	4-morpholino
16	2-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
17	2-pyridyl	4-morpholinocarbonyl
18	2-pyridyl	2-methyl-1-imidazolyl
19	2-pyridyl	5-methyl-1-imidazolyl
20	2-pyridyl	2-methylsulfonyl-1-imidazolyl
21	3-pyridyl	2-(aminosulfonyl)phenyl
22	3-pyridyl	2-(methylaminosulfonyl)phenyl
23	3-pyridyl	1-pyrrolidinocarbonyl
24	3-pyridyl	2-(methylsulfonyl)phenyl
25	3-pyridyl	4-morpholino
26	3-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
27	3-pyridyl	4-morpholinocarbonyl
28	3-pyridyl	2-methyl-1-imidazolyl
29	3-pyridyl	5-methyl-1-imidazolyl
30	3-pyridyl	2-methylsulfonyl-1-imidazolyl
31	2-pyrimidyl	2-(aminosulfonyl)phenyl
32	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
33	2-pyrimidyl	1-pyrrolidinocarbonyl
34	2-pyrimidyl	2-(methylsulfonyl)phenyl
35	2-pyrimidyl	4-morpholino
36	2-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
37	2-pyrimidyl	4-morpholinocarbonyl
38	2-pyrimidyl	2-methyl-1-imidazolyl
39	2-pyrimidyl	5-methyl-1-imidazolyl
40	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl

41	5-pyrimidyl	2-(aminosulfonyl)phenyl
42	5-pyrimidyl	2-(methylaninosulfonyl)phenyl
43	5-pyrimidyl	1-pyrrolidinocarbonyl
44	5-pyrimidyl	2-(methylsulfonyl)phenyl
45	5-pyrimidyl	4-morpholino
46	5-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
47	5-pyrimidyl	4-morpholinocarbonyl
48	5-pyrimidyl	2-methyl-1-imidazolyl
49	5-pyrimidyl	5-methyl-1-imidazolyl
50	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
51	2-Cl-phenyl	2-(aminosulfonyl)phenyl
52	2-Cl-phenyl	2-(methylaninosulfonyl)phenyl
53	2-Cl-phenyl	1-pyrrolidinocarbonyl
54	2-Cl-phenyl	2-(methylsulfonyl)phenyl
55	2-Cl-phenyl	4-morpholino
56	2-Cl-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
57	2-Cl-phenyl	4-morpholinocarbonyl
58	2-Cl-phenyl	2-methyl-1-imidazolyl
59	2-Cl-phenyl	5-methyl-1-imidazolyl
60	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
61	2-F-phenyl	2-(aminosulfonyl)phenyl
62	2-F-phenyl	2-(methylaninosulfonyl)phenyl
63	2-F-phenyl	1-pyrrolidinocarbonyl
64	2-F-phenyl	2-(methylsulfonyl)phenyl
65	2-F-phenyl	4-morpholino
66	2-F-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
67	2-F-phenyl	4-morpholinocarbonyl
68	2-F-phenyl	2-methyl-1-imidazolyl
69	2-F-phenyl	5-methyl-1-imidazolyl
70	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
71	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
72	2,6-diF-phenyl	2-(methylaninosulfonyl)phenyl
73	2,6-diF-phenyl	1-pyrrolidinocarbonyl
74	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
75	2,6-diF-phenyl	4-morpholino
76	2,6-diF-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
77	2,6-diF-phenyl	4-morpholinocarbonyl
78	2,6-diF-phenyl	2-methyl-1-imidazolyl
79	2,6-diF-phenyl	5-methyl-1-imidazolyl
80	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl

Table 4



For each example, DE is:

- (A) pyridin-4-yl-CH₂,
- (B) 2-amino-pyrimidin-4-yl,
- (C) 6-amino-pyridin-2-yl,
- (D) 3-amidino-4-F-phenyl, or
- (E) N-amidino-3-piperidinyl.

Ex #	R ^{1a}	A	B
1	CH ₃	phenyl	2-(aminosulfonyl)phenyl
2	CH ₃	phenyl	2-(methylaminosulfonyl)phenyl
3	CH ₃	phenyl	1-pyrrolidinocarbonyl
4	CH ₃	phenyl	2-(methylsulfonyl)phenyl
5	CH ₃	phenyl	4-morpholino
6	CH ₃	phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
7	CH ₃	phenyl	4-morpholinocarbonyl
8	CH ₃	phenyl	2-methyl-1-imidazolyl
9	CH ₃	phenyl	5-methyl-1-imidazolyl
10	CH ₃	phenyl	2-methylsulfonyl-1-imidazolyl
11	CH ₃	2-pyridyl	2-(aminosulfonyl)phenyl
12	CH ₃	2-pyridyl	2-(methylaminosulfonyl)phenyl
13	CH ₃	2-pyridyl	1-pyrrolidinocarbonyl
14	CH ₃	2-pyridyl	2-(methylsulfonyl)phenyl
15	CH ₃	2-pyridyl	4-morpholino
16	CH ₃	2-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
17	CH ₃	2-pyridyl	4-morpholinocarbonyl
18	CH ₃	2-pyridyl	2-methyl-1-imidazolyl
19	CH ₃	2-pyridyl	5-methyl-1-imidazolyl
20	CH ₃	2-pyridyl	2-methylsulfonyl-1-imidazolyl
21	CH ₃	3-pyridyl	2-(aminosulfonyl)phenyl
22	CH ₃	3-pyridyl	2-(methylaminosulfonyl)phenyl
23	CH ₃	3-pyridyl	1-pyrrolidinocarbonyl
24	CH ₃	3-pyridyl	2-(methylsulfonyl)phenyl
25	CH ₃	3-pyridyl	4-morpholino
26	CH ₃	3-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
27	CH ₃	3-pyridyl	4-morpholinocarbonyl
28	CH ₃	3-pyridyl	2-methyl-1-imidazolyl
29	CH ₃	3-pyridyl	5-methyl-1-imidazolyl
30	CH ₃	3-pyridyl	2-methylsulfonyl-1-imidazolyl
31	CH ₃	2-pyrimidyl	2-(aminosulfonyl)phenyl
32	CH ₃	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
33	CH ₃	2-pyrimidyl	1-pyrrolidinocarbonyl
34	CH ₃	2-pyrimidyl	2-(methylsulfonyl)phenyl
35	CH ₃	2-pyrimidyl	4-morpholino
36	CH ₃	2-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
37	CH ₃	2-pyrimidyl	4-morpholinocarbonyl
38	CH ₃	2-pyrimidyl	2-methyl-1-imidazolyl
39	CH ₃	2-pyrimidyl	5-methyl-1-imidazolyl
40	CH ₃	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl

41	CH ₃	5-pyrimidyl	2-(aminosulfonyl)phenyl
42	CH ₃	5-pyrimidyl	2-(methylaminosulfonyl)phenyl
43	CH ₃	5-pyrimidyl	1-pyrrolidinocarbonyl
44	CH ₃	5-pyrimidyl	2-(methylsulfonyl)phenyl
45	CH ₃	5-pyrimidyl	4-morpholino
46	CH ₃	5-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
47	CH ₃	5-pyrimidyl	4-morpholinocarbonyl
48	CH ₃	5-pyrimidyl	2-methyl-1-imidazolyl
49	CH ₃	5-pyrimidyl	5-methyl-1-imidazolyl
50	CH ₃	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
51	CH ₃	2-Cl-phenyl	2-(aminosulfonyl)phenyl
52	CH ₃	2-Cl-phenyl	2-(methylaminosulfonyl)phenyl
53	CH ₃	2-Cl-phenyl	1-pyrrolidinocarbonyl
54	CH ₃	2-Cl-phenyl	2-(methylsulfonyl)phenyl
55	CH ₃	2-Cl-phenyl	4-morpholino
56	CH ₃	2-Cl-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
57	CH ₃	2-Cl-phenyl	4-morpholinocarbonyl
58	CH ₃	2-Cl-phenyl	2-methyl-1-imidazolyl
59	CH ₃	2-Cl-phenyl	5-methyl-1-imidazolyl
60	CH ₃	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
61	CH ₃	2-F-phenyl	2-(aminosulfonyl)phenyl
62	CH ₃	2-F-phenyl	2-(methylaminosulfonyl)phenyl
63	CH ₃	2-F-phenyl	1-pyrrolidinocarbonyl
64	CH ₃	2-F-phenyl	2-(methylsulfonyl)phenyl
65	CH ₃	2-F-phenyl	4-morpholino
66	CH ₃	2-F-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
67	CH ₃	2-F-phenyl	4-morpholinocarbonyl
68	CH ₃	2-F-phenyl	2-methyl-1-imidazolyl
69	CH ₃	2-F-phenyl	5-methyl-1-imidazolyl
70	CH ₃	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
71	CH ₃	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
72	CH ₃	2,6-diF-phenyl	2-(methylaminosulfonyl)phenyl
73	CH ₃	2,6-diF-phenyl	1-pyrrolidinocarbonyl
74	CH ₃	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
75	CH ₃	2,6-diF-phenyl	4-morpholino
76	CH ₃	2,6-diF-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
77	CH ₃	2,6-diF-phenyl	4-morpholinocarbonyl
78	CH ₃	2,6-diF-phenyl	2-methyl-1-imidazolyl
79	CH ₃	2,6-diF-phenyl	5-methyl-1-imidazolyl
80	CH ₃	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
81	CH ₂ CH ₃	phenyl	2-(aminosulfonyl)phenyl
82	CH ₂ CH ₃	phenyl	2-(methylaminosulfonyl)phenyl
83	CH ₂ CH ₃	phenyl	1-pyrrolidinocarbonyl
84	CH ₂ CH ₃	phenyl	2-(methylsulfonyl)phenyl
85	CH ₂ CH ₃	phenyl	4-morpholino
86	CH ₂ CH ₃	phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
87	CH ₂ CH ₃	phenyl	4-morpholinocarbonyl
88	CH ₂ CH ₃	phenyl	2-methyl-1-imidazolyl

89	CH ₂ CH ₃	phenyl	5-methyl-1-imidazolyl
90	CH ₂ CH ₃	phenyl	2-methylsulfonyl-1-imidazolyl
91	CH ₂ CH ₃	2-pyridyl	2-(aminosulfonyl)phenyl
92	CH ₂ CH ₃	2-pyridyl	2-(methylaninosulfonyl)phenyl
93	CH ₂ CH ₃	2-pyridyl	1-pyrrolidinocarbonyl
94	CH ₂ CH ₃	2-pyridyl	2-(methylsulfonyl)phenyl
95	CH ₂ CH ₃	2-pyridyl	4-morpholino
96	CH ₂ CH ₃	2-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
97	CH ₂ CH ₃	2-pyridyl	4-morpholinocarbonyl
98	CH ₂ CH ₃	2-pyridyl	2-methyl-1-imidazolyl
99	CH ₂ CH ₃	2-pyridyl	5-methyl-1-imidazolyl
100	CH ₂ CH ₃	2-pyridyl	2-methylsulfonyl-1-imidazolyl
101	CH ₂ CH ₃	3-pyridyl	2-(aminosulfonyl)phenyl
102	CH ₂ CH ₃	3-pyridyl	2-(methylaninosulfonyl)phenyl
103	CH ₂ CH ₃	3-pyridyl	1-pyrrolidinocarbonyl
104	CH ₂ CH ₃	3-pyridyl	2-(methylsulfonyl)phenyl
105	CH ₂ CH ₃	3-pyridyl	4-morpholino
106	CH ₂ CH ₃	3-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
107	CH ₂ CH ₃	3-pyridyl	4-morpholinocarbonyl
108	CH ₂ CH ₃	3-pyridyl	2-methyl-1-imidazolyl
109	CH ₂ CH ₃	3-pyridyl	5-methyl-1-imidazolyl
110	CH ₂ CH ₃	3-pyridyl	2-methylsulfonyl-1-imidazolyl
111	CH ₂ CH ₃	2-pyrimidyl	2-(aminosulfonyl)phenyl
112	CH ₂ CH ₃	2-pyrimidyl	2-(methylaninosulfonyl)phenyl
113	CH ₂ CH ₃	2-pyrimidyl	1-pyrrolidinocarbonyl
114	CH ₂ CH ₃	2-pyrimidyl	2-(methylsulfonyl)phenyl
115	CH ₂ CH ₃	2-pyrimidyl	4-morpholino
116	CH ₂ CH ₃	2-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
117	CH ₂ CH ₃	2-pyrimidyl	4-morpholinocarbonyl
118	CH ₂ CH ₃	2-pyrimidyl	2-methyl-1-imidazolyl
119	CH ₂ CH ₃	2-pyrimidyl	5-methyl-1-imidazolyl
120	CH ₂ CH ₃	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
121	CH ₂ CH ₃	5-pyrimidyl	2-(aminosulfonyl)phenyl
122	CH ₂ CH ₃	5-pyrimidyl	2-(methylaninosulfonyl)phenyl
123	CH ₂ CH ₃	5-pyrimidyl	1-pyrrolidinocarbonyl
124	CH ₂ CH ₃	5-pyrimidyl	2-(methylsulfonyl)phenyl
125	CH ₂ CH ₃	5-pyrimidyl	4-morpholino
126	CH ₂ CH ₃	5-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
127	CH ₂ CH ₃	5-pyrimidyl	4-morpholinocarbonyl
128	CH ₂ CH ₃	5-pyrimidyl	2-methyl-1-imidazolyl
129	CH ₂ CH ₃	5-pyrimidyl	5-methyl-1-imidazolyl
130	CH ₂ CH ₃	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
131	CH ₂ CH ₃	2-Cl-phenyl	2-(aminosulfonyl)phenyl
132	CH ₂ CH ₃	2-Cl-phenyl	2-(methylaninosulfonyl)phenyl
133	CH ₂ CH ₃	2-Cl-phenyl	1-pyrrolidinocarbonyl
134	CH ₂ CH ₃	2-Cl-phenyl	2-(methylsulfonyl)phenyl
135	CH ₂ CH ₃	2-Cl-phenyl	4-morpholino
136	CH ₂ CH ₃	2-Cl-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl

137	CH ₂ CH ₃	2-Cl-phenyl	4-morpholinocarbonyl
138	CH ₂ CH ₃	2-Cl-phenyl	2-methyl-1-imidazolyl
139	CH ₂ CH ₃	2-Cl-phenyl	5-methyl-1-imidazolyl
140	CH ₂ CH ₃	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
141	CH ₂ CH ₃	2-F-phenyl	2-(aminosulfonyl)phenyl
142	CH ₂ CH ₃	2-F-phenyl	2-(methylaminosulfonyl)phenyl
143	CH ₂ CH ₃	2-F-phenyl	1-pyrrolidinocarbonyl
144	CH ₂ CH ₃	2-F-phenyl	2-(methylsulfonyl)phenyl
145	CH ₂ CH ₃	2-F-phenyl	4-morpholino
146	CH ₂ CH ₃	2-F-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
147	CH ₂ CH ₃	2-F-phenyl	4-morpholinocarbonyl
148	CH ₂ CH ₃	2-F-phenyl	2-methyl-1-imidazolyl
149	CH ₂ CH ₃	2-F-phenyl	5-methyl-1-imidazolyl
150	CH ₂ CH ₃	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
151	CH ₂ CH ₃	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
152	CH ₂ CH ₃	2,6-diF-phenyl	2-(methylaminosulfonyl)phenyl
153	CH ₂ CH ₃	2,6-diF-phenyl	1-pyrrolidinocarbonyl
154	CH ₂ CH ₃	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
155	CH ₂ CH ₃	2,6-diF-phenyl	4-morpholino
156	CH ₂ CH ₃	2,6-diF-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
157	CH ₂ CH ₃	2,6-diF-phenyl	4-morpholinocarbonyl
158	CH ₂ CH ₃	2,6-diF-phenyl	2-methyl-1-imidazolyl
159	CH ₂ CH ₃	2,6-diF-phenyl	5-methyl-1-imidazolyl
160	CH ₂ CH ₃	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
161	CF ₃	phenyl	2-(aminosulfonyl)phenyl
162	CF ₃	phenyl	2-(methylaminosulfonyl)phenyl
163	CF ₃	phenyl	1-pyrrolidinocarbonyl
164	CF ₃	phenyl	2-(methylsulfonyl)phenyl
165	CF ₃	phenyl	4-morpholino
166	CF ₃	phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
167	CF ₃	phenyl	4-morpholinocarbonyl
168	CF ₃	phenyl	2-methyl-1-imidazolyl
169	CF ₃	phenyl	5-methyl-1-imidazolyl
170	CF ₃	phenyl	2-methylsulfonyl-1-imidazolyl
171	CF ₃	2-pyridyl	2-(aminosulfonyl)phenyl
172	CF ₃	2-pyridyl	2-(methylaminosulfonyl)phenyl
173	CF ₃	2-pyridyl	1-pyrrolidinocarbonyl
174	CF ₃	2-pyridyl	2-(methylsulfonyl)phenyl
175	CF ₃	2-pyridyl	4-morpholino
176	CF ₃	2-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
177	CF ₃	2-pyridyl	4-morpholinocarbonyl
178	CF ₃	2-pyridyl	2-methyl-1-imidazolyl
179	CF ₃	2-pyridyl	5-methyl-1-imidazolyl
180	CF ₃	2-pyridyl	2-methylsulfonyl-1-imidazolyl
181	CF ₃	3-pyridyl	2-(aminosulfonyl)phenyl
182	CF ₃	3-pyridyl	2-(methylaminosulfonyl)phenyl
183	CF ₃	3-pyridyl	1-pyrrolidinocarbonyl
184	CF ₃	3-pyridyl	2-(methylsulfonyl)phenyl

185	CF ₃	3-pyridyl	4-morpholino
186	CF ₃	3-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
187	CF ₃	3-pyridyl	4-morpholinocarbonyl
188	CF ₃	3-pyridyl	2-methyl-1-imidazolyl
189	CF ₃	3-pyridyl	5-methyl-1-imidazolyl
190	CF ₃	3-pyridyl	2-methylsulfonyl-1-imidazolyl
191	CF ₃	2-pyrimidyl	2-(aminosulfonyl)phenyl
192	CF ₃	2-pyrimidyl	2-(methylaninosulfonyl)phenyl
193	CF ₃	2-pyrimidyl	1-pyrrolidinocarbonyl
194	CF ₃	2-pyrimidyl	2-(methylsulfonyl)phenyl
195	CF ₃	2-pyrimidyl	4-morpholino
196	CF ₃	2-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
197	CF ₃	2-pyrimidyl	4-morpholinocarbonyl
198	CF ₃	2-pyrimidyl	2-methyl-1-imidazolyl
199	CF ₃	2-pyrimidyl	5-methyl-1-imidazolyl
200	CF ₃	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
201	CF ₃	5-pyrimidyl	2-(aminosulfonyl)phenyl
202	CF ₃	5-pyrimidyl	2-(methylaninosulfonyl)phenyl
203	CF ₃	5-pyrimidyl	1-pyrrolidinocarbonyl
204	CF ₃	5-pyrimidyl	2-(methylsulfonyl)phenyl
205	CF ₃	5-pyrimidyl	4-morpholino
206	CF ₃	5-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
207	CF ₃	5-pyrimidyl	4-morpholinocarbonyl
208	CF ₃	5-pyrimidyl	2-methyl-1-imidazolyl
209	CF ₃	5-pyrimidyl	5-methyl-1-imidazolyl
210	CF ₃	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
211	CF ₃	2-Cl-phenyl	2-(aminosulfonyl)phenyl
212	CF ₃	2-Cl-phenyl	2-(methylaninosulfonyl)phenyl
213	CF ₃	2-Cl-phenyl	1-pyrrolidinocarbonyl
214	CF ₃	2-Cl-phenyl	2-(methylsulfonyl)phenyl
215	CF ₃	2-Cl-phenyl	4-morpholino
216	CF ₃	2-Cl-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
217	CF ₃	2-Cl-phenyl	4-morpholinocarbonyl
218	CF ₃	2-Cl-phenyl	2-methyl-1-imidazolyl
219	CF ₃	2-Cl-phenyl	5-methyl-1-imidazolyl
220	CF ₃	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
221	CF ₃	2-F-phenyl	2-(aminosulfonyl)phenyl
222	CF ₃	2-F-phenyl	2-(methylaninosulfonyl)phenyl
223	CF ₃	2-F-phenyl	1-pyrrolidinocarbonyl
224	CF ₃	2-F-phenyl	2-(methylsulfonyl)phenyl
225	CF ₃	2-F-phenyl	4-morpholino
226	CF ₃	2-F-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
227	CF ₃	2-F-phenyl	4-morpholinocarbonyl
228	CF ₃	2-F-phenyl	2-methyl-1-imidazolyl
229	CF ₃	2-F-phenyl	5-methyl-1-imidazolyl
230	CF ₃	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
231	CF ₃	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
232	CF ₃	2,6-diF-phenyl	2-(methylaninosulfonyl)phenyl

233	CF ₃	2,6-diF-phenyl	1-pyrrolidinocarbonyl
234	CF ₃	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
235	CF ₃	2,6-diF-phenyl	4-morpholino
236	CF ₃	2,6-diF-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
237	CF ₃	2,6-diF-phenyl	4-morpholinocarbonyl
238	CF ₃	2,6-diF-phenyl	2-methyl-1-imidazolyl
239	CF ₃	2,6-diF-phenyl	5-methyl-1-imidazolyl
240	CF ₃	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
241	SCH ₃	phenyl	2-(aminosulfonyl)phenyl
242	SCH ₃	phenyl	2-(methylaminosulfonyl)phenyl
243	SCH ₃	phenyl	1-pyrrolidinocarbonyl
244	SCH ₃	phenyl	2-(methylsulfonyl)phenyl
245	SCH ₃	phenyl	4-morpholino
246	SCH ₃	phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
247	SCH ₃	phenyl	4-morpholinocarbonyl
248	SCH ₃	phenyl	2-methyl-1-imidazolyl
249	SCH ₃	phenyl	5-methyl-1-imidazolyl
250	SCH ₃	phenyl	2-methylsulfonyl-1-imidazolyl
251	SCH ₃	2-pyridyl	2-(aminosulfonyl)phenyl
252	SCH ₃	2-pyridyl	2-(methylaminosulfonyl)phenyl
253	SCH ₃	2-pyridyl	1-pyrrolidinocarbonyl
254	SCH ₃	2-pyridyl	2-(methylsulfonyl)phenyl
255	SCH ₃	2-pyridyl	4-morpholino
256	SCH ₃	2-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
257	SCH ₃	2-pyridyl	4-morpholinocarbonyl
258	SCH ₃	2-pyridyl	2-methyl-1-imidazolyl
259	SCH ₃	2-pyridyl	5-methyl-1-imidazolyl
260	SCH ₃	2-pyridyl	2-methylsulfonyl-1-imidazolyl
261	SCH ₃	3-pyridyl	2-(aminosulfonyl)phenyl
262	SCH ₃	3-pyridyl	2-(methylaminosulfonyl)phenyl
263	SCH ₃	3-pyridyl	1-pyrrolidinocarbonyl
264	SCH ₃	3-pyridyl	2-(methylsulfonyl)phenyl
265	SCH ₃	3-pyridyl	4-morpholino
266	SCH ₃	3-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
267	SCH ₃	3-pyridyl	4-morpholinocarbonyl
268	SCH ₃	3-pyridyl	2-methyl-1-imidazolyl
269	SCH ₃	3-pyridyl	5-methyl-1-imidazolyl
270	SCH ₃	3-pyridyl	2-methylsulfonyl-1-imidazolyl
271	SCH ₃	2-pyrimidyl	2-(aminosulfonyl)phenyl
272	SCH ₃	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
273	SCH ₃	2-pyrimidyl	1-pyrrolidinocarbonyl
274	SCH ₃	2-pyrimidyl	2-(methylsulfonyl)phenyl
275	SCH ₃	2-pyrimidyl	4-morpholino
276	SCH ₃	2-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
277	SCH ₃	2-pyrimidyl	4-morpholinocarbonyl
278	SCH ₃	2-pyrimidyl	2-methyl-1-imidazolyl
279	SCH ₃	2-pyrimidyl	5-methyl-1-imidazolyl
280	SCH ₃	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl

281	SCH ₃	5-pyrimidyl	2-(aminosulfonyl)phenyl
282	SCH ₃	5-pyrimidyl	2-(methylaninosulfonyl)phenyl
283	SCH ₃	5-pyrimidyl	1-pyrrolidinocarbonyl
284	SCH ₃	5-pyrimidyl	2-(methysulfonyl)phenyl
285	SCH ₃	5-pyrimidyl	4-morpholino
286	SCH ₃	5-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
287	SCH ₃	5-pyrimidyl	4-morpholinocarbonyl
288	SCH ₃	5-pyrimidyl	2-methyl-1-imidazolyl
289	SCH ₃	5-pyrimidyl	5-methyl-1-imidazolyl
290	SCH ₃	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
291	SCH ₃	2-Cl-phenyl	2-(aminosulfonyl)phenyl
292	SCH ₃	2-Cl-phenyl	2-(methylaninosulfonyl)phenyl
293	SCH ₃	2-Cl-phenyl	1-pyrrolidinocarbonyl
294	SCH ₃	2-Cl-phenyl	2-(methysulfonyl)phenyl
295	SCH ₃	2-Cl-phenyl	4-morpholino
296	SCH ₃	2-Cl-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
297	SCH ₃	2-Cl-phenyl	4-morpholinocarbonyl
298	SCH ₃	2-Cl-phenyl	2-methyl-1-imidazolyl
299	SCH ₃	2-Cl-phenyl	5-methyl-1-imidazolyl
300	SCH ₃	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
301	SCH ₃	2-F-phenyl	2-(aminosulfonyl)phenyl
302	SCH ₃	2-F-phenyl	2-(methylaninosulfonyl)phenyl
303	SCH ₃	2-F-phenyl	1-pyrrolidinocarbonyl
304	SCH ₃	2-F-phenyl	2-(methysulfonyl)phenyl
305	SCH ₃	2-F-phenyl	4-morpholino
306	SCH ₃	2-F-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
307	SCH ₃	2-F-phenyl	4-morpholinocarbonyl
308	SCH ₃	2-F-phenyl	2-methyl-1-imidazolyl
309	SCH ₃	2-F-phenyl	5-methyl-1-imidazolyl
310	SCH ₃	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
311	SCH ₃	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
312	SCH ₃	2,6-diF-phenyl	2-(methylaninosulfonyl)phenyl
313	SCH ₃	2,6-diF-phenyl	1-pyrrolidinocarbonyl
314	SCH ₃	2,6-diF-phenyl	2-(methysulfonyl)phenyl
315	SCH ₃	2,6-diF-phenyl	4-morpholino
316	SCH ₃	2,6-diF-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
317	SCH ₃	2,6-diF-phenyl	4-morpholinocarbonyl
318	SCH ₃	2,6-diF-phenyl	2-methyl-1-imidazolyl
319	SCH ₃	2,6-diF-phenyl	5-methyl-1-imidazolyl
320	SCH ₃	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
321	SOCH ₃	phenyl	2-(aminosulfonyl)phenyl
322	SOCH ₃	phenyl	2-(methylaninosulfonyl)phenyl
323	SOCH ₃	phenyl	1-pyrrolidinocarbonyl
324	SOCH ₃	phenyl	2-(methysulfonyl)phenyl
325	SOCH ₃	phenyl	4-morpholino
326	SOCH ₃	phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
327	SOCH ₃	phenyl	4-morpholinocarbonyl
328	SOCH ₃	phenyl	2-methyl-1-imidazolyl

WO 98/28282

329	SOCH ₃	phenyl	5-methyl-1-imidazolyl
330	SOCH ₃	phenyl	2-methylsulfonyl-1-imidazolyl
331	SOCH ₃	2-pyridyl	2-(aminosulfonyl)phenyl
332	SOCH ₃	2-pyridyl	2-(methylaminosulfonyl)phenyl
333	SOCH ₃	2-pyridyl	1-pyrrolidinocarbonyl
334	SOCH ₃	2-pyridyl	2-(methylsulfonyl)phenyl
335	SOCH ₃	2-pyridyl	4-morpholino
336	SOCH ₃	2-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
337	SOCH ₃	2-pyridyl	4-morpholinocarbonyl
338	SOCH ₃	2-pyridyl	2-methyl-1-imidazolyl
339	SOCH ₃	2-pyridyl	5-methyl-1-imidazolyl
340	SOCH ₃	2-pyridyl	2-methylsulfonyl-1-imidazolyl
341	SOCH ₃	3-pyridyl	2-(aminosulfonyl)phenyl
342	SOCH ₃	3-pyridyl	2-(methylaminosulfonyl)phenyl
343	SOCH ₃	3-pyridyl	1-pyrrolidinocarbonyl
344	SOCH ₃	3-pyridyl	2-(methylsulfonyl)phenyl
345	SOCH ₃	3-pyridyl	4-morpholino
346	SOCH ₃	3-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
347	SOCH ₃	3-pyridyl	4-morpholinocarbonyl
348	SOCH ₃	3-pyridyl	2-methyl-1-imidazolyl
349	SOCH ₃	3-pyridyl	5-methyl-1-imidazolyl
350	SOCH ₃	3-pyridyl	2-methylsulfonyl-1-imidazolyl
351	SOCH ₃	2-pyrimidyl	2-(aminosulfonyl)phenyl
352	SOCH ₃	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
353	SOCH ₃	2-pyrimidyl	1-pyrrolidinocarbonyl
354	SOCH ₃	2-pyrimidyl	2-(methylsulfonyl)phenyl
355	SOCH ₃	2-pyrimidyl	4-morpholino
356	SOCH ₃	2-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
357	SOCH ₃	2-pyrimidyl	4-morpholinocarbonyl
358	SOCH ₃	2-pyrimidyl	2-methyl-1-imidazolyl
359	SOCH ₃	2-pyrimidyl	5-methyl-1-imidazolyl
360	SOCH ₃	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
361	SOCH ₃	5-pyrimidyl	2-(aminosulfonyl)phenyl
362	SOCH ₃	5-pyrimidyl	2-(methylaminosulfonyl)phenyl
363	SOCH ₃	5-pyrimidyl	1-pyrrolidinocarbonyl
364	SOCH ₃	5-pyrimidyl	2-(methylsulfonyl)phenyl
365	SOCH ₃	5-pyrimidyl	4-morpholino
366	SOCH ₃	5-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
367	SOCH ₃	5-pyrimidyl	4-morpholinocarbonyl
368	SOCH ₃	5-pyrimidyl	2-methyl-1-imidazolyl
369	SOCH ₃	5-pyrimidyl	5-methyl-1-imidazolyl
370	SOCH ₃	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
371	SOCH ₃	2-Cl-phenyl	2-(aminosulfonyl)phenyl
372	SOCH ₃	2-Cl-phenyl	2-(methylaminosulfonyl)phenyl
373	SOCH ₃	2-Cl-phenyl	1-pyrrolidinocarbonyl
374	SOCH ₃	2-Cl-phenyl	2-(methylsulfonyl)phenyl
375	SOCH ₃	2-Cl-phenyl	4-morpholino
376	SOCH ₃	2-Cl-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl

377	SOCH ₃	2-Cl-phenyl	4-morpholinocarbonyl
378	SOCH ₃	2-Cl-phenyl	2-methyl-1-imidazolyl
379	SOCH ₃	2-Cl-phenyl	5-methyl-1-imidazolyl
380	SOCH ₃	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
381	SOCH ₃	2-F-phenyl	2-(aminosulfonyl)phenyl
382	SOCH ₃	2-F-phenyl	2-(methylaninosulfonyl)phenyl
383	SOCH ₃	2-F-phenyl	1-pyrrolidinocarbonyl
384	SOCH ₃	2-F-phenyl	2-(methylsulfonyl)phenyl
385	SOCH ₃	2-F-phenyl	4-morpholino
386	SOCH ₃	2-F-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
387	SOCH ₃	2-F-phenyl	4-morpholinocarbonyl
388	SOCH ₃	2-F-phenyl	2-methyl-1-imidazolyl
389	SOCH ₃	2-F-phenyl	5-methyl-1-imidazolyl
390	SOCH ₃	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
391	SOCH ₃	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
392	SOCH ₃	2,6-diF-phenyl	2-(methylaninosulfonyl)phenyl
393	SOCH ₃	2,6-diF-phenyl	1-pyrrolidinocarbonyl
394	SOCH ₃	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
395	SOCH ₃	2,6-diF-phenyl	4-morpholino
396	SOCH ₃	2,6-diF-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
397	SOCH ₃	2,6-diF-phenyl	4-morpholinocarbonyl
398	SOCH ₃	2,6-diF-phenyl	2-methyl-1-imidazolyl
399	SOCH ₃	2,6-diF-phenyl	5-methyl-1-imidazolyl
400	SOCH ₃	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
401	SO ₂ CH ₃	phenyl	2-(aminosulfonyl)phenyl
402	SO ₂ CH ₃	phenyl	2-(methylaninosulfonyl)phenyl
403	SO ₂ CH ₃	phenyl	1-pyrrolidinocarbonyl
404	SO ₂ CH ₃	phenyl	2-(methylsulfonyl)phenyl
405	SO ₂ CH ₃	phenyl	4-morpholino
406	SO ₂ CH ₃	phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
407	SO ₂ CH ₃	phenyl	4-morpholinocarbonyl
408	SO ₂ CH ₃	phenyl	2-methyl-1-imidazolyl
409	SO ₂ CH ₃	phenyl	5-methyl-1-imidazolyl
410	SO ₂ CH ₃	phenyl	2-methylsulfonyl-1-imidazolyl
411	SO ₂ CH ₃	2-pyridyl	2-(aminosulfonyl)phenyl
412	SO ₂ CH ₃	2-pyridyl	2-(methylaninosulfonyl)phenyl
413	SO ₂ CH ₃	2-pyridyl	1-pyrrolidinocarbonyl
414	SO ₂ CH ₃	2-pyridyl	2-(methylsulfonyl)phenyl
415	SO ₂ CH ₃	2-pyridyl	4-morpholino
416	SO ₂ CH ₃	2-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
417	SO ₂ CH ₃	2-pyridyl	4-morpholinocarbonyl
418	SO ₂ CH ₃	2-pyridyl	2-methyl-1-imidazolyl
419	SO ₂ CH ₃	2-pyridyl	5-methyl-1-imidazolyl
420	SO ₂ CH ₃	2-pyridyl	2-methylsulfonyl-1-imidazolyl
421	SO ₂ CH ₃	3-pyridyl	2-(aminosulfonyl)phenyl
422	SO ₂ CH ₃	3-pyridyl	2-(methylaninosulfonyl)phenyl
423	SO ₂ CH ₃	3-pyridyl	1-pyrrolidinocarbonyl
424	SO ₂ CH ₃	3-pyridyl	2-(methylsulfonyl)phenyl

425	SO ₂ CH ₃	3-pyridyl	4-morpholino
426	SO ₂ CH ₃	3-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
427	SO ₂ CH ₃	3-pyridyl	4-morpholinocarbonyl
428	SO ₂ CH ₃	3-pyridyl	2-methyl-1-imidazolyl
429	SO ₂ CH ₃	3-pyridyl	5-methyl-1-imidazolyl
430	SO ₂ CH ₃	3-pyridyl	2-methylsulfonyl-1-imidazolyl
431	SO ₂ CH ₃	2-pyrimidyl	2-(aminosulfonyl)phenyl
432	SO ₂ CH ₃	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
433	SO ₂ CH ₃	2-pyrimidyl	1-pyrrolidinocarbonyl
434	SO ₂ CH ₃	2-pyrimidyl	2-(methylsulfonyl)phenyl
435	SO ₂ CH ₃	2-pyrimidyl	4-morpholino
436	SO ₂ CH ₃	2-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
437	SO ₂ CH ₃	2-pyrimidyl	4-morpholinocarbonyl
438	SO ₂ CH ₃	2-pyrimidyl	2-methyl-1-imidazolyl
439	SO ₂ CH ₃	2-pyrimidyl	5-methyl-1-imidazolyl
440	SO ₂ CH ₃	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
441	SO ₂ CH ₃	5-pyrimidyl	2-(aminosulfonyl)phenyl
442	SO ₂ CH ₃	5-pyrimidyl	2-(methylaminosulfonyl)phenyl
443	SO ₂ CH ₃	5-pyrimidyl	1-pyrrolidinocarbonyl
444	SO ₂ CH ₃	5-pyrimidyl	2-(methylsulfonyl)phenyl
445	SO ₂ CH ₃	5-pyrimidyl	4-morpholino
446	SO ₂ CH ₃	5-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
447	SO ₂ CH ₃	5-pyrimidyl	4-morpholinocarbonyl
448	SO ₂ CH ₃	5-pyrimidyl	2-methyl-1-imidazolyl
449	SO ₂ CH ₃	5-pyrimidyl	5-methyl-1-imidazolyl
450	SO ₂ CH ₃	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
451	SO ₂ CH ₃	2-Cl-phenyl	2-(aminosulfonyl)phenyl
452	SO ₂ CH ₃	2-Cl-phenyl	2-(methylaminosulfonyl)phenyl
453	SO ₂ CH ₃	2-Cl-phenyl	1-pyrrolidinocarbonyl
454	SO ₂ CH ₃	2-Cl-phenyl	2-(methylsulfonyl)phenyl
455	SO ₂ CH ₃	2-Cl-phenyl	4-morpholino
456	SO ₂ CH ₃	2-Cl-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
457	SO ₂ CH ₃	2-Cl-phenyl	4-morpholinocarbonyl
458	SO ₂ CH ₃	2-Cl-phenyl	2-methyl-1-imidazolyl
459	SO ₂ CH ₃	2-Cl-phenyl	5-methyl-1-imidazolyl
460	SO ₂ CH ₃	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
461	SO ₂ CH ₃	2-F-phenyl	2-(aminosulfonyl)phenyl
462	SO ₂ CH ₃	2-F-phenyl	2-(methylaminosulfonyl)phenyl
463	SO ₂ CH ₃	2-F-phenyl	1-pyrrolidinocarbonyl
464	SO ₂ CH ₃	2-F-phenyl	2-(methylsulfonyl)phenyl
465	SO ₂ CH ₃	2-F-phenyl	4-morpholino
466	SO ₂ CH ₃	2-F-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
467	SO ₂ CH ₃	2-F-phenyl	4-morpholinocarbonyl
468	SO ₂ CH ₃	2-F-phenyl	2-methyl-1-imidazolyl
469	SO ₂ CH ₃	2-F-phenyl	5-methyl-1-imidazolyl
470	SO ₂ CH ₃	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
471	SO ₂ CH ₃	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
472	SO ₂ CH ₃	2,6-diF-phenyl	2-(methylaminosulfonyl)phenyl

473	SO ₂ CH ₃	2,6-diF-phenyl	1-pyrrolidinocarbonyl
474	SO ₂ CH ₃	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
475	SO ₂ CH ₃	2,6-diF-phenyl	4-morpholino
476	SO ₂ CH ₃	2,6-diF-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
477	SO ₂ CH ₃	2,6-diF-phenyl	4-morpholinocarbonyl
478	SO ₂ CH ₃	2,6-diF-phenyl	2-methyl-1-imidazolyl
479	SO ₂ CH ₃	2,6-diF-phenyl	5-methyl-1-imidazolyl
480	SO ₂ CH ₃	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
481	CH ₂ NH-SO ₂ CH ₃	phenyl	2-(aminosulfonyl)phenyl
482	CH ₂ NH-SO ₂ CH ₃	phenyl	2-(methylaminosulfonyl)phenyl
483	CH ₂ NH-SO ₂ CH ₃	phenyl	1-pyrrolidinocarbonyl
484	CH ₂ NH-SO ₂ CH ₃	phenyl	2-(methylsulfonyl)phenyl
485	CH ₂ NH-SO ₂ CH ₃	phenyl	4-morpholino
486	CH ₂ NH-SO ₂ CH ₃	phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
487	CH ₂ NH-SO ₂ CH ₃	phenyl	4-morpholinocarbonyl
488	CH ₂ NH-SO ₂ CH ₃	phenyl	2-methyl-1-imidazolyl
489	CH ₂ NH-SO ₂ CH ₃	phenyl	5-methyl-1-imidazolyl
490	CH ₂ NH-SO ₂ CH ₃	phenyl	2-methylsulfonyl-1-imidazolyl
491	CH ₂ NH-SO ₂ CH ₃	2-pyridyl	2-(aminosulfonyl)phenyl
492	CH ₂ NH-SO ₂ CH ₃	2-pyridyl	2-(methylaminosulfonyl)phenyl
493	CH ₂ NH-SO ₂ CH ₃	2-pyridyl	1-pyrrolidinocarbonyl
494	CH ₂ NH-SO ₂ CH ₃	2-pyridyl	2-(methylsulfonyl)phenyl
495	CH ₂ NH-SO ₂ CH ₃	2-pyridyl	4-morpholino
496	CH ₂ NH-SO ₂ CH ₃	2-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
497	CH ₂ NH-SO ₂ CH ₃	2-pyridyl	4-morpholinocarbonyl
498	CH ₂ NH-SO ₂ CH ₃	2-pyridyl	2-methyl-1-imidazolyl
499	CH ₂ NH-SO ₂ CH ₃	2-pyridyl	5-methyl-1-imidazolyl
500	CH ₂ NH-SO ₂ CH ₃	2-pyridyl	2-methylsulfonyl-1-imidazolyl

501	CH ₂ NH- SO ₂ CH ₃	3-pyridyl	2-(aminosulfonyl)phenyl
502	CH ₂ NH- SO ₂ CH ₃	3-pyridyl	2-(methylaninosulfonyl)phenyl
503	CH ₂ NH- SO ₂ CH ₃	3-pyridyl	1-pyrrolidinocarbonyl
504	CH ₂ NH- SO ₂ CH ₃	3-pyridyl	2-(methylsulfonyl)phenyl
505	CH ₂ NH- SO ₂ CH ₃	3-pyridyl	4-morpholino
506	CH ₂ NH- SO ₂ CH ₃	3-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
507	CH ₂ NH- SO ₂ CH ₃	3-pyridyl	4-morpholinocarbonyl
508	CH ₂ NH- SO ₂ CH ₃	3-pyridyl	2-methyl-1-imidazolyl
509	CH ₂ NH- SO ₂ CH ₃	3-pyridyl	5-methyl-1-imidazolyl
510	CH ₂ NH- SO ₂ CH ₃	3-pyridyl	2-methylsulfonyl-1-imidazolyl
511	CH ₂ NH- SO ₂ CH ₃	2-pyrimidyl	2-(aminosulfonyl)phenyl
512	CH ₂ NH- SO ₂ CH ₃	2-pyrimidyl	2-(methylaninosulfonyl)phenyl
513	CH ₂ NH- SO ₂ CH ₃	2-pyrimidyl	1-pyrrolidinocarbonyl
514	CH ₂ NH- SO ₂ CH ₃	2-pyrimidyl	2-(methylsulfonyl)phenyl
515	CH ₂ NH- SO ₂ CH ₃	2-pyrimidyl	4-morpholino
516	CH ₂ NH- SO ₂ CH ₃	2-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
517	CH ₂ NH- SO ₂ CH ₃	2-pyrimidyl	4-morpholinocarbonyl
518	CH ₂ NH- SO ₂ CH ₃	2-pyrimidyl	2-methyl-1-imidazolyl
519	CH ₂ NH- SO ₂ CH ₃	2-pyrimidyl	5-methyl-1-imidazolyl
520	CH ₂ NH- SO ₂ CH ₃	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
521	CH ₂ NH- SO ₂ CH ₃	5-pyrimidyl	2-(aminosulfonyl)phenyl
522	CH ₂ NH- SO ₂ CH ₃	5-pyrimidyl	2-(methylaninosulfonyl)phenyl
523	CH ₂ NH- SO ₂ CH ₃	5-pyrimidyl	1-pyrrolidinocarbonyl
524	CH ₂ NH- SO ₂ CH ₃	5-pyrimidyl	2-(methylsulfonyl)phenyl

525	CH ₂ NH-SO ₂ CH ₃	5-pyrimidyl	4-morpholino
526	CH ₂ NH-SO ₂ CH ₃	5-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
527	CH ₂ NH-SO ₂ CH ₃	5-pyrimidyl	4-morpholinocarbonyl
528	CH ₂ NH-SO ₂ CH ₃	5-pyrimidyl	2-methyl-1-imidazolyl
529	CH ₂ NH-SO ₂ CH ₃	5-pyrimidyl	5-methyl-1-imidazolyl
530	CH ₂ NH-SO ₂ CH ₃	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
531	CH ₂ NH-SO ₂ CH ₃	2-Cl-phenyl	2-(aminosulfonyl)phenyl
532	CH ₂ NH-SO ₂ CH ₃	2-Cl-phenyl	2-(methylaninosulfonyl)phenyl
533	CH ₂ NH-SO ₂ CH ₃	2-Cl-phenyl	1-pyrrolidinocarbonyl
534	CH ₂ NH-SO ₂ CH ₃	2-Cl-phenyl	2-(methylsulfonyl)phenyl
535	CH ₂ NH-SO ₂ CH ₃	2-Cl-phenyl	4-morpholino
536	CH ₂ NH-SO ₂ CH ₃	2-Cl-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
537	CH ₂ NH-SO ₂ CH ₃	2-Cl-phenyl	4-morpholinocarbonyl
538	CH ₂ NH-SO ₂ CH ₃	2-Cl-phenyl	2-methyl-1-imidazolyl
539	CH ₂ NH-SO ₂ CH ₃	2-Cl-phenyl	5-methyl-1-imidazolyl
540	CH ₂ NH-SO ₂ CH ₃	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
541	CH ₂ NH-SO ₂ CH ₃	2-F-phenyl	2-(aminosulfonyl)phenyl
542	CH ₂ NH-SO ₂ CH ₃	2-F-phenyl	2-(methylaninosulfonyl)phenyl
543	CH ₂ NH-SO ₂ CH ₃	2-F-phenyl	1-pyrrolidinocarbonyl
544	CH ₂ NH-SO ₂ CH ₃	2-F-phenyl	2-(methylsulfonyl)phenyl
545	CH ₂ NH-SO ₂ CH ₃	2-F-phenyl	4-morpholino
546	CH ₂ NH-SO ₂ CH ₃	2-F-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
547	CH ₂ NH-SO ₂ CH ₃	2-F-phenyl	4-morpholinocarbonyl
548	CH ₂ NH-SO ₂ CH ₃	2-F-phenyl	2-methyl-1-imidazolyl

549	CH ₂ NH-SO ₂ CH ₃	2-F-phenyl	5-methyl-1-imidazolyl
550	CH ₂ NH-SO ₂ CH ₃	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
551	CH ₂ NH-SO ₂ CH ₃	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
552	CH ₂ NH-SO ₂ CH ₃	2,6-diF-phenyl	2-(methylaminosulfonyl)phenyl
553	CH ₂ NH-SO ₂ CH ₃	2,6-diF-phenyl	1-pyrrolidinocarbonyl
554	CH ₂ NH-SO ₂ CH ₃	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
555	CH ₂ NH-SO ₂ CH ₃	2,6-diF-phenyl	4-morpholino
556	CH ₂ NH-SO ₂ CH ₃	2,6-diF-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
557	CH ₂ NH-SO ₂ CH ₃	2,6-diF-phenyl	4-morpholinocarbonyl
558	CH ₂ NH-SO ₂ CH ₃	2,6-diF-phenyl	2-methyl-1-imidazolyl
559	CH ₂ NH-SO ₂ CH ₃	2,6-diF-phenyl	5-methyl-1-imidazolyl
560	CH ₂ NH-SO ₂ CH ₃	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
561	Cl	phenyl	2-(aminosulfonyl)phenyl
562	Cl	phenyl	2-(methylaminosulfonyl)phenyl
563	Cl	phenyl	1-pyrrolidinocarbonyl
564	Cl	phenyl	2-(methylsulfonyl)phenyl
565	Cl	phenyl	4-morpholino
566	Cl	phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
567	Cl	phenyl	4-morpholinocarbonyl
568	Cl	phenyl	2-methyl-1-imidazolyl
569	Cl	phenyl	5-methyl-1-imidazolyl
570	Cl	phenyl	2-methylsulfonyl-1-imidazolyl
571	Cl	2-pyridyl	2-(aminosulfonyl)phenyl
572	Cl	2-pyridyl	2-(methylaminosulfonyl)phenyl
573	Cl	2-pyridyl	1-pyrrolidinocarbonyl
574	Cl	2-pyridyl	2-(methylsulfonyl)phenyl
575	Cl	2-pyridyl	4-morpholino
576	Cl	2-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
577	Cl	2-pyridyl	4-morpholinocarbonyl
578	Cl	2-pyridyl	2-methyl-1-imidazolyl
579	Cl	2-pyridyl	5-methyl-1-imidazolyl
580	Cl	2-pyridyl	2-methylsulfonyl-1-imidazolyl
581	Cl	3-pyridyl	2-(aminosulfonyl)phenyl
582	Cl	3-pyridyl	2-(methylaminosulfonyl)phenyl
583	Cl	3-pyridyl	1-pyrrolidinocarbonyl
584	Cl	3-pyridyl	2-(methylsulfonyl)phenyl
585	Cl	3-pyridyl	4-morpholino
586	Cl	3-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
587	Cl	3-pyridyl	4-morpholinocarbonyl
588	Cl	3-pyridyl	2-methyl-1-imidazolyl

589	Cl	3-pyridyl	5-methyl-1-imidazolyl
590	Cl	3-pyridyl	2-methylsulfonyl-1-imidazolyl
591	Cl	2-pyrimidyl	2-(aminosulfonyl)phenyl
592	Cl	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
593	Cl	2-pyrimidyl	1-pyrrolidinocarbonyl
594	Cl	2-pyrimidyl	2-(methylsulfonyl)phenyl
595	Cl	2-pyrimidyl	4-morpholino
596	Cl	2-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
597	Cl	2-pyrimidyl	4-morpholinocarbonyl
598	Cl	2-pyrimidyl	2-methyl-1-imidazolyl
599	Cl	2-pyrimidyl	5-methyl-1-imidazolyl
600	Cl	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
601	Cl	5-pyrimidyl	2-(aminosulfonyl)phenyl
602	Cl	5-pyrimidyl	2-(methylaminosulfonyl)phenyl
603	Cl	5-pyrimidyl	1-pyrrolidinocarbonyl
604	Cl	5-pyrimidyl	2-(methylsulfonyl)phenyl
605	Cl	5-pyrimidyl	4-morpholino
606	Cl	5-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
607	Cl	5-pyrimidyl	4-morpholinocarbonyl
608	Cl	5-pyrimidyl	2-methyl-1-imidazolyl
609	Cl	5-pyrimidyl	5-methyl-1-imidazolyl
610	Cl	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
611	Cl	2-Cl-phenyl	2-(aminosulfonyl)phenyl
612	Cl	2-Cl-phenyl	2-(methylaminosulfonyl)phenyl
613	Cl	2-Cl-phenyl	1-pyrrolidinocarbonyl
614	Cl	2-Cl-phenyl	2-(methylsulfonyl)phenyl
615	Cl	2-Cl-phenyl	4-morpholino
616	Cl	2-Cl-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
617	Cl	2-Cl-phenyl	4-morpholinocarbonyl
618	Cl	2-Cl-phenyl	2-methyl-1-imidazolyl
619	Cl	2-Cl-phenyl	5-methyl-1-imidazolyl
620	Cl	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
621	Cl	2-F-phenyl	2-(aminosulfonyl)phenyl
622	Cl	2-F-phenyl	2-(methylaminosulfonyl)phenyl
623	Cl	2-F-phenyl	1-pyrrolidinocarbonyl
624	Cl	2-F-phenyl	2-(methylsulfonyl)phenyl
625	Cl	2-F-phenyl	4-morpholino
626	Cl	2-F-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
627	Cl	2-F-phenyl	4-morpholinocarbonyl
628	Cl	2-F-phenyl	2-methyl-1-imidazolyl
629	Cl	2-F-phenyl	5-methyl-1-imidazolyl
630	Cl	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
631	Cl	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
632	Cl	2,6-diF-phenyl	2-(methylaminosulfonyl)phenyl
633	Cl	2,6-diF-phenyl	1-pyrrolidinocarbonyl
634	Cl	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
635	Cl	2,6-diF-phenyl	4-morpholino
636	Cl	2,6-diF-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
637	Cl	2,6-diF-phenyl	4-morpholinocarbonyl
638	Cl	2,6-diF-phenyl	2-methyl-1-imidazolyl
639	Cl	2,6-diF-phenyl	5-methyl-1-imidazolyl
640	Cl	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
641	F	phenyl	2-(aminosulfonyl)phenyl
642	F	phenyl	2-(methylaminosulfonyl)phenyl
643	F	phenyl	1-pyrrolidinocarbonyl

644	F	phenyl	2-(methylsulfonyl)phenyl
645	F	phenyl	4-morpholino
646	F	phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
647	F	phenyl	4-morpholinocarbonyl
648	F	phenyl	2-methyl-1-imidazolyl
649	F	phenyl	5-methyl-1-imidazolyl
650	F	phenyl	2-methylsulfonyl-1-imidazolyl
651	F	2-pyridyl	2-(aminosulfonyl)phenyl
652	F	2-pyridyl	2-(methylaminosulfonyl)phenyl
653	F	2-pyridyl	1-pyrrolidinocarbonyl
654	F	2-pyridyl	2-(methylsulfonyl)phenyl
655	F	2-pyridyl	4-morpholino
656	F	2-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
657	F	2-pyridyl	4-morpholinocarbonyl
658	F	2-pyridyl	2-methyl-1-imidazolyl
659	F	2-pyridyl	5-methyl-1-imidazolyl
660	F	2-pyridyl	2-methylsulfonyl-1-imidazolyl
661	F	3-pyridyl	2-(aminosulfonyl)phenyl
662	F	3-pyridyl	2-(methylaminosulfonyl)phenyl
663	F	3-pyridyl	1-pyrrolidinocarbonyl
664	F	3-pyridyl	2-(methylsulfonyl)phenyl
665	F	3-pyridyl	4-morpholino
666	F	3-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
667	F	3-pyridyl	4-morpholinocarbonyl
668	F	3-pyridyl	2-methyl-1-imidazolyl
669	F	3-pyridyl	5-methyl-1-imidazolyl
670	F	3-pyridyl	2-methylsulfonyl-1-imidazolyl
671	F	2-pyrimidyl	2-(aminosulfonyl)phenyl
672	F	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
673	F	2-pyrimidyl	1-pyrrolidinocarbonyl
674	F	2-pyrimidyl	2-(methylsulfonyl)phenyl
675	F	2-pyrimidyl	4-morpholino
676	F	2-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
677	F	2-pyrimidyl	4-morpholinocarbonyl
678	F	2-pyrimidyl	2-methyl-1-imidazolyl
679	F	2-pyrimidyl	5-methyl-1-imidazolyl
680	F	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
681	F	5-pyrimidyl	2-(aminosulfonyl)phenyl
682	F	5-pyrimidyl	2-(methylaminosulfonyl)phenyl
683	F	5-pyrimidyl	1-pyrrolidinocarbonyl
684	F	5-pyrimidyl	2-(methylsulfonyl)phenyl
685	F	5-pyrimidyl	4-morpholino
686	F	5-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
687	F	5-pyrimidyl	4-morpholinocarbonyl
688	F	5-pyrimidyl	2-methyl-1-imidazolyl
689	F	5-pyrimidyl	5-methyl-1-imidazolyl
690	F	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
691	F	2-Cl-phenyl	2-(aminosulfonyl)phenyl
692	F	2-Cl-phenyl	2-(methylaminosulfonyl)phenyl
693	F	2-Cl-phenyl	1-pyrrolidinocarbonyl
694	F	2-Cl-phenyl	2-(methylsulfonyl)phenyl
695	F	2-Cl-phenyl	4-morpholino
696	F	2-Cl-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
697	F	2-Cl-phenyl	4-morpholinocarbonyl
698	F	2-Cl-phenyl	2-methyl-1-imidazolyl

699	F	2-Cl-phenyl	5-methyl-1-imidazolyl
700	F	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
701	F	2-F-phenyl	2-(aminosulfonyl)phenyl
702	F	2-F-phenyl	2-(methylaminosulfonyl)phenyl
703	F	2-F-phenyl	1-pyrrolidinocarbonyl
704	F	2-F-phenyl	2-(methylsulfonyl)phenyl
705	F	2-F-phenyl	4-morpholino
706	F	2-F-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
707	F	2-F-phenyl	4-morpholinocarbonyl
708	F	2-F-phenyl	2-methyl-1-imidazolyl
709	F	2-F-phenyl	5-methyl-1-imidazolyl
710	F	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
711	F	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
712	F	2,6-diF-phenyl	2-(methylaminosulfonyl)phenyl
713	F	2,6-diF-phenyl	1-pyrrolidinocarbonyl
714	F	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
715	F	2,6-diF-phenyl	4-morpholino
716	F	2,6-diF-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
717	F	2,6-diF-phenyl	4-morpholinocarbonyl
718	F	2,6-diF-phenyl	2-methyl-1-imidazolyl
719	F	2,6-diF-phenyl	5-methyl-1-imidazolyl
720	F	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
721	CO ₂ CH ₃	phenyl	2-(aminosulfonyl)phenyl
722	CO ₂ CH ₃	phenyl	2-(methylaminosulfonyl)phenyl
723	CO ₂ CH ₃	phenyl	1-pyrrolidinocarbonyl
724	CO ₂ CH ₃	phenyl	2-(methylsulfonyl)phenyl
725	CO ₂ CH ₃	phenyl	4-morpholino
726	CO ₂ CH ₃	phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
727	CO ₂ CH ₃	phenyl	4-morpholinocarbonyl
728	CO ₂ CH ₃	phenyl	2-methyl-1-imidazolyl
729	CO ₂ CH ₃	phenyl	5-methyl-1-imidazolyl
730	CO ₂ CH ₃	phenyl	2-methylsulfonyl-1-imidazolyl
731	CO ₂ CH ₃	2-pyridyl	2-(aminosulfonyl)phenyl
732	CO ₂ CH ₃	2-pyridyl	2-(methylaminosulfonyl)phenyl
733	CO ₂ CH ₃	2-pyridyl	1-pyrrolidinocarbonyl
734	CO ₂ CH ₃	2-pyridyl	2-(methylsulfonyl)phenyl
735	CO ₂ CH ₃	2-pyridyl	4-morpholino
736	CO ₂ CH ₃	2-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
737	CO ₂ CH ₃	2-pyridyl	4-morpholinocarbonyl
738	CO ₂ CH ₃	2-pyridyl	2-methyl-1-imidazolyl
739	CO ₂ CH ₃	2-pyridyl	5-methyl-1-imidazolyl
740	CO ₂ CH ₃	2-pyridyl	2-methylsulfonyl-1-imidazolyl
741	CO ₂ CH ₃	3-pyridyl	2-(aminosulfonyl)phenyl
742	CO ₂ CH ₃	3-pyridyl	2-(methylaminosulfonyl)phenyl
743	CO ₂ CH ₃	3-pyridyl	1-pyrrolidinocarbonyl
744	CO ₂ CH ₃	3-pyridyl	2-(methylsulfonyl)phenyl
745	CO ₂ CH ₃	3-pyridyl	4-morpholino
746	CO ₂ CH ₃	3-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
747	CO ₂ CH ₃	3-pyridyl	4-morpholinocarbonyl
748	CO ₂ CH ₃	3-pyridyl	2-methyl-1-imidazolyl
749	CO ₂ CH ₃	3-pyridyl	5-methyl-1-imidazolyl

750	CO ₂ CH ₃	3-pyridyl	2-methylsulfonyl-1-imidazolyl
751	CO ₂ CH ₃	2-pyrimidyl	2-(aminosulfonyl)phenyl
752	CO ₂ CH ₃	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
753	CO ₂ CH ₃	2-pyrimidyl	1-pyrrolidinocarbonyl
754	CO ₂ CH ₃	2-pyrimidyl	2-(methylsulfonyl)phenyl
755	CO ₂ CH ₃	2-pyrimidyl	4-morpholino
756	CO ₂ CH ₃	2-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
757	CO ₂ CH ₃	2-pyrimidyl	4-morpholinocarbonyl
758	CO ₂ CH ₃	2-pyrimidyl	2-methyl-1-imidazolyl
759	CO ₂ CH ₃	2-pyrimidyl	5-methyl-1-imidazolyl
760	CO ₂ CH ₃	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
761	CO ₂ CH ₃	5-pyrimidyl	2-(aminosulfonyl)phenyl
762	CO ₂ CH ₃	5-pyrimidyl	2-(methylaminosulfonyl)phenyl
763	CO ₂ CH ₃	5-pyrimidyl	1-pyrrolidinocarbonyl
764	CO ₂ CH ₃	5-pyrimidyl	2-(methylsulfonyl)phenyl
765	CO ₂ CH ₃	5-pyrimidyl	4-morpholino
766	CO ₂ CH ₃	5-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
767	CO ₂ CH ₃	5-pyrimidyl	4-morpholinocarbonyl
768	CO ₂ CH ₃	5-pyrimidyl	2-methyl-1-imidazolyl
769	CO ₂ CH ₃	5-pyrimidyl	5-methyl-1-imidazolyl
770	CO ₂ CH ₃	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
771	CO ₂ CH ₃	2-Cl-phenyl	2-(aminosulfonyl)phenyl
772	CO ₂ CH ₃	2-Cl-phenyl	2-(methylaminosulfonyl)phenyl
773	CO ₂ CH ₃	2-Cl-phenyl	1-pyrrolidinocarbonyl
774	CO ₂ CH ₃	2-Cl-phenyl	2-(methylsulfonyl)phenyl
775	CO ₂ CH ₃	2-Cl-phenyl	4-morpholino
776	CO ₂ CH ₃	2-Cl-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
777	CO ₂ CH ₃	2-Cl-phenyl	4-morpholinocarbonyl
778	CO ₂ CH ₃	2-Cl-phenyl	2-methyl-1-imidazolyl
779	CO ₂ CH ₃	2-Cl-phenyl	5-methyl-1-imidazolyl
780	CO ₂ CH ₃	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
781	CO ₂ CH ₃	2-F-phenyl	2-(aminosulfonyl)phenyl
782	CO ₂ CH ₃	2-F-phenyl	2-(methylaminosulfonyl)phenyl
783	CO ₂ CH ₃	2-F-phenyl	1-pyrrolidinocarbonyl
784	CO ₂ CH ₃	2-F-phenyl	2-(methylsulfonyl)phenyl
785	CO ₂ CH ₃	2-F-phenyl	4-morpholino
786	CO ₂ CH ₃	2-F-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
787	CO ₂ CH ₃	2-F-phenyl	4-morpholinocarbonyl
788	CO ₂ CH ₃	2-F-phenyl	2-methyl-1-imidazolyl
789	CO ₂ CH ₃	2-F-phenyl	5-methyl-1-imidazolyl
790	CO ₂ CH ₃	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
791	CO ₂ CH ₃	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
792	CO ₂ CH ₃	2,6-diF-phenyl	2-(methylaminosulfonyl)phenyl
793	CO ₂ CH ₃	2,6-diF-phenyl	1-pyrrolidinocarbonyl
794	CO ₂ CH ₃	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
795	CO ₂ CH ₃	2,6-diF-phenyl	4-morpholino
796	CO ₂ CH ₃	2,6-diF-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
797	CO ₂ CH ₃	2,6-diF-phenyl	4-morpholinocarbonyl


798	CO ₂ CH ₃	2,6-diF-phenyl	2-methyl-1-imidazolyl
799	CO ₂ CH ₃	2,6-diF-phenyl	5-methyl-1-imidazolyl
800	CO ₂ CH ₃	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
801	CH ₂ OCH ₃	phenyl	2-(aminosulfonyl)phenyl
802	CH ₂ OCH ₃	phenyl	2-(methylaminosulfonyl)phenyl
803	CH ₂ OCH ₃	phenyl	1-pyrrolidinocarbonyl
804	CH ₂ OCH ₃	phenyl	2-(methylsulfonyl)phenyl
805	CH ₂ OCH ₃	phenyl	4-morpholino
806	CH ₂ OCH ₃	phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
807	CH ₂ OCH ₃	phenyl	4-morpholinocarbonyl
808	CH ₂ OCH ₃	phenyl	2-methyl-1-imidazolyl
809	CH ₂ OCH ₃	phenyl	5-methyl-1-imidazolyl
810	CH ₂ OCH ₃	phenyl	2-methylsulfonyl-1-imidazolyl
811	CH ₂ OCH ₃	2-pyridyl	2-(aminosulfonyl)phenyl
812	CH ₂ OCH ₃	2-pyridyl	2-(methylaminosulfonyl)phenyl
813	CH ₂ OCH ₃	2-pyridyl	1-pyrrolidinocarbonyl
814	CH ₂ OCH ₃	2-pyridyl	2-(methylsulfonyl)phenyl
815	CH ₂ OCH ₃	2-pyridyl	4-morpholino
816	CH ₂ OCH ₃	2-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
817	CH ₂ OCH ₃	2-pyridyl	4-morpholinocarbonyl
818	CH ₂ OCH ₃	2-pyridyl	2-methyl-1-imidazolyl
819	CH ₂ OCH ₃	2-pyridyl	5-methyl-1-imidazolyl
820	CH ₂ OCH ₃	2-pyridyl	2-methylsulfonyl-1-imidazolyl
821	CH ₂ OCH ₃	3-pyridyl	2-(aminosulfonyl)phenyl
822	CH ₂ OCH ₃	3-pyridyl	2-(methylaminosulfonyl)phenyl
823	CH ₂ OCH ₃	3-pyridyl	1-pyrrolidinocarbonyl
824	CH ₂ OCH ₃	3-pyridyl	2-(methylsulfonyl)phenyl
825	CH ₂ OCH ₃	3-pyridyl	4-morpholino
826	CH ₂ OCH ₃	3-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
827	CH ₂ OCH ₃	3-pyridyl	4-morpholinocarbonyl
828	CH ₂ OCH ₃	3-pyridyl	2-methyl-1-imidazolyl
829	CH ₂ OCH ₃	3-pyridyl	5-methyl-1-imidazolyl
830	CH ₂ OCH ₃	3-pyridyl	2-methylsulfonyl-1-imidazolyl
831	CH ₂ OCH ₃	2-pyrimidyl	2-(aminosulfonyl)phenyl
832	CH ₂ OCH ₃	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
833	CH ₂ OCH ₃	2-pyrimidyl	1-pyrrolidinocarbonyl
834	CH ₂ OCH ₃	2-pyrimidyl	2-(methylsulfonyl)phenyl
835	CH ₂ OCH ₃	2-pyrimidyl	4-morpholino
836	CH ₂ OCH ₃	2-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
837	CH ₂ OCH ₃	2-pyrimidyl	4-morpholinocarbonyl
838	CH ₂ OCH ₃	2-pyrimidyl	2-methyl-1-imidazolyl
839	CH ₂ OCH ₃	2-pyrimidyl	5-methyl-1-imidazolyl
840	CH ₂ OCH ₃	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
841	CH ₂ OCH ₃	5-pyrimidyl	2-(aminosulfonyl)phenyl
842	CH ₂ OCH ₃	5-pyrimidyl	2-(methylaminosulfonyl)phenyl
843	CH ₂ OCH ₃	5-pyrimidyl	1-pyrrolidinocarbonyl
844	CH ₂ OCH ₃	5-pyrimidyl	2-(methylsulfonyl)phenyl
845	CH ₂ OCH ₃	5-pyrimidyl	4-morpholino

846	CH ₂ OCH ₃	5-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
847	CH ₂ OCH ₃	5-pyrimidyl	4-morpholinocarbonyl
848	CH ₂ OCH ₃	5-pyrimidyl	2-methyl-1-imidazolyl
849	CH ₂ OCH ₃	5-pyrimidyl	5-methyl-1-imidazolyl
850	CH ₂ OCH ₃	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
851	CH ₂ OCH ₃	2-Cl-phenyl	2-(aminosulfonyl)phenyl
852	CH ₂ OCH ₃	2-Cl-phenyl	2-(methylaminosulfonyl)phenyl
853	CH ₂ OCH ₃	2-Cl-phenyl	1-pyrrolidinocarbonyl
854	CH ₂ OCH ₃	2-Cl-phenyl	2-(methylsulfonyl)phenyl
855	CH ₂ OCH ₃	2-Cl-phenyl	4-morpholino
856	CH ₂ OCH ₃	2-Cl-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
857	CH ₂ OCH ₃	2-Cl-phenyl	4-morpholinocarbonyl
858	CH ₂ OCH ₃	2-Cl-phenyl	2-methyl-1-imidazolyl
859	CH ₂ OCH ₃	2-Cl-phenyl	5-methyl-1-imidazolyl
860	CH ₂ OCH ₃	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
861	CH ₂ OCH ₃	2-F-phenyl	2-(aminosulfonyl)phenyl
862	CH ₂ OCH ₃	2-F-phenyl	2-(methylaminosulfonyl)phenyl
863	CH ₂ OCH ₃	2-F-phenyl	1-pyrrolidinocarbonyl
864	CH ₂ OCH ₃	2-F-phenyl	2-(methylsulfonyl)phenyl
865	CH ₂ OCH ₃	2-F-phenyl	4-morpholino
866	CH ₂ OCH ₃	2-F-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
867	CH ₂ OCH ₃	2-F-phenyl	4-morpholinocarbonyl
868	CH ₂ OCH ₃	2-F-phenyl	2-methyl-1-imidazolyl
869	CH ₂ OCH ₃	2-F-phenyl	5-methyl-1-imidazolyl
870	CH ₂ OCH ₃	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
871	CH ₂ OCH ₃	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
872	CH ₂ OCH ₃	2,6-diF-phenyl	2-(methylaminosulfonyl)phenyl
873	CH ₂ OCH ₃	2,6-diF-phenyl	1-pyrrolidinocarbonyl
874	CH ₂ OCH ₃	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
875	CH ₂ OCH ₃	2,6-diF-phenyl	4-morpholino
876	CH ₂ OCH ₃	2,6-diF-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
877	CH ₂ OCH ₃	2,6-diF-phenyl	4-morpholinocarbonyl
878	CH ₂ OCH ₃	2,6-diF-phenyl	2-methyl-1-imidazolyl
879	CH ₂ OCH ₃	2,6-diF-phenyl	5-methyl-1-imidazolyl
880	CH ₂ OCH ₃	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
881	CONH ₂	phenyl	2-(aminosulfonyl)phenyl
882	CONH ₂	phenyl	2-(methylaminosulfonyl)phenyl
883	CONH ₂	phenyl	1-pyrrolidinocarbonyl
884	CONH ₂	phenyl	2-(methylsulfonyl)phenyl
885	CONH ₂	phenyl	4-morpholino
886	CONH ₂	phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
887	CONH ₂	phenyl	4-morpholinocarbonyl
888	CONH ₂	phenyl	2-methyl-1-imidazolyl
889	CONH ₂	phenyl	5-methyl-1-imidazolyl
890	CONH ₂	phenyl	2-methylsulfonyl-1-imidazolyl
891	CONH ₂	2-pyridyl	2-(aminosulfonyl)phenyl
892	CONH ₂	2-pyridyl	2-(methylaminosulfonyl)phenyl
893	CONH ₂	2-pyridyl	1-pyrrolidinocarbonyl

894	CONH ₂	2-pyridyl	2-(methylsulfonyl)phenyl
895	CONH ₂	2-pyridyl	4-morpholino
896	CONH ₂	2-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
897	CONH ₂	2-pyridyl	4-morpholinocarbonyl
898	CONH ₂	2-pyridyl	2-methyl-1-imidazolyl
899	CONH ₂	2-pyridyl	5-methyl-1-imidazolyl
900	CONH ₂	2-pyridyl	2-methylsulfonyl-1-imidazolyl
901	CONH ₂	3-pyridyl	2-(aminosulfonyl)phenyl
902	CONH ₂	3-pyridyl	2-(methylaminosulfonyl)phenyl
903	CONH ₂	3-pyridyl	1-pyrrolidinocarbonyl
904	CONH ₂	3-pyridyl	2-(methylsulfonyl)phenyl
905	CONH ₂	3-pyridyl	4-morpholino
906	CONH ₂	3-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
907	CONH ₂	3-pyridyl	4-morpholinocarbonyl
908	CONH ₂	3-pyridyl	2-methyl-1-imidazolyl
909	CONH ₂	3-pyridyl	5-methyl-1-imidazolyl
910	CONH ₂	3-pyridyl	2-methylsulfonyl-1-imidazolyl
911	CONH ₂	2-pyrimidyl	2-(aminosulfonyl)phenyl
912	CONH ₂	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
913	CONH ₂	2-pyrimidyl	1-pyrrolidinocarbonyl
914	CONH ₂	2-pyrimidyl	2-(methylsulfonyl)phenyl
915	CONH ₂	2-pyrimidyl	4-morpholino
916	CONH ₂	2-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
917	CONH ₂	2-pyrimidyl	4-morpholinocarbonyl
918	CONH ₂	2-pyrimidyl	2-methyl-1-imidazolyl
919	CONH ₂	2-pyrimidyl	5-methyl-1-imidazolyl
920	CONH ₂	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
921	CONH ₂	5-pyrimidyl	2-(aminosulfonyl)phenyl
922	CONH ₂	5-pyrimidyl	2-(methylaminosulfonyl)phenyl
923	CONH ₂	5-pyrimidyl	1-pyrrolidinocarbonyl
924	CONH ₂	5-pyrimidyl	2-(methylsulfonyl)phenyl
925	CONH ₂	5-pyrimidyl	4-morpholino
926	CONH ₂	5-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
927	CONH ₂	5-pyrimidyl	4-morpholinocarbonyl
928	CONH ₂	5-pyrimidyl	2-methyl-1-imidazolyl
929	CONH ₂	5-pyrimidyl	5-methyl-1-imidazolyl
930	CONH ₂	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
931	CONH ₂	2-Cl-phenyl	2-(aminosulfonyl)phenyl
932	CONH ₂	2-Cl-phenyl	2-(methylaminosulfonyl)phenyl
933	CONH ₂	2-Cl-phenyl	1-pyrrolidinocarbonyl
934	CONH ₂	2-Cl-phenyl	2-(methylsulfonyl)phenyl
935	CONH ₂	2-Cl-phenyl	4-morpholino
936	CONH ₂	2-Cl-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
937	CONH ₂	2-Cl-phenyl	4-morpholinocarbonyl
938	CONH ₂	2-Cl-phenyl	2-methyl-1-imidazolyl
939	CONH ₂	2-Cl-phenyl	5-methyl-1-imidazolyl
940	CONH ₂	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
941	CONH ₂	2-F-phenyl	2-(aminosulfonyl)phenyl

942	CONH ₂	2-F-phenyl	2-(methylaminosulfonyl)phenyl
943	CONH ₂	2-F-phenyl	1-pyrrolidinocarbonyl
944	CONH ₂	2-F-phenyl	2-(methylsulfonyl)phenyl
945	CONH ₂	2-F-phenyl	4-morpholino
946	CONH ₂	2-F-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
947	CONH ₂	2-F-phenyl	4-morpholinocarbonyl
948	CONH ₂	2-F-phenyl	2-methyl-1-imidazolyl
949	CONH ₂	2-F-phenyl	5-methyl-1-imidazolyl
950	CONH ₂	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
951	CONH ₂	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
952	CONH ₂	2,6-diF-phenyl	2-(methylaminosulfonyl)phenyl
953	CONH ₂	2,6-diF-phenyl	1-pyrrolidinocarbonyl
954	CONH ₂	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
955	CONH ₂	2,6-diF-phenyl	4-morpholino
956	CONH ₂	2,6-diF-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
957	CONH ₂	2,6-diF-phenyl	4-morpholinocarbonyl
958	CONH ₂	2,6-diF-phenyl	2-methyl-1-imidazolyl
959	CONH ₂	2,6-diF-phenyl	5-methyl-1-imidazolyl
960	CONH ₂	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl

Table 5




For each example, DE is:


- (A) pyridin-4-yl-CH₂,
- (B) 2-amino-pyrimidin-4-yl,
- (C) 6-amino-pyridin-2-yl,
- (D) 3-amidino-4-F-phenyl, or
- (E) N-amidino-3-piperidinyl.


Ex #	A	B
1	phenyl	2-(aminosulfonyl)phenyl
2	phenyl	2-(methylaminosulfonyl)phenyl
3	phenyl	1-pyrrolidinocarbonyl
4	phenyl	2-(methylsulfonyl)phenyl
5	phenyl	4-morpholino
6	phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
7	phenyl	4-morpholinocarbonyl
8	phenyl	2-methyl-1-imidazolyl
9	phenyl	5-methyl-1-imidazolyl
10	phenyl	2-methylsulfonyl-1-imidazolyl
11	2-pyridyl	2-(aminosulfonyl)phenyl
12	2-pyridyl	2-(methylaminosulfonyl)phenyl
13	2-pyridyl	1-pyrrolidinocarbonyl
14	2-pyridyl	2-(methylsulfonyl)phenyl
15	2-pyridyl	4-morpholino
16	2-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
17	2-pyridyl	4-morpholinocarbonyl
18	2-pyridyl	2-methyl-1-imidazolyl
19	2-pyridyl	5-methyl-1-imidazolyl
20	2-pyridyl	2-methylsulfonyl-1-imidazolyl
21	3-pyridyl	2-(aminosulfonyl)phenyl
22	3-pyridyl	2-(methylaminosulfonyl)phenyl
23	3-pyridyl	1-pyrrolidinocarbonyl
24	3-pyridyl	2-(methylsulfonyl)phenyl
25	3-pyridyl	4-morpholino
26	3-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
27	3-pyridyl	4-morpholinocarbonyl
28	3-pyridyl	2-methyl-1-imidazolyl
29	3-pyridyl	5-methyl-1-imidazolyl
30	3-pyridyl	2-methylsulfonyl-1-imidazolyl
31	2-pyrimidyl	2-(aminosulfonyl)phenyl
32	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
33	2-pyrimidyl	1-pyrrolidinocarbonyl
34	2-pyrimidyl	2-(methylsulfonyl)phenyl


35	2-pyrimidyl	4-morpholino
36	2-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
37	2-pyrimidyl	4-morpholinocarbonyl
38	2-pyrimidyl	2-methyl-1-imidazolyl
39	2-pyrimidyl	5-methyl-1-imidazolyl
40	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
41	5-pyrimidyl	2-(aminosulfonyl)phenyl
42	5-pyrimidyl	2-(methylaminosulfonyl)phenyl
43	5-pyrimidyl	1-pyrrolidinocarbonyl
44	5-pyrimidyl	2-(methylsulfonyl)phenyl
45	5-pyrimidyl	4-morpholino
46	5-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
47	5-pyrimidyl	4-morpholinocarbonyl
48	5-pyrimidyl	2-methyl-1-imidazolyl
49	5-pyrimidyl	5-methyl-1-imidazolyl
50	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
51	2-Cl-phenyl	2-(aminosulfonyl)phenyl
52	2-Cl-phenyl	2-(methylaminosulfonyl)phenyl
53	2-Cl-phenyl	1-pyrrolidinocarbonyl
54	2-Cl-phenyl	2-(methylsulfonyl)phenyl
55	2-Cl-phenyl	4-morpholino
56	2-Cl-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
57	2-Cl-phenyl	4-morpholinocarbonyl
58	2-Cl-phenyl	2-methyl-1-imidazolyl
59	2-Cl-phenyl	5-methyl-1-imidazolyl
60	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
61	2-F-phenyl	2-(aminosulfonyl)phenyl
62	2-F-phenyl	2-(methylaminosulfonyl)phenyl
63	2-F-phenyl	1-pyrrolidinocarbonyl
64	2-F-phenyl	2-(methylsulfonyl)phenyl
65	2-F-phenyl	4-morpholino
66	2-F-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
67	2-F-phenyl	4-morpholinocarbonyl
68	2-F-phenyl	2-methyl-1-imidazolyl
69	2-F-phenyl	5-methyl-1-imidazolyl
70	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
71	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
72	2,6-diF-phenyl	2-(methylaminosulfonyl)phenyl
73	2,6-diF-phenyl	1-pyrrolidinocarbonyl
74	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
75	2,6-diF-phenyl	4-morpholino
76	2,6-diF-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
77	2,6-diF-phenyl	4-morpholinocarbonyl
78	2,6-diF-phenyl	2-methyl-1-imidazolyl
79	2,6-diF-phenyl	5-methyl-1-imidazolyl
80	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl


Table 6














oo




pp




qq




rr




ss



tt



uu



vv

Ex #	R ^{1a}	A	B
1	CH ₃	phenyl	2-(aminosulfonyl)phenyl
2	CH ₃	phenyl	2-(methylaminosulfonyl)phenyl
3	CH ₃	phenyl	1-pyrrolidinocarbonyl
4	CH ₃	phenyl	2-(methylsulfonyl)phenyl
5	CH ₃	phenyl	4-morpholino
6	CH ₃	phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
7	CH ₃	phenyl	4-morpholinocarbonyl
8	CH ₃	phenyl	2-methyl-1-imidazolyl
9	CH ₃	phenyl	5-methyl-1-imidazolyl
10	CH ₃	phenyl	2-methylsulfonyl-1-imidazolyl
11	CH ₃	2-pyridyl	2-(aminosulfonyl)phenyl
12	CH ₃	2-pyridyl	2-(methylaminosulfonyl)phenyl
13	CH ₃	2-pyridyl	1-pyrrolidinocarbonyl
14	CH ₃	2-pyridyl	2-(methylsulfonyl)phenyl
15	CH ₃	2-pyridyl	4-morpholino
16	CH ₃	2-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
17	CH ₃	2-pyridyl	4-morpholinocarbonyl
18	CH ₃	2-pyridyl	2-methyl-1-imidazolyl
19	CH ₃	2-pyridyl	5-methyl-1-imidazolyl
20	CH ₃	2-pyridyl	2-methylsulfonyl-1-imidazolyl
21	CH ₃	3-pyridyl	2-(aminosulfonyl)phenyl
22	CH ₃	3-pyridyl	2-(methylaminosulfonyl)phenyl
23	CH ₃	3-pyridyl	1-pyrrolidinocarbonyl
24	CH ₃	3-pyridyl	2-(methylsulfonyl)phenyl
25	CH ₃	3-pyridyl	4-morpholino
26	CH ₃	3-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl

27	CH ₃	3-pyridyl	4-morpholinocarbonyl
28	CH ₃	3-pyridyl	2-methyl-1-imidazolyl
29	CH ₃	3-pyridyl	5-methyl-1-imidazolyl
30	CH ₃	3-pyridyl	2-methylsulfonyl-1-imidazolyl
31	CH ₃	2-pyrimidyl	2-(aminosulfonyl)phenyl
32	CH ₃	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
33	CH ₃	2-pyrimidyl	1-pyrrolidinocarbonyl
34	CH ₃	2-pyrimidyl	2-(methylsulfonyl)phenyl
35	CH ₃	2-pyrimidyl	4-morpholino
36	CH ₃	2-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
37	CH ₃	2-pyrimidyl	4-morpholinocarbonyl
38	CH ₃	2-pyrimidyl	2-methyl-1-imidazolyl
39	CH ₃	2-pyrimidyl	5-methyl-1-imidazolyl
40	CH ₃	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
41	CH ₃	5-pyrimidyl	2-(aminosulfonyl)phenyl
42	CH ₃	5-pyrimidyl	2-(methylaminosulfonyl)phenyl
43	CH ₃	5-pyrimidyl	1-pyrrolidinocarbonyl
44	CH ₃	5-pyrimidyl	2-(methylsulfonyl)phenyl
45	CH ₃	5-pyrimidyl	4-morpholino
46	CH ₃	5-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
47	CH ₃	5-pyrimidyl	4-morpholinocarbonyl
48	CH ₃	5-pyrimidyl	2-methyl-1-imidazolyl
49	CH ₃	5-pyrimidyl	5-methyl-1-imidazolyl
50	CH ₃	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
51	CH ₃	2-Cl-phenyl	2-(aminosulfonyl)phenyl
52	CH ₃	2-Cl-phenyl	2-(methylaminosulfonyl)phenyl
53	CH ₃	2-Cl-phenyl	1-pyrrolidinocarbonyl
54	CH ₃	2-Cl-phenyl	2-(methylsulfonyl)phenyl
55	CH ₃	2-Cl-phenyl	4-morpholino
56	CH ₃	2-Cl-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
57	CH ₃	2-Cl-phenyl	4-morpholinocarbonyl
58	CH ₃	2-Cl-phenyl	2-methyl-1-imidazolyl
59	CH ₃	2-Cl-phenyl	5-methyl-1-imidazolyl
60	CH ₃	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
61	CH ₃	2-F-phenyl	2-(aminosulfonyl)phenyl
62	CH ₃	2-F-phenyl	2-(methylaminosulfonyl)phenyl
63	CH ₃	2-F-phenyl	1-pyrrolidinocarbonyl
64	CH ₃	2-F-phenyl	2-(methylsulfonyl)phenyl
65	CH ₃	2-F-phenyl	4-morpholino
66	CH ₃	2-F-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
67	CH ₃	2-F-phenyl	4-morpholinocarbonyl
68	CH ₃	2-F-phenyl	2-methyl-1-imidazolyl
69	CH ₃	2-F-phenyl	5-methyl-1-imidazolyl
70	CH ₃	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
71	CH ₃	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
72	CH ₃	2,6-diF-phenyl	2-(methylaminosulfonyl)phenyl
73	CH ₃	2,6-diF-phenyl	1-pyrrolidinocarbonyl
74	CH ₃	2,6-diF-phenyl	2-(methylsulfonyl)phenyl

75	CH ₃	2,6-diF-phenyl	4-morpholino
76	CH ₃	2,6-diF-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
77	CH ₃	2,6-diF-phenyl	4-morpholinocarbonyl
78	CH ₃	2,6-diF-phenyl	2-methyl-1-imidazolyl
79	CH ₃	2,6-diF-phenyl	5-methyl-1-imidazolyl
80	CH ₃	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
81	CH ₂ CH ₃	phenyl	2-(aminosulfonyl)phenyl
82	CH ₂ CH ₃	phenyl	2-(methylaminosulfonyl)phenyl
83	CH ₂ CH ₃	phenyl	1-pyrrolidinocarbonyl
84	CH ₂ CH ₃	phenyl	2-(methylsulfonyl)phenyl
85	CH ₂ CH ₃	phenyl	4-morpholino
86	CH ₂ CH ₃	phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
87	CH ₂ CH ₃	phenyl	4-morpholinocarbonyl
88	CH ₂ CH ₃	phenyl	2-methyl-1-imidazolyl
89	CH ₂ CH ₃	phenyl	5-methyl-1-imidazolyl
90	CH ₂ CH ₃	phenyl	2-methylsulfonyl-1-imidazolyl
91	CH ₂ CH ₃	2-pyridyl	2-(aminosulfonyl)phenyl
92	CH ₂ CH ₃	2-pyridyl	2-(methylaminosulfonyl)phenyl
93	CH ₂ CH ₃	2-pyridyl	1-pyrrolidinocarbonyl
94	CH ₂ CH ₃	2-pyridyl	2-(methylsulfonyl)phenyl
95	CH ₂ CH ₃	2-pyridyl	4-morpholino
96	CH ₂ CH ₃	2-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
97	CH ₂ CH ₃	2-pyridyl	4-morpholinocarbonyl
98	CH ₂ CH ₃	2-pyridyl	2-methyl-1-imidazolyl
99	CH ₂ CH ₃	2-pyridyl	5-methyl-1-imidazolyl
100	CH ₂ CH ₃	2-pyridyl	2-methylsulfonyl-1-imidazolyl
101	CH ₂ CH ₃	3-pyridyl	2-(aminosulfonyl)phenyl
102	CH ₂ CH ₃	3-pyridyl	2-(methylaminosulfonyl)phenyl
103	CH ₂ CH ₃	3-pyridyl	1-pyrrolidinocarbonyl
104	CH ₂ CH ₃	3-pyridyl	2-(methylsulfonyl)phenyl
105	CH ₂ CH ₃	3-pyridyl	4-morpholino
106	CH ₂ CH ₃	3-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
107	CH ₂ CH ₃	3-pyridyl	4-morpholinocarbonyl
108	CH ₂ CH ₃	3-pyridyl	2-methyl-1-imidazolyl
109	CH ₂ CH ₃	3-pyridyl	5-methyl-1-imidazolyl
110	CH ₂ CH ₃	3-pyridyl	2-methylsulfonyl-1-imidazolyl
111	CH ₂ CH ₃	2-pyrimidyl	2-(aminosulfonyl)phenyl
112	CH ₂ CH ₃	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
113	CH ₂ CH ₃	2-pyrimidyl	1-pyrrolidinocarbonyl
114	CH ₂ CH ₃	2-pyrimidyl	2-(methylsulfonyl)phenyl
115	CH ₂ CH ₃	2-pyrimidyl	4-morpholino
116	CH ₂ CH ₃	2-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
117	CH ₂ CH ₃	2-pyrimidyl	4-morpholinocarbonyl
118	CH ₂ CH ₃	2-pyrimidyl	2-methyl-1-imidazolyl
119	CH ₂ CH ₃	2-pyrimidyl	5-methyl-1-imidazolyl
120	CH ₂ CH ₃	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
121	CH ₂ CH ₃	5-pyrimidyl	2-(aminosulfonyl)phenyl
122	CH ₂ CH ₃	5-pyrimidyl	2-(methylaminosulfonyl)phenyl

123	CH ₂ CH ₃	5-pyrimidyl	1-pyrrolidinocarbonyl
124	CH ₂ CH ₃	5-pyrimidyl	2-(methylsulfonyl)phenyl
125	CH ₂ CH ₃	5-pyrimidyl	4-morpholino
126	CH ₂ CH ₃	5-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
127	CH ₂ CH ₃	5-pyrimidyl	4-morpholinocarbonyl
128	CH ₂ CH ₃	5-pyrimidyl	2-methyl-1-imidazolyl
129	CH ₂ CH ₃	5-pyrimidyl	5-methyl-1-imidazolyl
130	CH ₂ CH ₃	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
131	CH ₂ CH ₃	2-Cl-phenyl	2-(aminosulfonyl)phenyl
132	CH ₂ CH ₃	2-Cl-phenyl	2-(methylaminosulfonyl)phenyl
133	CH ₂ CH ₃	2-Cl-phenyl	1-pyrrolidinocarbonyl
134	CH ₂ CH ₃	2-Cl-phenyl	2-(methylsulfonyl)phenyl
135	CH ₂ CH ₃	2-Cl-phenyl	4-morpholino
136	CH ₂ CH ₃	2-Cl-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
137	CH ₂ CH ₃	2-Cl-phenyl	4-morpholinocarbonyl
138	CH ₂ CH ₃	2-Cl-phenyl	2-methyl-1-imidazolyl
139	CH ₂ CH ₃	2-Cl-phenyl	5-methyl-1-imidazolyl
140	CH ₂ CH ₃	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
141	CH ₂ CH ₃	2-F-phenyl	2-(aminosulfonyl)phenyl
142	CH ₂ CH ₃	2-F-phenyl	2-(methylaminosulfonyl)phenyl
143	CH ₂ CH ₃	2-F-phenyl	1-pyrrolidinocarbonyl
144	CH ₂ CH ₃	2-F-phenyl	2-(methylsulfonyl)phenyl
145	CH ₂ CH ₃	2-F-phenyl	4-morpholino
146	CH ₂ CH ₃	2-F-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
147	CH ₂ CH ₃	2-F-phenyl	4-morpholinocarbonyl
148	CH ₂ CH ₃	2-F-phenyl	2-methyl-1-imidazolyl
149	CH ₂ CH ₃	2-F-phenyl	5-methyl-1-imidazolyl
150	CH ₂ CH ₃	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
151	CH ₂ CH ₃	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
152	CH ₂ CH ₃	2,6-diF-phenyl	2-(methylaminosulfonyl)phenyl
153	CH ₂ CH ₃	2,6-diF-phenyl	1-pyrrolidinocarbonyl
154	CH ₂ CH ₃	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
155	CH ₂ CH ₃	2,6-diF-phenyl	4-morpholino
156	CH ₂ CH ₃	2,6-diF-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
157	CH ₂ CH ₃	2,6-diF-phenyl	4-morpholinocarbonyl
158	CH ₂ CH ₃	2,6-diF-phenyl	2-methyl-1-imidazolyl
159	CH ₂ CH ₃	2,6-diF-phenyl	5-methyl-1-imidazolyl
160	CH ₂ CH ₃	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
161	CF ₃	phenyl	2-(aminosulfonyl)phenyl
162	CF ₃	phenyl	2-(methylaminosulfonyl)phenyl
163	CF ₃	phenyl	1-pyrrolidinocarbonyl
164	CF ₃	phenyl	2-(methylsulfonyl)phenyl
165	CF ₃	phenyl	4-morpholino
166	CF ₃	phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
167	CF ₃	phenyl	4-morpholinocarbonyl
168	CF ₃	phenyl	2-methyl-1-imidazolyl
169	CF ₃	phenyl	5-methyl-1-imidazolyl
170	CF ₃	phenyl	2-methylsulfonyl-1-imidazolyl

171	CF ₃	2-pyridyl	2-(aminosulfonyl)phenyl
172	CF ₃	2-pyridyl	2-(methylaminosulfonyl)phenyl
173	CF ₃	2-pyridyl	1-pyrrolidinocarbonyl
174	CF ₃	2-pyridyl	2-(methylsulfonyl)phenyl
175	CF ₃	2-pyridyl	4-morpholino
176	CF ₃	2-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
177	CF ₃	2-pyridyl	4-morpholinocarbonyl
178	CF ₃	2-pyridyl	2-methyl-1-imidazolyl
179	CF ₃	2-pyridyl	5-methyl-1-imidazolyl
180	CF ₃	2-pyridyl	2-methylsulfonyl-1-imidazolyl
181	CF ₃	3-pyridyl	2-(aminosulfonyl)phenyl
182	CF ₃	3-pyridyl	2-(methylaminosulfonyl)phenyl
183	CF ₃	3-pyridyl	1-pyrrolidinocarbonyl
184	CF ₃	3-pyridyl	2-(methylsulfonyl)phenyl
185	CF ₃	3-pyridyl	4-morpholino
186	CF ₃	3-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
187	CF ₃	3-pyridyl	4-morpholinocarbonyl
188	CF ₃	3-pyridyl	2-methyl-1-imidazolyl
189	CF ₃	3-pyridyl	5-methyl-1-imidazolyl
190	CF ₃	3-pyridyl	2-methylsulfonyl-1-imidazolyl
191	CF ₃	2-pyrimidyl	2-(aminosulfonyl)phenyl
192	CF ₃	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
193	CF ₃	2-pyrimidyl	1-pyrrolidinocarbonyl
194	CF ₃	2-pyrimidyl	2-(methylsulfonyl)phenyl
195	CF ₃	2-pyrimidyl	4-morpholino
196	CF ₃	2-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
197	CF ₃	2-pyrimidyl	4-morpholinocarbonyl
198	CF ₃	2-pyrimidyl	2-methyl-1-imidazolyl
199	CF ₃	2-pyrimidyl	5-methyl-1-imidazolyl
200	CF ₃	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
201	CF ₃	5-pyrimidyl	2-(aminosulfonyl)phenyl
202	CF ₃	5-pyrimidyl	2-(methylaminosulfonyl)phenyl
203	CF ₃	5-pyrimidyl	1-pyrrolidinocarbonyl
204	CF ₃	5-pyrimidyl	2-(methylsulfonyl)phenyl
205	CF ₃	5-pyrimidyl	4-morpholino
206	CF ₃	5-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
207	CF ₃	5-pyrimidyl	4-morpholinocarbonyl
208	CF ₃	5-pyrimidyl	2-methyl-1-imidazolyl
209	CF ₃	5-pyrimidyl	5-methyl-1-imidazolyl
210	CF ₃	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
211	CF ₃	2-Cl-phenyl	2-(aminosulfonyl)phenyl
212	CF ₃	2-Cl-phenyl	2-(methylaminosulfonyl)phenyl
213	CF ₃	2-Cl-phenyl	1-pyrrolidinocarbonyl
214	CF ₃	2-Cl-phenyl	2-(methylsulfonyl)phenyl
215	CF ₃	2-Cl-phenyl	4-morpholino
216	CF ₃	2-Cl-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
217	CF ₃	2-Cl-phenyl	4-morpholinocarbonyl
218	CF ₃	2-Cl-phenyl	2-methyl-1-imidazolyl

219	CF ₃	2-Cl-phenyl	5-methyl-1-imidazolyl
220	CF ₃	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
221	CF ₃	2-F-phenyl	2-(aminosulfonyl)phenyl
222	CF ₃	2-F-phenyl	2-(methyaminosulfonyl)phenyl
223	CF ₃	2-F-phenyl	1-pyrrolidinocarbonyl
224	CF ₃	2-F-phenyl	2-(methylsulfonyl)phenyl
225	CF ₃	2-F-phenyl	4-morpholino
226	CF ₃	2-F-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
227	CF ₃	2-F-phenyl	4-morpholinocarbonyl
228	CF ₃	2-F-phenyl	2-methyl-1-imidazolyl
229	CF ₃	2-F-phenyl	5-methyl-1-imidazolyl
230	CF ₃	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
231	CF ₃	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
232	CF ₃	2,6-diF-phenyl	2-(methyaminosulfonyl)phenyl
233	CF ₃	2,6-diF-phenyl	1-pyrrolidinocarbonyl
234	CF ₃	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
235	CF ₃	2,6-diF-phenyl	4-morpholino
236	CF ₃	2,6-diF-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
237	CF ₃	2,6-diF-phenyl	4-morpholinocarbonyl
238	CF ₃	2,6-diF-phenyl	2-methyl-1-imidazolyl
239	CF ₃	2,6-diF-phenyl	5-methyl-1-imidazolyl
240	CF ₃	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
241	SCH ₃	phenyl	2-(aminosulfonyl)phenyl
242	SCH ₃	phenyl	2-(methyaminosulfonyl)phenyl
243	SCH ₃	phenyl	1-pyrrolidinocarbonyl
244	SCH ₃	phenyl	2-(methylsulfonyl)phenyl
245	SCH ₃	phenyl	4-morpholino
246	SCH ₃	phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
247	SCH ₃	phenyl	4-morpholinocarbonyl
248	SCH ₃	phenyl	2-methyl-1-imidazolyl
249	SCH ₃	phenyl	5-methyl-1-imidazolyl
250	SCH ₃	phenyl	2-methylsulfonyl-1-imidazolyl
251	SCH ₃	2-pyridyl	2-(aminosulfonyl)phenyl
252	SCH ₃	2-pyridyl	2-(methyaminosulfonyl)phenyl
253	SCH ₃	2-pyridyl	1-pyrrolidinocarbonyl
254	SCH ₃	2-pyridyl	2-(methylsulfonyl)phenyl
255	SCH ₃	2-pyridyl	4-morpholino
256	SCH ₃	2-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
257	SCH ₃	2-pyridyl	4-morpholinocarbonyl
258	SCH ₃	2-pyridyl	2-methyl-1-imidazolyl
259	SCH ₃	2-pyridyl	5-methyl-1-imidazolyl
260	SCH ₃	2-pyridyl	2-methylsulfonyl-1-imidazolyl
261	SCH ₃	3-pyridyl	2-(aminosulfonyl)phenyl
262	SCH ₃	3-pyridyl	2-(methyaminosulfonyl)phenyl
263	SCH ₃	3-pyridyl	1-pyrrolidinocarbonyl
264	SCH ₃	3-pyridyl	2-(methylsulfonyl)phenyl
265	SCH ₃	3-pyridyl	4-morpholino
266	SCH ₃	3-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl

267	SCH ₃	3-pyridyl	4-morpholinocarbonyl
268	SCH ₃	3-pyridyl	2-methyl-1-imidazolyl
269	SCH ₃	3-pyridyl	5-methyl-1-imidazolyl
270	SCH ₃	3-pyridyl	2-methylsulfonyl-1-imidazolyl
271	SCH ₃	2-pyrimidyl	2-(aminosulfonyl)phenyl
272	SCH ₃	2-pyrimidyl	2-(methylaninosulfonyl)phenyl
273	SCH ₃	2-pyrimidyl	1-pyrrolidinocarbonyl
274	SCH ₃	2-pyrimidyl	2-(methylsulfonyl)phenyl
275	SCH ₃	2-pyrimidyl	4-morpholino
276	SCH ₃	2-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
277	SCH ₃	2-pyrimidyl	4-morpholinocarbonyl
278	SCH ₃	2-pyrimidyl	2-methyl-1-imidazolyl
279	SCH ₃	2-pyrimidyl	5-methyl-1-imidazolyl
280	SCH ₃	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
281	SCH ₃	5-pyrimidyl	2-(aminosulfonyl)phenyl
282	SCH ₃	5-pyrimidyl	2-(methylaninosulfonyl)phenyl
283	SCH ₃	5-pyrimidyl	1-pyrrolidinocarbonyl
284	SCH ₃	5-pyrimidyl	2-(methylsulfonyl)phenyl
285	SCH ₃	5-pyrimidyl	4-morpholino
286	SCH ₃	5-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
287	SCH ₃	5-pyrimidyl	4-morpholinocarbonyl
288	SCH ₃	5-pyrimidyl	2-methyl-1-imidazolyl
289	SCH ₃	5-pyrimidyl	5-methyl-1-imidazolyl
290	SCH ₃	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
291	SCH ₃	2-Cl-phenyl	2-(aminosulfonyl)phenyl
292	SCH ₃	2-Cl-phenyl	2-(methylaninosulfonyl)phenyl
293	SCH ₃	2-Cl-phenyl	1-pyrrolidinocarbonyl
294	SCH ₃	2-Cl-phenyl	2-(methylsulfonyl)phenyl
295	SCH ₃	2-Cl-phenyl	4-morpholino
296	SCH ₃	2-Cl-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
297	SCH ₃	2-Cl-phenyl	4-morpholinocarbonyl
298	SCH ₃	2-Cl-phenyl	2-methyl-1-imidazolyl
299	SCH ₃	2-Cl-phenyl	5-methyl-1-imidazolyl
300	SCH ₃	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
301	SCH ₃	2-F-phenyl	2-(aminosulfonyl)phenyl
302	SCH ₃	2-F-phenyl	2-(methylaninosulfonyl)phenyl
303	SCH ₃	2-F-phenyl	1-pyrrolidinocarbonyl
304	SCH ₃	2-F-phenyl	2-(methylsulfonyl)phenyl
305	SCH ₃	2-F-phenyl	4-morpholino
306	SCH ₃	2-F-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
307	SCH ₃	2-F-phenyl	4-morpholinocarbonyl
308	SCH ₃	2-F-phenyl	2-methyl-1-imidazolyl
309	SCH ₃	2-F-phenyl	5-methyl-1-imidazolyl
310	SCH ₃	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
311	SCH ₃	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
312	SCH ₃	2,6-diF-phenyl	2-(methylaninosulfonyl)phenyl
313	SCH ₃	2,6-diF-phenyl	1-pyrrolidinocarbonyl
314	SCH ₃	2,6-diF-phenyl	2-(methylsulfonyl)phenyl

315	SCH ₃	2,6-diF-phenyl	4-morpholino
316	SCH ₃	2,6-diF-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
317	SCH ₃	2,6-diF-phenyl	4-morpholinocarbonyl
318	SCH ₃	2,6-diF-phenyl	2-methyl-1-imidazolyl
319	SCH ₃	2,6-diF-phenyl	5-methyl-1-imidazolyl
320	SCH ₃	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
321	SOCH ₃	phenyl	2-(aminosulfonyl)phenyl
322	SOCH ₃	phenyl	2-(methylaminosulfonyl)phenyl
323	SOCH ₃	phenyl	1-pyrrolidinocarbonyl
324	SOCH ₃	phenyl	2-(methylsulfonyl)phenyl
325	SOCH ₃	phenyl	4-morpholino
326	SOCH ₃	phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
327	SOCH ₃	phenyl	4-morpholinocarbonyl
328	SOCH ₃	phenyl	2-methyl-1-imidazolyl
329	SOCH ₃	phenyl	5-methyl-1-imidazolyl
330	SOCH ₃	phenyl	2-methylsulfonyl-1-imidazolyl
331	SOCH ₃	2-pyridyl	2-(aminosulfonyl)phenyl
332	SOCH ₃	2-pyridyl	2-(methylaminosulfonyl)phenyl
333	SOCH ₃	2-pyridyl	1-pyrrolidinocarbonyl
334	SOCH ₃	2-pyridyl	2-(methylsulfonyl)phenyl
335	SOCH ₃	2-pyridyl	4-morpholino
336	SOCH ₃	2-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
337	SOCH ₃	2-pyridyl	4-morpholinocarbonyl
338	SOCH ₃	2-pyridyl	2-methyl-1-imidazolyl
339	SOCH ₃	2-pyridyl	5-methyl-1-imidazolyl
340	SOCH ₃	2-pyridyl	2-methylsulfonyl-1-imidazolyl
341	SOCH ₃	3-pyridyl	2-(aminosulfonyl)phenyl
342	SOCH ₃	3-pyridyl	2-(methylaminosulfonyl)phenyl
343	SOCH ₃	3-pyridyl	1-pyrrolidinocarbonyl
344	SOCH ₃	3-pyridyl	2-(methylsulfonyl)phenyl
345	SOCH ₃	3-pyridyl	4-morpholino
346	SOCH ₃	3-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
347	SOCH ₃	3-pyridyl	4-morpholinocarbonyl
348	SOCH ₃	3-pyridyl	2-methyl-1-imidazolyl
349	SOCH ₃	3-pyridyl	5-methyl-1-imidazolyl
350	SOCH ₃	3-pyridyl	2-methylsulfonyl-1-imidazolyl
351	SOCH ₃	2-pyrimidyl	2-(aminosulfonyl)phenyl
352	SOCH ₃	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
353	SOCH ₃	2-pyrimidyl	1-pyrrolidinocarbonyl
354	SOCH ₃	2-pyrimidyl	2-(methylsulfonyl)phenyl
355	SOCH ₃	2-pyrimidyl	4-morpholino
356	SOCH ₃	2-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
357	SOCH ₃	2-pyrimidyl	4-morpholinocarbonyl
358	SOCH ₃	2-pyrimidyl	2-methyl-1-imidazolyl
359	SOCH ₃	2-pyrimidyl	5-methyl-1-imidazolyl
360	SOCH ₃	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
361	SOCH ₃	5-pyrimidyl	2-(aminosulfonyl)phenyl
362	SOCH ₃	5-pyrimidyl	2-(methylaminosulfonyl)phenyl

363	SOCH ₃	5-pyrimidyl	1-pyrrolidinocarbonyl
364	SOCH ₃	5-pyrimidyl	2-(methylsulfonyl)phenyl
365	SOCH ₃	5-pyrimidyl	4-morpholino
366	SOCH ₃	5-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
367	SOCH ₃	5-pyrimidyl	4-morpholinocarbonyl
368	SOCH ₃	5-pyrimidyl	2-methyl-1-imidazolyl
369	SOCH ₃	5-pyrimidyl	5-methyl-1-imidazolyl
370	SOCH ₃	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
371	SOCH ₃	2-Cl-phenyl	2-(aminosulfonyl)phenyl
372	SOCH ₃	2-Cl-phenyl	2-(methylaminosulfonyl)phenyl
373	SOCH ₃	2-Cl-phenyl	1-pyrrolidinocarbonyl
374	SOCH ₃	2-Cl-phenyl	2-(methylsulfonyl)phenyl
375	SOCH ₃	2-Cl-phenyl	4-morpholino
376	SOCH ₃	2-Cl-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
377	SOCH ₃	2-Cl-phenyl	4-morpholinocarbonyl
378	SOCH ₃	2-Cl-phenyl	2-methyl-1-imidazolyl
379	SOCH ₃	2-Cl-phenyl	5-methyl-1-imidazolyl
380	SOCH ₃	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
381	SOCH ₃	2-F-phenyl	2-(aminosulfonyl)phenyl
382	SOCH ₃	2-F-phenyl	2-(methylaminosulfonyl)phenyl
383	SOCH ₃	2-F-phenyl	1-pyrrolidinocarbonyl
384	SOCH ₃	2-F-phenyl	2-(methylsulfonyl)phenyl
385	SOCH ₃	2-F-phenyl	4-morpholino
386	SOCH ₃	2-F-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
387	SOCH ₃	2-F-phenyl	4-morpholinocarbonyl
388	SOCH ₃	2-F-phenyl	2-methyl-1-imidazolyl
389	SOCH ₃	2-F-phenyl	5-methyl-1-imidazolyl
390	SOCH ₃	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
391	SOCH ₃	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
392	SOCH ₃	2,6-diF-phenyl	2-(methylaminosulfonyl)phenyl
393	SOCH ₃	2,6-diF-phenyl	1-pyrrolidinocarbonyl
394	SOCH ₃	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
395	SOCH ₃	2,6-diF-phenyl	4-morpholino
396	SOCH ₃	2,6-diF-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
397	SOCH ₃	2,6-diF-phenyl	4-morpholinocarbonyl
398	SOCH ₃	2,6-diF-phenyl	2-methyl-1-imidazolyl
399	SOCH ₃	2,6-diF-phenyl	5-methyl-1-imidazolyl
400	SOCH ₃	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
401	SO ₂ CH ₃	phenyl	2-(aminosulfonyl)phenyl
402	SO ₂ CH ₃	phenyl	2-(methylaminosulfonyl)phenyl
403	SO ₂ CH ₃	phenyl	1-pyrrolidinocarbonyl
404	SO ₂ CH ₃	phenyl	2-(methylsulfonyl)phenyl
405	SO ₂ CH ₃	phenyl	4-morpholino
406	SO ₂ CH ₃	phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
407	SO ₂ CH ₃	phenyl	4-morpholinocarbonyl
408	SO ₂ CH ₃	phenyl	2-methyl-1-imidazolyl
409	SO ₂ CH ₃	phenyl	5-methyl-1-imidazolyl
410	SO ₂ CH ₃	phenyl	2-methylsulfonyl-1-imidazolyl

411	SO ₂ CH ₃	2-pyridyl	2-(aminosulfonyl)phenyl
412	SO ₂ CH ₃	2-pyridyl	2-(methylaninosulfonyl)phenyl
413	SO ₂ CH ₃	2-pyridyl	1-pyrrolidinocarbonyl
414	SO ₂ CH ₃	2-pyridyl	2-(methylsulfonyl)phenyl
415	SO ₂ CH ₃	2-pyridyl	4-morpholino
416	SO ₂ CH ₃	2-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
417	SO ₂ CH ₃	2-pyridyl	4-morpholinocarbonyl
418	SO ₂ CH ₃	2-pyridyl	2-methyl-1-imidazolyl
419	SO ₂ CH ₃	2-pyridyl	5-methyl-1-imidazolyl
420	SO ₂ CH ₃	2-pyridyl	2-methylsulfonyl-1-imidazolyl
421	SO ₂ CH ₃	3-pyridyl	2-(aminosulfonyl)phenyl
422	SO ₂ CH ₃	3-pyridyl	2-(methylaninosulfonyl)phenyl
423	SO ₂ CH ₃	3-pyridyl	1-pyrrolidinocarbonyl
424	SO ₂ CH ₃	3-pyridyl	2-(methylsulfonyl)phenyl
425	SO ₂ CH ₃	3-pyridyl	4-morpholino
426	SO ₂ CH ₃	3-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
427	SO ₂ CH ₃	3-pyridyl	4-morpholinocarbonyl
428	SO ₂ CH ₃	3-pyridyl	2-methyl-1-imidazolyl
429	SO ₂ CH ₃	3-pyridyl	5-methyl-1-imidazolyl
430	SO ₂ CH ₃	3-pyridyl	2-methylsulfonyl-1-imidazolyl
431	SO ₂ CH ₃	2-pyrimidyl	2-(aminosulfonyl)phenyl
432	SO ₂ CH ₃	2-pyrimidyl	2-(methylaninosulfonyl)phenyl
433	SO ₂ CH ₃	2-pyrimidyl	1-pyrrolidinocarbonyl
434	SO ₂ CH ₃	2-pyrimidyl	2-(methylsulfonyl)phenyl
435	SO ₂ CH ₃	2-pyrimidyl	4-morpholino
436	SO ₂ CH ₃	2-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
437	SO ₂ CH ₃	2-pyrimidyl	4-morpholinocarbonyl
438	SO ₂ CH ₃	2-pyrimidyl	2-methyl-1-imidazolyl
439	SO ₂ CH ₃	2-pyrimidyl	5-methyl-1-imidazolyl
440	SO ₂ CH ₃	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
441	SO ₂ CH ₃	5-pyrimidyl	2-(aminosulfonyl)phenyl
442	SO ₂ CH ₃	5-pyrimidyl	2-(methylaninosulfonyl)phenyl
443	SO ₂ CH ₃	5-pyrimidyl	1-pyrrolidinocarbonyl
444	SO ₂ CH ₃	5-pyrimidyl	2-(methylsulfonyl)phenyl
445	SO ₂ CH ₃	5-pyrimidyl	4-morpholino
446	SO ₂ CH ₃	5-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
447	SO ₂ CH ₃	5-pyrimidyl	4-morpholinocarbonyl
448	SO ₂ CH ₃	5-pyrimidyl	2-methyl-1-imidazolyl
449	SO ₂ CH ₃	5-pyrimidyl	5-methyl-1-imidazolyl
450	SO ₂ CH ₃	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
451	SO ₂ CH ₃	2-Cl-phenyl	2-(aminosulfonyl)phenyl
452	SO ₂ CH ₃	2-Cl-phenyl	2-(methylaninosulfonyl)phenyl
453	SO ₂ CH ₃	2-Cl-phenyl	1-pyrrolidinocarbonyl
454	SO ₂ CH ₃	2-Cl-phenyl	2-(methylsulfonyl)phenyl
455	SO ₂ CH ₃	2-Cl-phenyl	4-morpholino
456	SO ₂ CH ₃	2-Cl-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
457	SO ₂ CH ₃	2-Cl-phenyl	4-morpholinocarbonyl
458	SO ₂ CH ₃	2-Cl-phenyl	2-methyl-1-imidazolyl

459	SO ₂ CH ₃	2-Cl-phenyl	5-methyl-1-imidazolyl
460	SO ₂ CH ₃	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
461	SO ₂ CH ₃	2-F-phenyl	2-(aminosulfonyl)phenyl
462	SO ₂ CH ₃	2-F-phenyl	2-(methyaminosulfonyl)phenyl
463	SO ₂ CH ₃	2-F-phenyl	1-pyrrolidinocarbonyl
464	SO ₂ CH ₃	2-F-phenyl	2-(methylsulfonyl)phenyl
465	SO ₂ CH ₃	2-F-phenyl	4-morpholino
466	SO ₂ CH ₃	2-F-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
467	SO ₂ CH ₃	2-F-phenyl	4-morpholinocarbonyl
468	SO ₂ CH ₃	2-F-phenyl	2-methyl-1-imidazolyl
469	SO ₂ CH ₃	2-F-phenyl	5-methyl-1-imidazolyl
470	SO ₂ CH ₃	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
471	SO ₂ CH ₃	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
472	SO ₂ CH ₃	2,6-diF-phenyl	2-(methyaminosulfonyl)phenyl
473	SO ₂ CH ₃	2,6-diF-phenyl	1-pyrrolidinocarbonyl
474	SO ₂ CH ₃	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
475	SO ₂ CH ₃	2,6-diF-phenyl	4-morpholino
476	SO ₂ CH ₃	2,6-diF-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
477	SO ₂ CH ₃	2,6-diF-phenyl	4-morpholinocarbonyl
478	SO ₂ CH ₃	2,6-diF-phenyl	2-methyl-1-imidazolyl
479	SO ₂ CH ₃	2,6-diF-phenyl	5-methyl-1-imidazolyl
480	SO ₂ CH ₃	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
481	CH ₂ NH- SO ₂ CH ₃	phenyl	2-(aminosulfonyl)phenyl
482	CH ₂ NH- SO ₂ CH ₃	phenyl	2-(methyaminosulfonyl)phenyl
483	CH ₂ NH- SO ₂ CH ₃	phenyl	1-pyrrolidinocarbonyl
484	CH ₂ NH- SO ₂ CH ₃	phenyl	2-(methylsulfonyl)phenyl
485	CH ₂ NH- SO ₂ CH ₃	phenyl	4-morpholino
486	CH ₂ NH- SO ₂ CH ₃	phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
487	CH ₂ NH- SO ₂ CH ₃	phenyl	4-morpholinocarbonyl
488	CH ₂ NH- SO ₂ CH ₃	phenyl	2-methyl-1-imidazolyl
489	CH ₂ NH- SO ₂ CH ₃	phenyl	5-methyl-1-imidazolyl
490	CH ₂ NH- SO ₂ CH ₃	phenyl	2-methylsulfonyl-1-imidazolyl
491	CH ₂ NH- SO ₂ CH ₃	2-pyridyl	2-(aminosulfonyl)phenyl
492	CH ₂ NH- SO ₂ CH ₃	2-pyridyl	2-(methyaminosulfonyl)phenyl
493	CH ₂ NH- SO ₂ CH ₃	2-pyridyl	1-pyrrolidinocarbonyl

494	CH ₂ NH-SO ₂ CH ₃	2-pyridyl	2-(methylsulfonyl)phenyl
495	CH ₂ NH-SO ₂ CH ₃	2-pyridyl	4-morpholino
496	CH ₂ NH-SO ₂ CH ₃	2-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
497	CH ₂ NH-SO ₂ CH ₃	2-pyridyl	4-morpholinocarbonyl
498	CH ₂ NH-SO ₂ CH ₃	2-pyridyl	2-methyl-1-imidazolyl
499	CH ₂ NH-SO ₂ CH ₃	2-pyridyl	5-methyl-1-imidazolyl
500	CH ₂ NH-SO ₂ CH ₃	2-pyridyl	2-methylsulfonyl-1-imidazolyl
501	CH ₂ NH-SO ₂ CH ₃	3-pyridyl	2-(aminosulfonyl)phenyl
502	CH ₂ NH-SO ₂ CH ₃	3-pyridyl	2-(methylaminosulfonyl)phenyl
503	CH ₂ NH-SO ₂ CH ₃	3-pyridyl	1-pyrrolidinocarbonyl
504	CH ₂ NH-SO ₂ CH ₃	3-pyridyl	2-(methylsulfonyl)phenyl
505	CH ₂ NH-SO ₂ CH ₃	3-pyridyl	4-morpholino
506	CH ₂ NH-SO ₂ CH ₃	3-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
507	CH ₂ NH-SO ₂ CH ₃	3-pyridyl	4-morpholinocarbonyl
508	CH ₂ NH-SO ₂ CH ₃	3-pyridyl	2-methyl-1-imidazolyl
509	CH ₂ NH-SO ₂ CH ₃	3-pyridyl	5-methyl-1-imidazolyl
510	CH ₂ NH-SO ₂ CH ₃	3-pyridyl	2-methylsulfonyl-1-imidazolyl
511	CH ₂ NH-SO ₂ CH ₃	2-pyrimidyl	2-(aminosulfonyl)phenyl
512	CH ₂ NH-SO ₂ CH ₃	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
513	CH ₂ NH-SO ₂ CH ₃	2-pyrimidyl	1-pyrrolidinocarbonyl
514	CH ₂ NH-SO ₂ CH ₃	2-pyrimidyl	2-(methylsulfonyl)phenyl
515	CH ₂ NH-SO ₂ CH ₃	2-pyrimidyl	4-morpholino
516	CH ₂ NH-SO ₂ CH ₃	2-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
517	CH ₂ NH-SO ₂ CH ₃	2-pyrimidyl	4-morpholinocarbonyl

518	CH ₂ NH-SO ₂ CH ₃	2-pyrimidyl	2-methyl-1-imidazolyl
519	CH ₂ NH-SO ₂ CH ₃	2-pyrimidyl	5-methyl-1-imidazolyl
520	CH ₂ NH-SO ₂ CH ₃	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
521	CH ₂ NH-SO ₂ CH ₃	5-pyrimidyl	2-(aminosulfonyl)phenyl
522	CH ₂ NH-SO ₂ CH ₃	5-pyrimidyl	2-(methylaminosulfonyl)phenyl
523	CH ₂ NH-SO ₂ CH ₃	5-pyrimidyl	1-pyrrolidinocarbonyl
524	CH ₂ NH-SO ₂ CH ₃	5-pyrimidyl	2-(methylsulfonyl)phenyl
525	CH ₂ NH-SO ₂ CH ₃	5-pyrimidyl	4-morpholino
526	CH ₂ NH-SO ₂ CH ₃	5-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
527	CH ₂ NH-SO ₂ CH ₃	5-pyrimidyl	4-morpholinocarbonyl
528	CH ₂ NH-SO ₂ CH ₃	5-pyrimidyl	2-methyl-1-imidazolyl
529	CH ₂ NH-SO ₂ CH ₃	5-pyrimidyl	5-methyl-1-imidazolyl
530	CH ₂ NH-SO ₂ CH ₃	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
531	CH ₂ NH-SO ₂ CH ₃	2-Cl-phenyl	2-(aminosulfonyl)phenyl
532	CH ₂ NH-SO ₂ CH ₃	2-Cl-phenyl	2-(methylaminosulfonyl)phenyl
533	CH ₂ NH-SO ₂ CH ₃	2-Cl-phenyl	1-pyrrolidinocarbonyl
534	CH ₂ NH-SO ₂ CH ₃	2-Cl-phenyl	2-(methylsulfonyl)phenyl
535	CH ₂ NH-SO ₂ CH ₃	2-Cl-phenyl	4-morpholino
536	CH ₂ NH-SO ₂ CH ₃	2-Cl-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
537	CH ₂ NH-SO ₂ CH ₃	2-Cl-phenyl	4-morpholinocarbonyl
538	CH ₂ NH-SO ₂ CH ₃	2-Cl-phenyl	2-methyl-1-imidazolyl
539	CH ₂ NH-SO ₂ CH ₃	2-Cl-phenyl	5-methyl-1-imidazolyl
540	CH ₂ NH-SO ₂ CH ₃	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
541	CH ₂ NH-SO ₂ CH ₃	2-F-phenyl	2-(aminosulfonyl)phenyl

WO 98/28282

542	CH ₂ NH-SO ₂ CH ₃	2-F-phenyl	2-(methylaminosulfonyl)phenyl
543	CH ₂ NH-SO ₂ CH ₃	2-F-phenyl	1-pyrrolidinocarbonyl
544	CH ₂ NH-SO ₂ CH ₃	2-F-phenyl	2-(methylsulfonyl)phenyl
545	CH ₂ NH-SO ₂ CH ₃	2-F-phenyl	4-morpholino
546	CH ₂ NH-SO ₂ CH ₃	2-F-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
547	CH ₂ NH-SO ₂ CH ₃	2-F-phenyl	4-morpholinocarbonyl
548	CH ₂ NH-SO ₂ CH ₃	2-F-phenyl	2-methyl-1-imidazolyl
549	CH ₂ NH-SO ₂ CH ₃	2-F-phenyl	5-methyl-1-imidazolyl
550	CH ₂ NH-SO ₂ CH ₃	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
551	CH ₂ NH-SO ₂ CH ₃	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
552	CH ₂ NH-SO ₂ CH ₃	2,6-diF-phenyl	2-(methylaminosulfonyl)phenyl
553	CH ₂ NH-SO ₂ CH ₃	2,6-diF-phenyl	1-pyrrolidinocarbonyl
554	CH ₂ NH-SO ₂ CH ₃	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
555	CH ₂ NH-SO ₂ CH ₃	2,6-diF-phenyl	4-morpholino
556	CH ₂ NH-SO ₂ CH ₃	2,6-diF-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
557	CH ₂ NH-SO ₂ CH ₃	2,6-diF-phenyl	4-morpholinocarbonyl
558	CH ₂ NH-SO ₂ CH ₃	2,6-diF-phenyl	2-methyl-1-imidazolyl
559	CH ₂ NH-SO ₂ CH ₃	2,6-diF-phenyl	5-methyl-1-imidazolyl
560	CH ₂ NH-SO ₂ CH ₃	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
561	Cl	phenyl	2-(aminosulfonyl)phenyl
562	Cl	phenyl	2-(methylaminosulfonyl)phenyl
563	Cl	phenyl	1-pyrrolidinocarbonyl
564	Cl	phenyl	2-(methylsulfonyl)phenyl
565	Cl	phenyl	4-morpholino
566	Cl	phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
567	Cl	phenyl	4-morpholinocarbonyl
568	Cl	phenyl	2-methyl-1-imidazolyl
569	Cl	phenyl	5-methyl-1-imidazolyl
570	Cl	phenyl	2-methylsulfonyl-1-imidazolyl
571	Cl	2-pyridyl	2-(aminosulfonyl)phenyl
572	Cl	2-pyridyl	2-(methylaminosulfonyl)phenyl

573	Cl	2-pyridyl	1-pyrrolidinocarbonyl
574	Cl	2-pyridyl	2-(methylsulfonyl)phenyl
575	Cl	2-pyridyl	4-morpholino
576	Cl	2-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
577	Cl	2-pyridyl	4-morpholinocarbonyl
578	Cl	2-pyridyl	2-methyl-1-imidazolyl
579	Cl	2-pyridyl	5-methyl-1-imidazolyl
580	Cl	2-pyridyl	2-methylsulfonyl-1-imidazolyl
581	Cl	3-pyridyl	2-(aminosulfonyl)phenyl
582	Cl	3-pyridyl	2-(methylaminosulfonyl)phenyl
583	Cl	3-pyridyl	1-pyrrolidinocarbonyl
584	Cl	3-pyridyl	2-(methylsulfonyl)phenyl
585	Cl	3-pyridyl	4-morpholino
586	Cl	3-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
587	Cl	3-pyridyl	4-morpholinocarbonyl
588	Cl	3-pyridyl	2-methyl-1-imidazolyl
589	Cl	3-pyridyl	5-methyl-1-imidazolyl
590	Cl	3-pyridyl	2-methylsulfonyl-1-imidazolyl
591	Cl	2-pyrimidyl	2-(aminosulfonyl)phenyl
592	Cl	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
593	Cl	2-pyrimidyl	1-pyrrolidinocarbonyl
594	Cl	2-pyrimidyl	2-(methylsulfonyl)phenyl
595	Cl	2-pyrimidyl	4-morpholino
596	Cl	2-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
597	Cl	2-pyrimidyl	4-morpholinocarbonyl
598	Cl	2-pyrimidyl	2-methyl-1-imidazolyl
599	Cl	2-pyrimidyl	5-methyl-1-imidazolyl
600	Cl	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
601	Cl	5-pyrimidyl	2-(aminosulfonyl)phenyl
602	Cl	5-pyrimidyl	2-(methylaminosulfonyl)phenyl
603	Cl	5-pyrimidyl	1-pyrrolidinocarbonyl
604	Cl	5-pyrimidyl	2-(methylsulfonyl)phenyl
605	Cl	5-pyrimidyl	4-morpholino
606	Cl	5-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
607	Cl	5-pyrimidyl	4-morpholinocarbonyl
608	Cl	5-pyrimidyl	2-methyl-1-imidazolyl
609	Cl	5-pyrimidyl	5-methyl-1-imidazolyl
610	Cl	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
611	Cl	2-Cl-phenyl	2-(aminosulfonyl)phenyl
612	Cl	2-Cl-phenyl	2-(methylaminosulfonyl)phenyl
613	Cl	2-Cl-phenyl	1-pyrrolidinocarbonyl
614	Cl	2-Cl-phenyl	2-(methylsulfonyl)phenyl
615	Cl	2-Cl-phenyl	4-morpholino
616	Cl	2-Cl-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
617	Cl	2-Cl-phenyl	4-morpholinocarbonyl
618	Cl	2-Cl-phenyl	2-methyl-1-imidazolyl
619	Cl	2-Cl-phenyl	5-methyl-1-imidazolyl
620	Cl	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
621	Cl	2-F-phenyl	2-(aminosulfonyl)phenyl
622	Cl	2-F-phenyl	2-(methylaminosulfonyl)phenyl
623	Cl	2-F-phenyl	1-pyrrolidinocarbonyl
624	Cl	2-F-phenyl	2-(methylsulfonyl)phenyl
625	Cl	2-F-phenyl	4-morpholino
626	Cl	2-F-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
627	Cl	2-F-phenyl	4-morpholinocarbonyl

WO 98/28282

628	Cl	2-F-phenyl	2-methyl-1-imidazolyl
629	Cl	2-F-phenyl	5-methyl-1-imidazolyl
630	Cl	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
631	Cl	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
632	Cl	2,6-diF-phenyl	2-(methyaminosulfonyl)phenyl
633	Cl	2,6-diF-phenyl	1-pyrrolidinocarbonyl
634	Cl	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
635	Cl	2,6-diF-phenyl	4-morpholino
636	Cl	2,6-diF-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
637	Cl	2,6-diF-phenyl	4-morpholinocarbonyl
638	Cl	2,6-diF-phenyl	2-methyl-1-imidazolyl
639	Cl	2,6-diF-phenyl	5-methyl-1-imidazolyl
640	Cl	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
641	F	phenyl	2-(aminosulfonyl)phenyl
642	F	phenyl	2-(methyaminosulfonyl)phenyl
643	F	phenyl	1-pyrrolidinocarbonyl
644	F	phenyl	2-(methylsulfonyl)phenyl
645	F	phenyl	4-morpholino
646	F	phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
647	F	phenyl	4-morpholinocarbonyl
648	F	phenyl	2-methyl-1-imidazolyl
649	F	phenyl	5-methyl-1-imidazolyl
650	F	phenyl	2-methylsulfonyl-1-imidazolyl
651	F	2-pyridyl	2-(aminosulfonyl)phenyl
652	F	2-pyridyl	2-(methyaminosulfonyl)phenyl
653	F	2-pyridyl	1-pyrrolidinocarbonyl
654	F	2-pyridyl	2-(methylsulfonyl)phenyl
655	F	2-pyridyl	4-morpholino
656	F	2-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
657	F	2-pyridyl	4-morpholinocarbonyl
658	F	2-pyridyl	2-methyl-1-imidazolyl
659	F	2-pyridyl	5-methyl-1-imidazolyl
660	F	2-pyridyl	2-methylsulfonyl-1-imidazolyl
661	F	3-pyridyl	2-(aminosulfonyl)phenyl
662	F	3-pyridyl	2-(methyaminosulfonyl)phenyl
663	F	3-pyridyl	1-pyrrolidinocarbonyl
664	F	3-pyridyl	2-(methylsulfonyl)phenyl
665	F	3-pyridyl	4-morpholino
666	F	3-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
667	F	3-pyridyl	4-morpholinocarbonyl
668	F	3-pyridyl	2-methyl-1-imidazolyl
669	F	3-pyridyl	5-methyl-1-imidazolyl
670	F	3-pyridyl	2-methylsulfonyl-1-imidazolyl
671	F	2-pyrimidyl	2-(aminosulfonyl)phenyl
672	F	2-pyrimidyl	2-(methyaminosulfonyl)phenyl
673	F	2-pyrimidyl	1-pyrrolidinocarbonyl
674	F	2-pyrimidyl	2-(methylsulfonyl)phenyl
675	F	2-pyrimidyl	4-morpholino
676	F	2-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
677	F	2-pyrimidyl	4-morpholinocarbonyl
678	F	2-pyrimidyl	2-methyl-1-imidazolyl
679	F	2-pyrimidyl	5-methyl-1-imidazolyl
680	F	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
681	F	5-pyrimidyl	2-(aminosulfonyl)phenyl
682	F	5-pyrimidyl	2-(methyaminosulfonyl)phenyl

683	F	5-pyrimidyl	1-pyrrolidinocarbonyl
684	F	5-pyrimidyl	2-(methanesulfonyl)phenyl
685	F	5-pyrimidyl	4-morpholino
686	F	5-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
687	F	5-pyrimidyl	4-morpholinocarbonyl
688	F	5-pyrimidyl	2-methyl-1-imidazolyl
689	F	5-pyrimidyl	5-methyl-1-imidazolyl
690	F	5-pyrimidyl	2-methanesulfonyl-1-imidazolyl
691	F	2-Cl-phenyl	2-(aminosulfonyl)phenyl
692	F	2-Cl-phenyl	2-(methanaminosulfonyl)phenyl
693	F	2-Cl-phenyl	1-pyrrolidinocarbonyl
694	F	2-Cl-phenyl	2-(methanesulfonyl)phenyl
695	F	2-Cl-phenyl	4-morpholino
696	F	2-Cl-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
697	F	2-Cl-phenyl	4-morpholinocarbonyl
698	F	2-Cl-phenyl	2-methyl-1-imidazolyl
699	F	2-Cl-phenyl	5-methyl-1-imidazolyl
700	F	2-Cl-phenyl	2-methanesulfonyl-1-imidazolyl
701	F	2-F-phenyl	2-(aminosulfonyl)phenyl
702	F	2-F-phenyl	2-(methanaminosulfonyl)phenyl
703	F	2-F-phenyl	1-pyrrolidinocarbonyl
704	F	2-F-phenyl	2-(methanesulfonyl)phenyl
705	F	2-F-phenyl	4-morpholino
706	F	2-F-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
707	F	2-F-phenyl	4-morpholinocarbonyl
708	F	2-F-phenyl	2-methyl-1-imidazolyl
709	F	2-F-phenyl	5-methyl-1-imidazolyl
710	F	2-F-phenyl	2-methanesulfonyl-1-imidazolyl
711	F	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
712	F	2,6-diF-phenyl	2-(methanaminosulfonyl)phenyl
713	F	2,6-diF-phenyl	1-pyrrolidinocarbonyl
714	F	2,6-diF-phenyl	2-(methanesulfonyl)phenyl
715	F	2,6-diF-phenyl	4-morpholino
716	F	2,6-diF-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
717	F	2,6-diF-phenyl	4-morpholinocarbonyl
718	F	2,6-diF-phenyl	2-methyl-1-imidazolyl
719	F	2,6-diF-phenyl	5-methyl-1-imidazolyl
720	F	2,6-diF-phenyl	2-methanesulfonyl-1-imidazolyl
721	CO ₂ CH ₃	phenyl	2-(aminosulfonyl)phenyl
722	CO ₂ CH ₃	phenyl	2-(methanaminosulfonyl)phenyl
723	CO ₂ CH ₃	phenyl	1-pyrrolidinocarbonyl
724	CO ₂ CH ₃	phenyl	2-(methanesulfonyl)phenyl
725	CO ₂ CH ₃	phenyl	4-morpholino
726	CO ₂ CH ₃	phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
727	CO ₂ CH ₃	phenyl	4-morpholinocarbonyl
728	CO ₂ CH ₃	phenyl	2-methyl-1-imidazolyl
729	CO ₂ CH ₃	phenyl	5-methyl-1-imidazolyl
730	CO ₂ CH ₃	phenyl	2-methanesulfonyl-1-imidazolyl
731	CO ₂ CH ₃	2-pyridyl	2-(aminosulfonyl)phenyl
732	CO ₂ CH ₃	2-pyridyl	2-(methanaminosulfonyl)phenyl
733	CO ₂ CH ₃	2-pyridyl	1-pyrrolidinocarbonyl
734	CO ₂ CH ₃	2-pyridyl	2-(methanesulfonyl)phenyl
735	CO ₂ CH ₃	2-pyridyl	4-morpholino

736	CO ₂ CH ₃	2-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
737	CO ₂ CH ₃	2-pyridyl	4-morpholinocarbonyl
738	CO ₂ CH ₃	2-pyridyl	2-methyl-1-imidazolyl
739	CO ₂ CH ₃	2-pyridyl	5-methyl-1-imidazolyl
740	CO ₂ CH ₃	2-pyridyl	2-methylsulfonyl-1-imidazolyl
741	CO ₂ CH ₃	3-pyridyl	2-(aminosulfonyl)phenyl
742	CO ₂ CH ₃	3-pyridyl	2-(methylaminosulfonyl)phenyl
743	CO ₂ CH ₃	3-pyridyl	1-pyrrolidinocarbonyl
744	CO ₂ CH ₃	3-pyridyl	2-(methylsulfonyl)phenyl
745	CO ₂ CH ₃	3-pyridyl	4-morpholino
746	CO ₂ CH ₃	3-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
747	CO ₂ CH ₃	3-pyridyl	4-morpholinocarbonyl
748	CO ₂ CH ₃	3-pyridyl	2-methyl-1-imidazolyl
749	CO ₂ CH ₃	3-pyridyl	5-methyl-1-imidazolyl
750	CO ₂ CH ₃	3-pyridyl	2-methylsulfonyl-1-imidazolyl
751	CO ₂ CH ₃	2-pyrimidyl	2-(aminosulfonyl)phenyl
752	CO ₂ CH ₃	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
753	CO ₂ CH ₃	2-pyrimidyl	1-pyrrolidinocarbonyl
754	CO ₂ CH ₃	2-pyrimidyl	2-(methylsulfonyl)phenyl
755	CO ₂ CH ₃	2-pyrimidyl	4-morpholino
756	CO ₂ CH ₃	2-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
757	CO ₂ CH ₃	2-pyrimidyl	4-morpholinocarbonyl
758	CO ₂ CH ₃	2-pyrimidyl	2-methyl-1-imidazolyl
759	CO ₂ CH ₃	2-pyrimidyl	5-methyl-1-imidazolyl
760	CO ₂ CH ₃	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
761	CO ₂ CH ₃	5-pyrimidyl	2-(aminosulfonyl)phenyl
762	CO ₂ CH ₃	5-pyrimidyl	2-(methylaminosulfonyl)phenyl
763	CO ₂ CH ₃	5-pyrimidyl	1-pyrrolidinocarbonyl
764	CO ₂ CH ₃	5-pyrimidyl	2-(methylsulfonyl)phenyl
765	CO ₂ CH ₃	5-pyrimidyl	4-morpholino
766	CO ₂ CH ₃	5-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
767	CO ₂ CH ₃	5-pyrimidyl	4-morpholinocarbonyl
768	CO ₂ CH ₃	5-pyrimidyl	2-methyl-1-imidazolyl
769	CO ₂ CH ₃	5-pyrimidyl	5-methyl-1-imidazolyl
770	CO ₂ CH ₃	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
771	CO ₂ CH ₃	2-Cl-phenyl	2-(aminosulfonyl)phenyl
772	CO ₂ CH ₃	2-Cl-phenyl	2-(methylaminosulfonyl)phenyl
773	CO ₂ CH ₃	2-Cl-phenyl	1-pyrrolidinocarbonyl
774	CO ₂ CH ₃	2-Cl-phenyl	2-(methylsulfonyl)phenyl
775	CO ₂ CH ₃	2-Cl-phenyl	4-morpholino
776	CO ₂ CH ₃	2-Cl-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
777	CO ₂ CH ₃	2-Cl-phenyl	4-morpholinocarbonyl
778	CO ₂ CH ₃	2-Cl-phenyl	2-methyl-1-imidazolyl
779	CO ₂ CH ₃	2-Cl-phenyl	5-methyl-1-imidazolyl
780	CO ₂ CH ₃	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
781	CO ₂ CH ₃	2-F-phenyl	2-(aminosulfonyl)phenyl
782	CO ₂ CH ₃	2-F-phenyl	2-(methylaminosulfonyl)phenyl
783	CO ₂ CH ₃	2-F-phenyl	1-pyrrolidinocarbonyl


784	CO ₂ CH ₃	2-F-phenyl	2-(methylsulfonyl)phenyl
785	CO ₂ CH ₃	2-F-phenyl	4-morpholino
786	CO ₂ CH ₃	2-F-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
787	CO ₂ CH ₃	2-F-phenyl	4-morpholinocarbonyl
788	CO ₂ CH ₃	2-F-phenyl	2-methyl-1-imidazolyl
789	CO ₂ CH ₃	2-F-phenyl	5-methyl-1-imidazolyl
790	CO ₂ CH ₃	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
791	CO ₂ CH ₃	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
792	CO ₂ CH ₃	2,6-diF-phenyl	2-(methylaminosulfonyl)phenyl
793	CO ₂ CH ₃	2,6-diF-phenyl	1-pyrrolidinocarbonyl
794	CO ₂ CH ₃	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
795	CO ₂ CH ₃	2,6-diF-phenyl	4-morpholino
796	CO ₂ CH ₃	2,6-diF-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
797	CO ₂ CH ₃	2,6-diF-phenyl	4-morpholinocarbonyl
798	CO ₂ CH ₃	2,6-diF-phenyl	2-methyl-1-imidazolyl
799	CO ₂ CH ₃	2,6-diF-phenyl	5-methyl-1-imidazolyl
800	CO ₂ CH ₃	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
801	CH ₂ OCH ₃	phenyl	2-(aminosulfonyl)phenyl
802	CH ₂ OCH ₃	phenyl	2-(methylaminosulfonyl)phenyl
803	CH ₂ OCH ₃	phenyl	1-pyrrolidinocarbonyl
804	CH ₂ OCH ₃	phenyl	2-(methylsulfonyl)phenyl
805	CH ₂ OCH ₃	phenyl	4-morpholino
806	CH ₂ OCH ₃	phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
807	CH ₂ OCH ₃	phenyl	4-morpholinocarbonyl
808	CH ₂ OCH ₃	phenyl	2-methyl-1-imidazolyl
809	CH ₂ OCH ₃	phenyl	5-methyl-1-imidazolyl
810	CH ₂ OCH ₃	phenyl	2-methylsulfonyl-1-imidazolyl
811	CH ₂ OCH ₃	2-pyridyl	2-(aminosulfonyl)phenyl
812	CH ₂ OCH ₃	2-pyridyl	2-(methylaminosulfonyl)phenyl
813	CH ₂ OCH ₃	2-pyridyl	1-pyrrolidinocarbonyl
814	CH ₂ OCH ₃	2-pyridyl	2-(methylsulfonyl)phenyl
815	CH ₂ OCH ₃	2-pyridyl	4-morpholino
816	CH ₂ OCH ₃	2-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
817	CH ₂ OCH ₃	2-pyridyl	4-morpholinocarbonyl
818	CH ₂ OCH ₃	2-pyridyl	2-methyl-1-imidazolyl
819	CH ₂ OCH ₃	2-pyridyl	5-methyl-1-imidazolyl
820	CH ₂ OCH ₃	2-pyridyl	2-methylsulfonyl-1-imidazolyl
821	CH ₂ OCH ₃	3-pyridyl	2-(aminosulfonyl)phenyl
822	CH ₂ OCH ₃	3-pyridyl	2-(methylaminosulfonyl)phenyl
823	CH ₂ OCH ₃	3-pyridyl	1-pyrrolidinocarbonyl
824	CH ₂ OCH ₃	3-pyridyl	2-(methylsulfonyl)phenyl
825	CH ₂ OCH ₃	3-pyridyl	4-morpholino
826	CH ₂ OCH ₃	3-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
827	CH ₂ OCH ₃	3-pyridyl	4-morpholinocarbonyl
828	CH ₂ OCH ₃	3-pyridyl	2-methyl-1-imidazolyl
829	CH ₂ OCH ₃	3-pyridyl	5-methyl-1-imidazolyl
830	CH ₂ OCH ₃	3-pyridyl	2-methylsulfonyl-1-imidazolyl
831	CH ₂ OCH ₃	2-pyrimidyl	2-(aminosulfonyl)phenyl


832	CH ₂ OCH ₃	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
833	CH ₂ OCH ₃	2-pyrimidyl	1-pyrrolidinocarbonyl
834	CH ₂ OCH ₃	2-pyrimidyl	2-(methylsulfonyl)phenyl
835	CH ₂ OCH ₃	2-pyrimidyl	4-morpholino
836	CH ₂ OCH ₃	2-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
837	CH ₂ OCH ₃	2-pyrimidyl	4-morpholinocarbonyl
838	CH ₂ OCH ₃	2-pyrimidyl	2-methyl-1-imidazolyl
839	CH ₂ OCH ₃	2-pyrimidyl	5-methyl-1-imidazolyl
840	CH ₂ OCH ₃	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
841	CH ₂ OCH ₃	5-pyrimidyl	2-(aminosulfonyl)phenyl
842	CH ₂ OCH ₃	5-pyrimidyl	2-(methylaminosulfonyl)phenyl
843	CH ₂ OCH ₃	5-pyrimidyl	1-pyrrolidinocarbonyl
844	CH ₂ OCH ₃	5-pyrimidyl	2-(methylsulfonyl)phenyl
845	CH ₂ OCH ₃	5-pyrimidyl	4-morpholino
846	CH ₂ OCH ₃	5-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
847	CH ₂ OCH ₃	5-pyrimidyl	4-morpholinocarbonyl
848	CH ₂ OCH ₃	5-pyrimidyl	2-methyl-1-imidazolyl
849	CH ₂ OCH ₃	5-pyrimidyl	5-methyl-1-imidazolyl
850	CH ₂ OCH ₃	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
851	CH ₂ OCH ₃	2-Cl-phenyl	2-(aminosulfonyl)phenyl
852	CH ₂ OCH ₃	2-Cl-phenyl	2-(methylaminosulfonyl)phenyl
853	CH ₂ OCH ₃	2-Cl-phenyl	1-pyrrolidinocarbonyl
854	CH ₂ OCH ₃	2-Cl-phenyl	2-(methylsulfonyl)phenyl
855	CH ₂ OCH ₃	2-Cl-phenyl	4-morpholino
856	CH ₂ OCH ₃	2-Cl-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
857	CH ₂ OCH ₃	2-Cl-phenyl	4-morpholinocarbonyl
858	CH ₂ OCH ₃	2-Cl-phenyl	2-methyl-1-imidazolyl
859	CH ₂ OCH ₃	2-Cl-phenyl	5-methyl-1-imidazolyl
860	CH ₂ OCH ₃	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
861	CH ₂ OCH ₃	2-F-phenyl	2-(aminosulfonyl)phenyl
862	CH ₂ OCH ₃	2-F-phenyl	2-(methylaminosulfonyl)phenyl
863	CH ₂ OCH ₃	2-F-phenyl	1-pyrrolidinocarbonyl
864	CH ₂ OCH ₃	2-F-phenyl	2-(methylsulfonyl)phenyl
865	CH ₂ OCH ₃	2-F-phenyl	4-morpholino
866	CH ₂ OCH ₃	2-F-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
867	CH ₂ OCH ₃	2-F-phenyl	4-morpholinocarbonyl
868	CH ₂ OCH ₃	2-F-phenyl	2-methyl-1-imidazolyl
869	CH ₂ OCH ₃	2-F-phenyl	5-methyl-1-imidazolyl
870	CH ₂ OCH ₃	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
871	CH ₂ OCH ₃	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
872	CH ₂ OCH ₃	2,6-diF-phenyl	2-(methylaminosulfonyl)phenyl
873	CH ₂ OCH ₃	2,6-diF-phenyl	1-pyrrolidinocarbonyl
874	CH ₂ OCH ₃	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
875	CH ₂ OCH ₃	2,6-diF-phenyl	4-morpholino
876	CH ₂ OCH ₃	2,6-diF-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
877	CH ₂ OCH ₃	2,6-diF-phenyl	4-morpholinocarbonyl
878	CH ₂ OCH ₃	2,6-diF-phenyl	2-methyl-1-imidazolyl
879	CH ₂ OCH ₃	2,6-diF-phenyl	5-methyl-1-imidazolyl


880	CH ₂ OCH ₃	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
881	CONH ₂	phenyl	2-(aminosulfonyl)phenyl
882	CONH ₂	phenyl	2-(methylaminosulfonyl)phenyl
883	CONH ₂	phenyl	1-pyrrolidinocarbonyl
884	CONH ₂	phenyl	2-(methylsulfonyl)phenyl
885	CONH ₂	phenyl	4-morpholino
886	CONH ₂	phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
887	CONH ₂	phenyl	4-morpholinocarbonyl
888	CONH ₂	phenyl	2-methyl-1-imidazolyl
889	CONH ₂	phenyl	5-methyl-1-imidazolyl
890	CONH ₂	phenyl	2-methylsulfonyl-1-imidazolyl
891	CONH ₂	2-pyridyl	2-(aminosulfonyl)phenyl
892	CONH ₂	2-pyridyl	2-(methylaminosulfonyl)phenyl
893	CONH ₂	2-pyridyl	1-pyrrolidinocarbonyl
894	CONH ₂	2-pyridyl	2-(methylsulfonyl)phenyl
895	CONH ₂	2-pyridyl	4-morpholino
896	CONH ₂	2-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
897	CONH ₂	2-pyridyl	4-morpholinocarbonyl
898	CONH ₂	2-pyridyl	2-methyl-1-imidazolyl
899	CONH ₂	2-pyridyl	5-methyl-1-imidazolyl
900	CONH ₂	2-pyridyl	2-methylsulfonyl-1-imidazolyl
901	CONH ₂	3-pyridyl	2-(aminosulfonyl)phenyl
902	CONH ₂	3-pyridyl	2-(methylaminosulfonyl)phenyl
903	CONH ₂	3-pyridyl	1-pyrrolidinocarbonyl
904	CONH ₂	3-pyridyl	2-(methylsulfonyl)phenyl
905	CONH ₂	3-pyridyl	4-morpholino
906	CONH ₂	3-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
907	CONH ₂	3-pyridyl	4-morpholinocarbonyl
908	CONH ₂	3-pyridyl	2-methyl-1-imidazolyl
909	CONH ₂	3-pyridyl	5-methyl-1-imidazolyl
910	CONH ₂	3-pyridyl	2-methylsulfonyl-1-imidazolyl
911	CONH ₂	2-pyrimidyl	2-(aminosulfonyl)phenyl
912	CONH ₂	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
913	CONH ₂	2-pyrimidyl	1-pyrrolidinocarbonyl
914	CONH ₂	2-pyrimidyl	2-(methylsulfonyl)phenyl
915	CONH ₂	2-pyrimidyl	4-morpholino
916	CONH ₂	2-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
917	CONH ₂	2-pyrimidyl	4-morpholinocarbonyl
918	CONH ₂	2-pyrimidyl	2-methyl-1-imidazolyl
919	CONH ₂	2-pyrimidyl	5-methyl-1-imidazolyl
920	CONH ₂	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
921	CONH ₂	5-pyrimidyl	2-(aminosulfonyl)phenyl
922	CONH ₂	5-pyrimidyl	2-(methylaminosulfonyl)phenyl
923	CONH ₂	5-pyrimidyl	1-pyrrolidinocarbonyl
924	CONH ₂	5-pyrimidyl	2-(methylsulfonyl)phenyl
925	CONH ₂	5-pyrimidyl	4-morpholino
926	CONH ₂	5-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
927	CONH ₂	5-pyrimidyl	4-morpholinocarbonyl




928	CONH ₂	5-pyrimidyl	2-methyl-1-imidazolyl
929	CONH ₂	5-pyrimidyl	5-methyl-1-imidazolyl
930	CONH ₂	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
931	CONH ₂	2-Cl-phenyl	2-(aminosulfonyl)phenyl
932	CONH ₂	2-Cl-phenyl	2-(methylaninosulfonyl)phenyl
933	CONH ₂	2-Cl-phenyl	1-pyrrolidinocarbonyl
934	CONH ₂	2-Cl-phenyl	2-(methylsulfonyl)phenyl
935	CONH ₂	2-Cl-phenyl	4-morpholino
936	CONH ₂	2-Cl-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
937	CONH ₂	2-Cl-phenyl	4-morpholinocarbonyl
938	CONH ₂	2-Cl-phenyl	2-methyl-1-imidazolyl
939	CONH ₂	2-Cl-phenyl	5-methyl-1-imidazolyl
940	CONH ₂	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
941	CONH ₂	2-F-phenyl	2-(aminosulfonyl)phenyl
942	CONH ₂	2-F-phenyl	2-(methylaninosulfonyl)phenyl
943	CONH ₂	2-F-phenyl	1-pyrrolidinocarbonyl
944	CONH ₂	2-F-phenyl	2-(methylsulfonyl)phenyl
945	CONH ₂	2-F-phenyl	4-morpholino
946	CONH ₂	2-F-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
947	CONH ₂	2-F-phenyl	4-morpholinocarbonyl
948	CONH ₂	2-F-phenyl	2-methyl-1-imidazolyl
949	CONH ₂	2-F-phenyl	5-methyl-1-imidazolyl
950	CONH ₂	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
951	CONH ₂	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
952	CONH ₂	2,6-diF-phenyl	2-(methylaninosulfonyl)phenyl
953	CONH ₂	2,6-diF-phenyl	1-pyrrolidinocarbonyl
954	CONH ₂	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
955	CONH ₂	2,6-diF-phenyl	4-morpholino
956	CONH ₂	2,6-diF-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
957	CONH ₂	2,6-diF-phenyl	4-morpholinocarbonyl
958	CONH ₂	2,6-diF-phenyl	2-methyl-1-imidazolyl
959	CONH ₂	2,6-diF-phenyl	5-methyl-1-imidazolyl
960	CONH ₂	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl


Table 7














oo




pp




qq




rr




ss



tt



uu



vv

Ex #	R ^{1a}	A	B
1	CH ₃	phenyl	2-(aminosulfonyl)phenyl
2	CH ₃	phenyl	2-(methylaninosulfonyl)phenyl
3	CH ₃	phenyl	1-pyrrolidinocarbonyl
4	CH ₃	phenyl	2-(methylsulfonyl)phenyl
5	CH ₃	phenyl	4-morpholino
6	CH ₃	phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
7	CH ₃	phenyl	4-morpholinocarbonyl
8	CH ₃	phenyl	2-methyl-1-imidazolyl
9	CH ₃	phenyl	5-methyl-1-imidazolyl
10	CH ₃	phenyl	2-methylsulfonyl-1-imidazolyl
11	CH ₃	2-pyridyl	2-(aminosulfonyl)phenyl
12	CH ₃	2-pyridyl	2-(methylaninosulfonyl)phenyl
13	CH ₃	2-pyridyl	1-pyrrolidinocarbonyl
14	CH ₃	2-pyridyl	2-(methylsulfonyl)phenyl
15	CH ₃	2-pyridyl	4-morpholino
16	CH ₃	2-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
17	CH ₃	2-pyridyl	4-morpholinocarbonyl
18	CH ₃	2-pyridyl	2-methyl-1-imidazolyl
19	CH ₃	2-pyridyl	5-methyl-1-imidazolyl
20	CH ₃	2-pyridyl	2-methylsulfonyl-1-imidazolyl
21	CH ₃	3-pyridyl	2-(aminosulfonyl)phenyl
22	CH ₃	3-pyridyl	2-(methylaninosulfonyl)phenyl
23	CH ₃	3-pyridyl	1-pyrrolidinocarbonyl
24	CH ₃	3-pyridyl	2-(methylsulfonyl)phenyl
25	CH ₃	3-pyridyl	4-morpholino

26	CH ₃	3-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
27	CH ₃	3-pyridyl	4-morpholinocarbonyl
28	CH ₃	3-pyridyl	2-methyl-1-imidazolyl
29	CH ₃	3-pyridyl	5-methyl-1-imidazolyl
30	CH ₃	3-pyridyl	2-methylsulfonyl-1-imidazolyl
31	CH ₃	2-pyrimidyl	2-(aminosulfonyl)phenyl
32	CH ₃	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
33	CH ₃	2-pyrimidyl	1-pyrrolidinocarbonyl
34	CH ₃	2-pyrimidyl	2-(methylsulfonyl)phenyl
35	CH ₃	2-pyrimidyl	4-morpholino
36	CH ₃	2-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
37	CH ₃	2-pyrimidyl	4-morpholinocarbonyl
38	CH ₃	2-pyrimidyl	2-methyl-1-imidazolyl
39	CH ₃	2-pyrimidyl	5-methyl-1-imidazolyl
40	CH ₃	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
41	CH ₃	5-pyrimidyl	2-(aminosulfonyl)phenyl
42	CH ₃	5-pyrimidyl	2-(methylaminosulfonyl)phenyl
43	CH ₃	5-pyrimidyl	1-pyrrolidinocarbonyl
44	CH ₃	5-pyrimidyl	2-(methylsulfonyl)phenyl
45	CH ₃	5-pyrimidyl	4-morpholino
46	CH ₃	5-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
47	CH ₃	5-pyrimidyl	4-morpholinocarbonyl
48	CH ₃	5-pyrimidyl	2-methyl-1-imidazolyl
49	CH ₃	5-pyrimidyl	5-methyl-1-imidazolyl
50	CH ₃	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
51	CH ₃	2-Cl-phenyl	2-(aminosulfonyl)phenyl
52	CH ₃	2-Cl-phenyl	2-(methylaminosulfonyl)phenyl
53	CH ₃	2-Cl-phenyl	1-pyrrolidinocarbonyl
54	CH ₃	2-Cl-phenyl	2-(methylsulfonyl)phenyl
55	CH ₃	2-Cl-phenyl	4-morpholino
56	CH ₃	2-Cl-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
57	CH ₃	2-Cl-phenyl	4-morpholinocarbonyl
58	CH ₃	2-Cl-phenyl	2-methyl-1-imidazolyl
59	CH ₃	2-Cl-phenyl	5-methyl-1-imidazolyl
60	CH ₃	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
61	CH ₃	2-F-phenyl	2-(aminosulfonyl)phenyl
62	CH ₃	2-F-phenyl	2-(methylaminosulfonyl)phenyl
63	CH ₃	2-F-phenyl	1-pyrrolidinocarbonyl
64	CH ₃	2-F-phenyl	2-(methylsulfonyl)phenyl
65	CH ₃	2-F-phenyl	4-morpholino
66	CH ₃	2-F-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
67	CH ₃	2-F-phenyl	4-morpholinocarbonyl
68	CH ₃	2-F-phenyl	2-methyl-1-imidazolyl
69	CH ₃	2-F-phenyl	5-methyl-1-imidazolyl
70	CH ₃	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
71	CH ₃	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
72	CH ₃	2,6-diF-phenyl	2-(methylaminosulfonyl)phenyl
73	CH ₃	2,6-diF-phenyl	1-pyrrolidinocarbonyl

74	CH ₃	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
75	CH ₃	2,6-diF-phenyl	4-morpholino
76	CH ₃	2,6-diF-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
77	CH ₃	2,6-diF-phenyl	4-morpholinocarbonyl
78	CH ₃	2,6-diF-phenyl	2-methyl-1-imidazolyl
79	CH ₃	2,6-diF-phenyl	5-methyl-1-imidazolyl
80	CH ₃	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
81	CH ₂ CH ₃	phenyl	2-(aminosulfonyl)phenyl
82	CH ₂ CH ₃	phenyl	2-(methylaminosulfonyl)phenyl
83	CH ₂ CH ₃	phenyl	1-pyrrolidinocarbonyl
84	CH ₂ CH ₃	phenyl	2-(methylsulfonyl)phenyl
85	CH ₂ CH ₃	phenyl	4-morpholino
86	CH ₂ CH ₃	phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
87	CH ₂ CH ₃	phenyl	4-morpholinocarbonyl
88	CH ₂ CH ₃	phenyl	2-methyl-1-imidazolyl
89	CH ₂ CH ₃	phenyl	5-methyl-1-imidazolyl
90	CH ₂ CH ₃	phenyl	2-methylsulfonyl-1-imidazolyl
91	CH ₂ CH ₃	2-pyridyl	2-(aminosulfonyl)phenyl
92	CH ₂ CH ₃	2-pyridyl	2-(methylaminosulfonyl)phenyl
93	CH ₂ CH ₃	2-pyridyl	1-pyrrolidinocarbonyl
94	CH ₂ CH ₃	2-pyridyl	2-(methylsulfonyl)phenyl
95	CH ₂ CH ₃	2-pyridyl	4-morpholino
96	CH ₂ CH ₃	2-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
97	CH ₂ CH ₃	2-pyridyl	4-morpholinocarbonyl
98	CH ₂ CH ₃	2-pyridyl	2-methyl-1-imidazolyl
99	CH ₂ CH ₃	2-pyridyl	5-methyl-1-imidazolyl
100	CH ₂ CH ₃	2-pyridyl	2-methylsulfonyl-1-imidazolyl
101	CH ₂ CH ₃	3-pyridyl	2-(aminosulfonyl)phenyl
102	CH ₂ CH ₃	3-pyridyl	2-(methylaminosulfonyl)phenyl
103	CH ₂ CH ₃	3-pyridyl	1-pyrrolidinocarbonyl
104	CH ₂ CH ₃	3-pyridyl	2-(methylsulfonyl)phenyl
105	CH ₂ CH ₃	3-pyridyl	4-morpholino
106	CH ₂ CH ₃	3-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
107	CH ₂ CH ₃	3-pyridyl	4-morpholinocarbonyl
108	CH ₂ CH ₃	3-pyridyl	2-methyl-1-imidazolyl
109	CH ₂ CH ₃	3-pyridyl	5-methyl-1-imidazolyl
110	CH ₂ CH ₃	3-pyridyl	2-methylsulfonyl-1-imidazolyl
111	CH ₂ CH ₃	2-pyrimidyl	2-(aminosulfonyl)phenyl
112	CH ₂ CH ₃	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
113	CH ₂ CH ₃	2-pyrimidyl	1-pyrrolidinocarbonyl
114	CH ₂ CH ₃	2-pyrimidyl	2-(methylsulfonyl)phenyl
115	CH ₂ CH ₃	2-pyrimidyl	4-morpholino
116	CH ₂ CH ₃	2-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
117	CH ₂ CH ₃	2-pyrimidyl	4-morpholinocarbonyl
118	CH ₂ CH ₃	2-pyrimidyl	2-methyl-1-imidazolyl
119	CH ₂ CH ₃	2-pyrimidyl	5-methyl-1-imidazolyl
120	CH ₂ CH ₃	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
121	CH ₂ CH ₃	5-pyrimidyl	2-(aminosulfonyl)phenyl

122	CH ₂ CH ₃	5-pyrimidyl	2-(methylaninosulfonyl)phenyl
123	CH ₂ CH ₃	5-pyrimidyl	1-pyrrolidinocarbonyl
124	CH ₂ CH ₃	5-pyrimidyl	2-(methylsulfonyl)phenyl
125	CH ₂ CH ₃	5-pyrimidyl	4-morpholino
126	CH ₂ CH ₃	5-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
127	CH ₂ CH ₃	5-pyrimidyl	4-morpholinocarbonyl
128	CH ₂ CH ₃	5-pyrimidyl	2-methyl-1-imidazolyl
129	CH ₂ CH ₃	5-pyrimidyl	5-methyl-1-imidazolyl
130	CH ₂ CH ₃	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
131	CH ₂ CH ₃	2-Cl-phenyl	2-(aminosulfonyl)phenyl
132	CH ₂ CH ₃	2-Cl-phenyl	2-(methylaninosulfonyl)phenyl
133	CH ₂ CH ₃	2-Cl-phenyl	1-pyrrolidinocarbonyl
134	CH ₂ CH ₃	2-Cl-phenyl	2-(methylsulfonyl)phenyl
135	CH ₂ CH ₃	2-Cl-phenyl	4-morpholino
136	CH ₂ CH ₃	2-Cl-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
137	CH ₂ CH ₃	2-Cl-phenyl	4-morpholinocarbonyl
138	CH ₂ CH ₃	2-Cl-phenyl	2-methyl-1-imidazolyl
139	CH ₂ CH ₃	2-Cl-phenyl	5-methyl-1-imidazolyl
140	CH ₂ CH ₃	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
141	CH ₂ CH ₃	2-F-phenyl	2-(aminosulfonyl)phenyl
142	CH ₂ CH ₃	2-F-phenyl	2-(methylaninosulfonyl)phenyl
143	CH ₂ CH ₃	2-F-phenyl	1-pyrrolidinocarbonyl
144	CH ₂ CH ₃	2-F-phenyl	2-(methylsulfonyl)phenyl
145	CH ₂ CH ₃	2-F-phenyl	4-morpholino
146	CH ₂ CH ₃	2-F-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
147	CH ₂ CH ₃	2-F-phenyl	4-morpholinocarbonyl
148	CH ₂ CH ₃	2-F-phenyl	2-methyl-1-imidazolyl
149	CH ₂ CH ₃	2-F-phenyl	5-methyl-1-imidazolyl
150	CH ₂ CH ₃	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
151	CH ₂ CH ₃	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
152	CH ₂ CH ₃	2,6-diF-phenyl	2-(methylaninosulfonyl)phenyl
153	CH ₂ CH ₃	2,6-diF-phenyl	1-pyrrolidinocarbonyl
154	CH ₂ CH ₃	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
155	CH ₂ CH ₃	2,6-diF-phenyl	4-morpholino
156	CH ₂ CH ₃	2,6-diF-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
157	CH ₂ CH ₃	2,6-diF-phenyl	4-morpholinocarbonyl
158	CH ₂ CH ₃	2,6-diF-phenyl	2-methyl-1-imidazolyl
159	CH ₂ CH ₃	2,6-diF-phenyl	5-methyl-1-imidazolyl
160	CH ₂ CH ₃	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
161	CF ₃	phenyl	2-(aminosulfonyl)phenyl
162	CF ₃	phenyl	2-(methylaninosulfonyl)phenyl
163	CF ₃	phenyl	1-pyrrolidinocarbonyl
164	CF ₃	phenyl	2-(methylsulfonyl)phenyl
165	CF ₃	phenyl	4-morpholino
166	CF ₃	phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
167	CF ₃	phenyl	4-morpholinocarbonyl
168	CF ₃	phenyl	2-methyl-1-imidazolyl
169	CF ₃	phenyl	5-methyl-1-imidazolyl

170	CF ₃	phenyl	2-methylsulfonyl-1-imidazolyl
171	CF ₃	2-pyridyl	2-(aminosulfonyl)phenyl
172	CF ₃	2-pyridyl	2-(methylaminosulfonyl)phenyl
173	CF ₃	2-pyridyl	1-pyrrolidinocarbonyl
174	CF ₃	2-pyridyl	2-(methylsulfonyl)phenyl
175	CF ₃	2-pyridyl	4-morpholino
176	CF ₃	2-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
177	CF ₃	2-pyridyl	4-morpholinocarbonyl
178	CF ₃	2-pyridyl	2-methyl-1-imidazolyl
179	CF ₃	2-pyridyl	5-methyl-1-imidazolyl
180	CF ₃	2-pyridyl	2-methylsulfonyl-1-imidazolyl
181	CF ₃	3-pyridyl	2-(aminosulfonyl)phenyl
182	CF ₃	3-pyridyl	2-(methylaminosulfonyl)phenyl
183	CF ₃	3-pyridyl	1-pyrrolidinocarbonyl
184	CF ₃	3-pyridyl	2-(methylsulfonyl)phenyl
185	CF ₃	3-pyridyl	4-morpholino
186	CF ₃	3-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
187	CF ₃	3-pyridyl	4-morpholinocarbonyl
188	CF ₃	3-pyridyl	2-methyl-1-imidazolyl
189	CF ₃	3-pyridyl	5-methyl-1-imidazolyl
190	CF ₃	3-pyridyl	2-methylsulfonyl-1-imidazolyl
191	CF ₃	2-pyrimidyl	2-(aminosulfonyl)phenyl
192	CF ₃	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
193	CF ₃	2-pyrimidyl	1-pyrrolidinocarbonyl
194	CF ₃	2-pyrimidyl	2-(methylsulfonyl)phenyl
195	CF ₃	2-pyrimidyl	4-morpholino
196	CF ₃	2-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
197	CF ₃	2-pyrimidyl	4-morpholinocarbonyl
198	CF ₃	2-pyrimidyl	2-methyl-1-imidazolyl
199	CF ₃	2-pyrimidyl	5-methyl-1-imidazolyl
200	CF ₃	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
201	CF ₃	5-pyrimidyl	2-(aminosulfonyl)phenyl
202	CF ₃	5-pyrimidyl	2-(methylaminosulfonyl)phenyl
203	CF ₃	5-pyrimidyl	1-pyrrolidinocarbonyl
204	CF ₃	5-pyrimidyl	2-(methylsulfonyl)phenyl
205	CF ₃	5-pyrimidyl	4-morpholino
206	CF ₃	5-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
207	CF ₃	5-pyrimidyl	4-morpholinocarbonyl
208	CF ₃	5-pyrimidyl	2-methyl-1-imidazolyl
209	CF ₃	5-pyrimidyl	5-methyl-1-imidazolyl
210	CF ₃	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
211	CF ₃	2-Cl-phenyl	2-(aminosulfonyl)phenyl
212	CF ₃	2-Cl-phenyl	2-(methylaminosulfonyl)phenyl
213	CF ₃	2-Cl-phenyl	1-pyrrolidinocarbonyl
214	CF ₃	2-Cl-phenyl	2-(methylsulfonyl)phenyl
215	CF ₃	2-Cl-phenyl	4-morpholino
216	CF ₃	2-Cl-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
217	CF ₃	2-Cl-phenyl	4-morpholinocarbonyl

218	CF ₃	2-Cl-phenyl	2-methyl-1-imidazolyl
219	CF ₃	2-Cl-phenyl	5-methyl-1-imidazolyl
220	CF ₃	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
221	CF ₃	2-F-phenyl	2-(aminosulfonyl)phenyl
222	CF ₃	2-F-phenyl	2-(methylaninosulfonyl)phenyl
223	CF ₃	2-F-phenyl	1-pyrrolidinocarbonyl
224	CF ₃	2-F-phenyl	2-(methylsulfonyl)phenyl
225	CF ₃	2-F-phenyl	4-morpholino
226	CF ₃	2-F-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
227	CF ₃	2-F-phenyl	4-morpholinocarbonyl
228	CF ₃	2-F-phenyl	2-methyl-1-imidazolyl
229	CF ₃	2-F-phenyl	5-methyl-1-imidazolyl
230	CF ₃	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
231	CF ₃	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
232	CF ₃	2,6-diF-phenyl	2-(methylaninosulfonyl)phenyl
233	CF ₃	2,6-diF-phenyl	1-pyrrolidinocarbonyl
234	CF ₃	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
235	CF ₃	2,6-diF-phenyl	4-morpholino
236	CF ₃	2,6-diF-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
237	CF ₃	2,6-diF-phenyl	4-morpholinocarbonyl
238	CF ₃	2,6-diF-phenyl	2-methyl-1-imidazolyl
239	CF ₃	2,6-diF-phenyl	5-methyl-1-imidazolyl
240	CF ₃	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
241	SCH ₃	phenyl	2-(aminosulfonyl)phenyl
242	SCH ₃	phenyl	2-(methylaninosulfonyl)phenyl
243	SCH ₃	phenyl	1-pyrrolidinocarbonyl
244	SCH ₃	phenyl	2-(methylsulfonyl)phenyl
245	SCH ₃	phenyl	4-morpholino
246	SCH ₃	phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
247	SCH ₃	phenyl	4-morpholinocarbonyl
248	SCH ₃	phenyl	2-methyl-1-imidazolyl
249	SCH ₃	phenyl	5-methyl-1-imidazolyl
250	SCH ₃	phenyl	2-methylsulfonyl-1-imidazolyl
251	SCH ₃	2-pyridyl	2-(aminosulfonyl)phenyl
252	SCH ₃	2-pyridyl	2-(methylaninosulfonyl)phenyl
253	SCH ₃	2-pyridyl	1-pyrrolidinocarbonyl
254	SCH ₃	2-pyridyl	2-(methylsulfonyl)phenyl
255	SCH ₃	2-pyridyl	4-morpholino
256	SCH ₃	2-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
257	SCH ₃	2-pyridyl	4-morpholinocarbonyl
258	SCH ₃	2-pyridyl	2-methyl-1-imidazolyl
259	SCH ₃	2-pyridyl	5-methyl-1-imidazolyl
260	SCH ₃	2-pyridyl	2-methylsulfonyl-1-imidazolyl
261	SCH ₃	3-pyridyl	2-(aminosulfonyl)phenyl
262	SCH ₃	3-pyridyl	2-(methylaninosulfonyl)phenyl
263	SCH ₃	3-pyridyl	1-pyrrolidinocarbonyl
264	SCH ₃	3-pyridyl	2-(methylsulfonyl)phenyl
265	SCH ₃	3-pyridyl	4-morpholino

266	SCH ₃	3-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
267	SCH ₃	3-pyridyl	4-morpholinocarbonyl
268	SCH ₃	3-pyridyl	2-methyl-1-imidazolyl
269	SCH ₃	3-pyridyl	5-methyl-1-imidazolyl
270	SCH ₃	3-pyridyl	2-methylsulfonyl-1-imidazolyl
271	SCH ₃	2-pyrimidyl	2-(aminosulfonyl)phenyl
272	SCH ₃	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
273	SCH ₃	2-pyrimidyl	1-pyrrolidinocarbonyl
274	SCH ₃	2-pyrimidyl	2-(methylsulfonyl)phenyl
275	SCH ₃	2-pyrimidyl	4-morpholino
276	SCH ₃	2-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
277	SCH ₃	2-pyrimidyl	4-morpholinocarbonyl
278	SCH ₃	2-pyrimidyl	2-methyl-1-imidazolyl
279	SCH ₃	2-pyrimidyl	5-methyl-1-imidazolyl
280	SCH ₃	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
281	SCH ₃	5-pyrimidyl	2-(aminosulfonyl)phenyl
282	SCH ₃	5-pyrimidyl	2-(methylaminosulfonyl)phenyl
283	SCH ₃	5-pyrimidyl	1-pyrrolidinocarbonyl
284	SCH ₃	5-pyrimidyl	2-(methylsulfonyl)phenyl
285	SCH ₃	5-pyrimidyl	4-morpholino
286	SCH ₃	5-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
287	SCH ₃	5-pyrimidyl	4-morpholinocarbonyl
288	SCH ₃	5-pyrimidyl	2-methyl-1-imidazolyl
289	SCH ₃	5-pyrimidyl	5-methyl-1-imidazolyl
290	SCH ₃	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
291	SCH ₃	2-Cl-phenyl	2-(aminosulfonyl)phenyl
292	SCH ₃	2-Cl-phenyl	2-(methylaminosulfonyl)phenyl
293	SCH ₃	2-Cl-phenyl	1-pyrrolidinocarbonyl
294	SCH ₃	2-Cl-phenyl	2-(methylsulfonyl)phenyl
295	SCH ₃	2-Cl-phenyl	4-morpholino
296	SCH ₃	2-Cl-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
297	SCH ₃	2-Cl-phenyl	4-morpholinocarbonyl
298	SCH ₃	2-Cl-phenyl	2-methyl-1-imidazolyl
299	SCH ₃	2-Cl-phenyl	5-methyl-1-imidazolyl
300	SCH ₃	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
301	SCH ₃	2-F-phenyl	2-(aminosulfonyl)phenyl
302	SCH ₃	2-F-phenyl	2-(methylaminosulfonyl)phenyl
303	SCH ₃	2-F-phenyl	1-pyrrolidinocarbonyl
304	SCH ₃	2-F-phenyl	2-(methylsulfonyl)phenyl
305	SCH ₃	2-F-phenyl	4-morpholino
306	SCH ₃	2-F-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
307	SCH ₃	2-F-phenyl	4-morpholinocarbonyl
308	SCH ₃	2-F-phenyl	2-methyl-1-imidazolyl
309	SCH ₃	2-F-phenyl	5-methyl-1-imidazolyl
310	SCH ₃	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
311	SCH ₃	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
312	SCH ₃	2,6-diF-phenyl	2-(methylaminosulfonyl)phenyl
313	SCH ₃	2,6-diF-phenyl	1-pyrrolidinocarbonyl

314	SCH ₃	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
315	SCH ₃	2,6-diF-phenyl	4-morpholino
316	SCH ₃	2,6-diF-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
317	SCH ₃	2,6-diF-phenyl	4-morpholinocarbonyl
318	SCH ₃	2,6-diF-phenyl	2-methyl-1-imidazolyl
319	SCH ₃	2,6-diF-phenyl	5-methyl-1-imidazolyl
320	SCH ₃	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
321	SOCH ₃	phenyl	2-(aminosulfonyl)phenyl
322	SOCH ₃	phenyl	2-(methylaminosulfonyl)phenyl
323	SOCH ₃	phenyl	1-pyrrolidinocarbonyl
324	SOCH ₃	phenyl	2-(methylsulfonyl)phenyl
325	SOCH ₃	phenyl	4-morpholino
326	SOCH ₃	phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
327	SOCH ₃	phenyl	4-morpholinocarbonyl
328	SOCH ₃	phenyl	2-methyl-1-imidazolyl
329	SOCH ₃	phenyl	5-methyl-1-imidazolyl
330	SOCH ₃	phenyl	2-methylsulfonyl-1-imidazolyl
331	SOCH ₃	2-pyridyl	2-(aminosulfonyl)phenyl
332	SOCH ₃	2-pyridyl	2-(methylaminosulfonyl)phenyl
333	SOCH ₃	2-pyridyl	1-pyrrolidinocarbonyl
334	SOCH ₃	2-pyridyl	2-(methylsulfonyl)phenyl
335	SOCH ₃	2-pyridyl	4-morpholino
336	SOCH ₃	2-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
337	SOCH ₃	2-pyridyl	4-morpholinocarbonyl
338	SOCH ₃	2-pyridyl	2-methyl-1-imidazolyl
339	SOCH ₃	2-pyridyl	5-methyl-1-imidazolyl
340	SOCH ₃	2-pyridyl	2-methylsulfonyl-1-imidazolyl
341	SOCH ₃	3-pyridyl	2-(aminosulfonyl)phenyl
342	SOCH ₃	3-pyridyl	2-(methylaminosulfonyl)phenyl
343	SOCH ₃	3-pyridyl	1-pyrrolidinocarbonyl
344	SOCH ₃	3-pyridyl	2-(methylsulfonyl)phenyl
345	SOCH ₃	3-pyridyl	4-morpholino
346	SOCH ₃	3-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
347	SOCH ₃	3-pyridyl	4-morpholinocarbonyl
348	SOCH ₃	3-pyridyl	2-methyl-1-imidazolyl
349	SOCH ₃	3-pyridyl	5-methyl-1-imidazolyl
350	SOCH ₃	3-pyridyl	2-methylsulfonyl-1-imidazolyl
351	SOCH ₃	2-pyrimidyl	2-(aminosulfonyl)phenyl
352	SOCH ₃	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
353	SOCH ₃	2-pyrimidyl	1-pyrrolidinocarbonyl
354	SOCH ₃	2-pyrimidyl	2-(methylsulfonyl)phenyl
355	SOCH ₃	2-pyrimidyl	4-morpholino
356	SOCH ₃	2-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
357	SOCH ₃	2-pyrimidyl	4-morpholinocarbonyl
358	SOCH ₃	2-pyrimidyl	2-methyl-1-imidazolyl
359	SOCH ₃	2-pyrimidyl	5-methyl-1-imidazolyl
360	SOCH ₃	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
361	SOCH ₃	5-pyrimidyl	2-(aminosulfonyl)phenyl

362	SOCH ₃	5-pyrimidyl	2-(methylaninosulfonyl)phenyl
363	SOCH ₃	5-pyrimidyl	1-pyrrolidinocarbonyl
364	SOCH ₃	5-pyrimidyl	2-(methylsulfonyl)phenyl
365	SOCH ₃	5-pyrimidyl	4-morpholino
366	SOCH ₃	5-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
367	SOCH ₃	5-pyrimidyl	4-morpholinocarbonyl
368	SOCH ₃	5-pyrimidyl	2-methyl-1-imidazolyl
369	SOCH ₃	5-pyrimidyl	5-methyl-1-imidazolyl
370	SOCH ₃	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
371	SOCH ₃	2-Cl-phenyl	2-(aminosulfonyl)phenyl
372	SOCH ₃	2-Cl-phenyl	2-(methylaninosulfonyl)phenyl
373	SOCH ₃	2-Cl-phenyl	1-pyrrolidinocarbonyl
374	SOCH ₃	2-Cl-phenyl	2-(methylsulfonyl)phenyl
375	SOCH ₃	2-Cl-phenyl	4-morpholino
376	SOCH ₃	2-Cl-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
377	SOCH ₃	2-Cl-phenyl	4-morpholinocarbonyl
378	SOCH ₃	2-Cl-phenyl	2-methyl-1-imidazolyl
379	SOCH ₃	2-Cl-phenyl	5-methyl-1-imidazolyl
380	SOCH ₃	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
381	SOCH ₃	2-F-phenyl	2-(aminosulfonyl)phenyl
382	SOCH ₃	2-F-phenyl	2-(methylaninosulfonyl)phenyl
383	SOCH ₃	2-F-phenyl	1-pyrrolidinocarbonyl
384	SOCH ₃	2-F-phenyl	2-(methylsulfonyl)phenyl
385	SOCH ₃	2-F-phenyl	4-morpholino
386	SOCH ₃	2-F-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
387	SOCH ₃	2-F-phenyl	4-morpholinocarbonyl
388	SOCH ₃	2-F-phenyl	2-methyl-1-imidazolyl
389	SOCH ₃	2-F-phenyl	5-methyl-1-imidazolyl
390	SOCH ₃	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
391	SOCH ₃	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
392	SOCH ₃	2,6-diF-phenyl	2-(methylaninosulfonyl)phenyl
393	SOCH ₃	2,6-diF-phenyl	1-pyrrolidinocarbonyl
394	SOCH ₃	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
395	SOCH ₃	2,6-diF-phenyl	4-morpholino
396	SOCH ₃	2,6-diF-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
397	SOCH ₃	2,6-diF-phenyl	4-morpholinocarbonyl
398	SOCH ₃	2,6-diF-phenyl	2-methyl-1-imidazolyl
399	SOCH ₃	2,6-diF-phenyl	5-methyl-1-imidazolyl
400	SOCH ₃	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
401	SO ₂ CH ₃	phenyl	2-(aminosulfonyl)phenyl
402	SO ₂ CH ₃	phenyl	2-(methylaninosulfonyl)phenyl
403	SO ₂ CH ₃	phenyl	1-pyrrolidinocarbonyl
404	SO ₂ CH ₃	phenyl	2-(methylsulfonyl)phenyl
405	SO ₂ CH ₃	phenyl	4-morpholino
406	SO ₂ CH ₃	phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
407	SO ₂ CH ₃	phenyl	4-morpholinocarbonyl
408	SO ₂ CH ₃	phenyl	2-methyl-1-imidazolyl
409	SO ₂ CH ₃	phenyl	5-methyl-1-imidazolyl

410	SO ₂ CH ₃	phenyl	2-methylsulfonyl-1-imidazolyl
411	SO ₂ CH ₃	2-pyridyl	2-(aminosulfonyl)phenyl
412	SO ₂ CH ₃	2-pyridyl	2-(methylaminosulfonyl)phenyl
413	SO ₂ CH ₃	2-pyridyl	1-pyrrolidinocarbonyl
414	SO ₂ CH ₃	2-pyridyl	2-(methylsulfonyl)phenyl
415	SO ₂ CH ₃	2-pyridyl	4-morpholino
416	SO ₂ CH ₃	2-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
417	SO ₂ CH ₃	2-pyridyl	4-morpholinocarbonyl
418	SO ₂ CH ₃	2-pyridyl	2-methyl-1-imidazolyl
419	SO ₂ CH ₃	2-pyridyl	5-methyl-1-imidazolyl
420	SO ₂ CH ₃	2-pyridyl	2-methylsulfonyl-1-imidazolyl
421	SO ₂ CH ₃	3-pyridyl	2-(aminosulfonyl)phenyl
422	SO ₂ CH ₃	3-pyridyl	2-(methylaminosulfonyl)phenyl
423	SO ₂ CH ₃	3-pyridyl	1-pyrrolidinocarbonyl
424	SO ₂ CH ₃	3-pyridyl	2-(methylsulfonyl)phenyl
425	SO ₂ CH ₃	3-pyridyl	4-morpholino
426	SO ₂ CH ₃	3-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
427	SO ₂ CH ₃	3-pyridyl	4-morpholinocarbonyl
428	SO ₂ CH ₃	3-pyridyl	2-methyl-1-imidazolyl
429	SO ₂ CH ₃	3-pyridyl	5-methyl-1-imidazolyl
430	SO ₂ CH ₃	3-pyridyl	2-methylsulfonyl-1-imidazolyl
431	SO ₂ CH ₃	2-pyrimidyl	2-(aminosulfonyl)phenyl
432	SO ₂ CH ₃	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
433	SO ₂ CH ₃	2-pyrimidyl	1-pyrrolidinocarbonyl
434	SO ₂ CH ₃	2-pyrimidyl	2-(methylsulfonyl)phenyl
435	SO ₂ CH ₃	2-pyrimidyl	4-morpholino
436	SO ₂ CH ₃	2-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
437	SO ₂ CH ₃	2-pyrimidyl	4-morpholinocarbonyl
438	SO ₂ CH ₃	2-pyrimidyl	2-methyl-1-imidazolyl
439	SO ₂ CH ₃	2-pyrimidyl	5-methyl-1-imidazolyl
440	SO ₂ CH ₃	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
441	SO ₂ CH ₃	5-pyrimidyl	2-(aminosulfonyl)phenyl
442	SO ₂ CH ₃	5-pyrimidyl	2-(methylaminosulfonyl)phenyl
443	SO ₂ CH ₃	5-pyrimidyl	1-pyrrolidinocarbonyl
444	SO ₂ CH ₃	5-pyrimidyl	2-(methylsulfonyl)phenyl
445	SO ₂ CH ₃	5-pyrimidyl	4-morpholino
446	SO ₂ CH ₃	5-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
447	SO ₂ CH ₃	5-pyrimidyl	4-morpholinocarbonyl
448	SO ₂ CH ₃	5-pyrimidyl	2-methyl-1-imidazolyl
449	SO ₂ CH ₃	5-pyrimidyl	5-methyl-1-imidazolyl
450	SO ₂ CH ₃	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
451	SO ₂ CH ₃	2-Cl-phenyl	2-(aminosulfonyl)phenyl
452	SO ₂ CH ₃	2-Cl-phenyl	2-(methylaminosulfonyl)phenyl
453	SO ₂ CH ₃	2-Cl-phenyl	1-pyrrolidinocarbonyl
454	SO ₂ CH ₃	2-Cl-phenyl	2-(methylsulfonyl)phenyl
455	SO ₂ CH ₃	2-Cl-phenyl	4-morpholino
456	SO ₂ CH ₃	2-Cl-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
457	SO ₂ CH ₃	2-Cl-phenyl	4-morpholinocarbonyl

458	SO ₂ CH ₃	2-Cl-phenyl	2-methyl-1-imidazolyl
459	SO ₂ CH ₃	2-Cl-phenyl	5-methyl-1-imidazolyl
460	SO ₂ CH ₃	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
461	SO ₂ CH ₃	2-F-phenyl	2-(aminosulfonyl)phenyl
462	SO ₂ CH ₃	2-F-phenyl	2-(methylaminosulfonyl)phenyl
463	SO ₂ CH ₃	2-F-phenyl	1-pyrrolidinocarbonyl
464	SO ₂ CH ₃	2-F-phenyl	2-(methylsulfonyl)phenyl
465	SO ₂ CH ₃	2-F-phenyl	4-morpholino
466	SO ₂ CH ₃	2-F-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
467	SO ₂ CH ₃	2-F-phenyl	4-morpholinocarbonyl
468	SO ₂ CH ₃	2-F-phenyl	2-methyl-1-imidazolyl
469	SO ₂ CH ₃	2-F-phenyl	5-methyl-1-imidazolyl
470	SO ₂ CH ₃	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
471	SO ₂ CH ₃	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
472	SO ₂ CH ₃	2,6-diF-phenyl	2-(methylaminosulfonyl)phenyl
473	SO ₂ CH ₃	2,6-diF-phenyl	1-pyrrolidinocarbonyl
474	SO ₂ CH ₃	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
475	SO ₂ CH ₃	2,6-diF-phenyl	4-morpholino
476	SO ₂ CH ₃	2,6-diF-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
477	SO ₂ CH ₃	2,6-diF-phenyl	4-morpholinocarbonyl
478	SO ₂ CH ₃	2,6-diF-phenyl	2-methyl-1-imidazolyl
479	SO ₂ CH ₃	2,6-diF-phenyl	5-methyl-1-imidazolyl
480	SO ₂ CH ₃	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
481	CH ₂ NH- SO ₂ CH ₃	phenyl	2-(aminosulfonyl)phenyl
482	CH ₂ NH- SO ₂ CH ₃	phenyl	2-(methylaminosulfonyl)phenyl
483	CH ₂ NH- SO ₂ CH ₃	phenyl	1-pyrrolidinocarbonyl
484	CH ₂ NH- SO ₂ CH ₃	phenyl	2-(methylsulfonyl)phenyl
485	CH ₂ NH- SO ₂ CH ₃	phenyl	4-morpholino
486	CH ₂ NH- SO ₂ CH ₃	phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
487	CH ₂ NH- SO ₂ CH ₃	phenyl	4-morpholinocarbonyl
488	CH ₂ NH- SO ₂ CH ₃	phenyl	2-methyl-1-imidazolyl
489	CH ₂ NH- SO ₂ CH ₃	phenyl	5-methyl-1-imidazolyl
490	CH ₂ NH- SO ₂ CH ₃	phenyl	2-methylsulfonyl-1-imidazolyl
491	CH ₂ NH- SO ₂ CH ₃	2-pyridyl	2-(aminosulfonyl)phenyl
492	CH ₂ NH- SO ₂ CH ₃	2-pyridyl	2-(methylaminosulfonyl)phenyl

493	CH ₂ NH-SO ₂ CH ₃	2-pyridyl	1-pyrrolidinocarbonyl
494	CH ₂ NH-SO ₂ CH ₃	2-pyridyl	2-(methylsulfonyl)phenyl
495	CH ₂ NH-SO ₂ CH ₃	2-pyridyl	4-morpholino
496	CH ₂ NH-SO ₂ CH ₃	2-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
497	CH ₂ NH-SO ₂ CH ₃	2-pyridyl	4-morpholinocarbonyl
498	CH ₂ NH-SO ₂ CH ₃	2-pyridyl	2-methyl-1-imidazolyl
499	CH ₂ NH-SO ₂ CH ₃	2-pyridyl	5-methyl-1-imidazolyl
500	CH ₂ NH-SO ₂ CH ₃	2-pyridyl	2-methylsulfonyl-1-imidazolyl
501	CH ₂ NH-SO ₂ CH ₃	3-pyridyl	2-(aminosulfonyl)phenyl
502	CH ₂ NH-SO ₂ CH ₃	3-pyridyl	2-(methylaminosulfonyl)phenyl
503	CH ₂ NH-SO ₂ CH ₃	3-pyridyl	1-pyrrolidinocarbonyl
504	CH ₂ NH-SO ₂ CH ₃	3-pyridyl	2-(methylsulfonyl)phenyl
505	CH ₂ NH-SO ₂ CH ₃	3-pyridyl	4-morpholino
506	CH ₂ NH-SO ₂ CH ₃	3-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
507	CH ₂ NH-SO ₂ CH ₃	3-pyridyl	4-morpholinocarbonyl
508	CH ₂ NH-SO ₂ CH ₃	3-pyridyl	2-methyl-1-imidazolyl
509	CH ₂ NH-SO ₂ CH ₃	3-pyridyl	5-methyl-1-imidazolyl
510	CH ₂ NH-SO ₂ CH ₃	3-pyridyl	2-methylsulfonyl-1-imidazolyl
511	CH ₂ NH-SO ₂ CH ₃	2-pyrimidyl	2-(aminosulfonyl)phenyl
512	CH ₂ NH-SO ₂ CH ₃	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
513	CH ₂ NH-SO ₂ CH ₃	2-pyrimidyl	1-pyrrolidinocarbonyl
514	CH ₂ NH-SO ₂ CH ₃	2-pyrimidyl	2-(methylsulfonyl)phenyl
515	CH ₂ NH-SO ₂ CH ₃	2-pyrimidyl	4-morpholino
516	CH ₂ NH-SO ₂ CH ₃	2-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl

517	CH ₂ NH-SO ₂ CH ₃	2-pyrimidyl	4-morpholinocarbonyl
518	CH ₂ NH-SO ₂ CH ₃	2-pyrimidyl	2-methyl-1-imidazolyl
519	CH ₂ NH-SO ₂ CH ₃	2-pyrimidyl	5-methyl-1-imidazolyl
520	CH ₂ NH-SO ₂ CH ₃	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
521	CH ₂ NH-SO ₂ CH ₃	5-pyrimidyl	2-(aminosulfonyl)phenyl
522	CH ₂ NH-SO ₂ CH ₃	5-pyrimidyl	2-(methylaminosulfonyl)phenyl
523	CH ₂ NH-SO ₂ CH ₃	5-pyrimidyl	1-pyrrolidinocarbonyl
524	CH ₂ NH-SO ₂ CH ₃	5-pyrimidyl	2-(methylsulfonyl)phenyl
525	CH ₂ NH-SO ₂ CH ₃	5-pyrimidyl	4-morpholino
526	CH ₂ NH-SO ₂ CH ₃	5-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
527	CH ₂ NH-SO ₂ CH ₃	5-pyrimidyl	4-morpholinocarbonyl
528	CH ₂ NH-SO ₂ CH ₃	5-pyrimidyl	2-methyl-1-imidazolyl
529	CH ₂ NH-SO ₂ CH ₃	5-pyrimidyl	5-methyl-1-imidazolyl
530	CH ₂ NH-SO ₂ CH ₃	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
531	CH ₂ NH-SO ₂ CH ₃	2-Cl-phenyl	2-(aminosulfonyl)phenyl
532	CH ₂ NH-SO ₂ CH ₃	2-Cl-phenyl	2-(methylaminosulfonyl)phenyl
533	CH ₂ NH-SO ₂ CH ₃	2-Cl-phenyl	1-pyrrolidinocarbonyl
534	CH ₂ NH-SO ₂ CH ₃	2-Cl-phenyl	2-(methylsulfonyl)phenyl
535	CH ₂ NH-SO ₂ CH ₃	2-Cl-phenyl	4-morpholino
536	CH ₂ NH-SO ₂ CH ₃	2-Cl-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
537	CH ₂ NH-SO ₂ CH ₃	2-Cl-phenyl	4-morpholinocarbonyl
538	CH ₂ NH-SO ₂ CH ₃	2-Cl-phenyl	2-methyl-1-imidazolyl
539	CH ₂ NH-SO ₂ CH ₃	2-Cl-phenyl	5-methyl-1-imidazolyl
540	CH ₂ NH-SO ₂ CH ₃	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl

541	CH ₂ NH-SO ₂ CH ₃	2-F-phenyl	2-(aminosulfonyl)phenyl
542	CH ₂ NH-SO ₂ CH ₃	2-F-phenyl	2-(methylaminosulfonyl)phenyl
543	CH ₂ NH-SO ₂ CH ₃	2-F-phenyl	1-pyrrolidinocarbonyl
544	CH ₂ NH-SO ₂ CH ₃	2-F-phenyl	2-(methylsulfonyl)phenyl
545	CH ₂ NH-SO ₂ CH ₃	2-F-phenyl	4-morpholino
546	CH ₂ NH-SO ₂ CH ₃	2-F-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
547	CH ₂ NH-SO ₂ CH ₃	2-F-phenyl	4-morpholinocarbonyl
548	CH ₂ NH-SO ₂ CH ₃	2-F-phenyl	2-methyl-1-imidazolyl
549	CH ₂ NH-SO ₂ CH ₃	2-F-phenyl	5-methyl-1-imidazolyl
550	CH ₂ NH-SO ₂ CH ₃	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
551	CH ₂ NH-SO ₂ CH ₃	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
552	CH ₂ NH-SO ₂ CH ₃	2,6-diF-phenyl	2-(methylaminosulfonyl)phenyl
553	CH ₂ NH-SO ₂ CH ₃	2,6-diF-phenyl	1-pyrrolidinocarbonyl
554	CH ₂ NH-SO ₂ CH ₃	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
555	CH ₂ NH-SO ₂ CH ₃	2,6-diF-phenyl	4-morpholino
556	CH ₂ NH-SO ₂ CH ₃	2,6-diF-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
557	CH ₂ NH-SO ₂ CH ₃	2,6-diF-phenyl	4-morpholinocarbonyl
558	CH ₂ NH-SO ₂ CH ₃	2,6-diF-phenyl	2-methyl-1-imidazolyl
559	CH ₂ NH-SO ₂ CH ₃	2,6-diF-phenyl	5-methyl-1-imidazolyl
560	CH ₂ NH-SO ₂ CH ₃	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
561	Cl	phenyl	2-(aminosulfonyl)phenyl
562	Cl	phenyl	2-(methylaminosulfonyl)phenyl
563	Cl	phenyl	1-pyrrolidinocarbonyl
564	Cl	phenyl	2-(methylsulfonyl)phenyl
565	Cl	phenyl	4-morpholino
566	Cl	phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
567	Cl	phenyl	4-morpholinocarbonyl
568	Cl	phenyl	2-methyl-1-imidazolyl
569	Cl	phenyl	5-methyl-1-imidazolyl

570	Cl	phenyl	2-methylsulfonyl-1-imidazolyl
571	Cl	2-pyridyl	2-(aminosulfonyl)phenyl
572	Cl	2-pyridyl	2-(methylaminosulfonyl)phenyl
573	Cl	2-pyridyl	1-pyrrolidinocarbonyl
574	Cl	2-pyridyl	2-(methylsulfonyl)phenyl
575	Cl	2-pyridyl	4-morpholino
576	Cl	2-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
577	Cl	2-pyridyl	4-morpholinocarbonyl
578	Cl	2-pyridyl	2-methyl-1-imidazolyl
579	Cl	2-pyridyl	5-methyl-1-imidazolyl
580	Cl	2-pyridyl	2-methylsulfonyl-1-imidazolyl
581	Cl	3-pyridyl	2-(aminosulfonyl)phenyl
582	Cl	3-pyridyl	2-(methylaminosulfonyl)phenyl
583	Cl	3-pyridyl	1-pyrrolidinocarbonyl
584	Cl	3-pyridyl	2-(methylsulfonyl)phenyl
585	Cl	3-pyridyl	4-morpholino
586	Cl	3-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
587	Cl	3-pyridyl	4-morpholinocarbonyl
588	Cl	3-pyridyl	2-methyl-1-imidazolyl
589	Cl	3-pyridyl	5-methyl-1-imidazolyl
590	Cl	3-pyridyl	2-methylsulfonyl-1-imidazolyl
591	Cl	2-pyrimidyl	2-(aminosulfonyl)phenyl
592	Cl	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
593	Cl	2-pyrimidyl	1-pyrrolidinocarbonyl
594	Cl	2-pyrimidyl	2-(methylsulfonyl)phenyl
595	Cl	2-pyrimidyl	4-morpholino
596	Cl	2-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
597	Cl	2-pyrimidyl	4-morpholinocarbonyl
598	Cl	2-pyrimidyl	2-methyl-1-imidazolyl
599	Cl	2-pyrimidyl	5-methyl-1-imidazolyl
600	Cl	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
601	Cl	5-pyrimidyl	2-(aminosulfonyl)phenyl
602	Cl	5-pyrimidyl	2-(methylaminosulfonyl)phenyl
603	Cl	5-pyrimidyl	1-pyrrolidinocarbonyl
604	Cl	5-pyrimidyl	2-(methylsulfonyl)phenyl
605	Cl	5-pyrimidyl	4-morpholino
606	Cl	5-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
607	Cl	5-pyrimidyl	4-morpholinocarbonyl
608	Cl	5-pyrimidyl	2-methyl-1-imidazolyl
609	Cl	5-pyrimidyl	5-methyl-1-imidazolyl
610	Cl	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
611	Cl	2-Cl-phenyl	2-(aminosulfonyl)phenyl
612	Cl	2-Cl-phenyl	2-(methylaminosulfonyl)phenyl
613	Cl	2-Cl-phenyl	1-pyrrolidinocarbonyl
614	Cl	2-Cl-phenyl	2-(methylsulfonyl)phenyl
615	Cl	2-Cl-phenyl	4-morpholino
616	Cl	2-Cl-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
617	Cl	2-Cl-phenyl	4-morpholinocarbonyl
618	Cl	2-Cl-phenyl	2-methyl-1-imidazolyl
619	Cl	2-Cl-phenyl	5-methyl-1-imidazolyl
620	Cl	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
621	Cl	2-F-phenyl	2-(aminosulfonyl)phenyl
622	Cl	2-F-phenyl	2-(methylaminosulfonyl)phenyl
623	Cl	2-F-phenyl	1-pyrrolidinocarbonyl
624	Cl	2-F-phenyl	2-(methylsulfonyl)phenyl

625	Cl	2-F-phenyl	4-morpholino
626	Cl	2-F-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
627	Cl	2-F-phenyl	4-morpholinocarbonyl
628	Cl	2-F-phenyl	2-methyl-1-imidazolyl
629	Cl	2-F-phenyl	5-methyl-1-imidazolyl
630	Cl	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
631	Cl	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
632	Cl	2,6-diF-phenyl	2-(methylaminosulfonyl)phenyl
633	Cl	2,6-diF-phenyl	1-pyrrolidinocarbonyl
634	Cl	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
635	Cl	2,6-diF-phenyl	4-morpholino
636	Cl	2,6-diF-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
637	Cl	2,6-diF-phenyl	4-morpholinocarbonyl
638	Cl	2,6-diF-phenyl	2-methyl-1-imidazolyl
639	Cl	2,6-diF-phenyl	5-methyl-1-imidazolyl
640	Cl	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
641	F	phenyl	2-(aminosulfonyl)phenyl
642	F	phenyl	2-(methylaminosulfonyl)phenyl
643	F	phenyl	1-pyrrolidinocarbonyl
644	F	phenyl	2-(methylsulfonyl)phenyl
645	F	phenyl	4-morpholino
646	F	phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
647	F	phenyl	4-morpholinocarbonyl
648	F	phenyl	2-methyl-1-imidazolyl
649	F	phenyl	5-methyl-1-imidazolyl
650	F	phenyl	2-methylsulfonyl-1-imidazolyl
651	F	2-pyridyl	2-(aminosulfonyl)phenyl
652	F	2-pyridyl	2-(methylaminosulfonyl)phenyl
653	F	2-pyridyl	1-pyrrolidinocarbonyl
654	F	2-pyridyl	2-(methylsulfonyl)phenyl
655	F	2-pyridyl	4-morpholino
656	F	2-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
657	F	2-pyridyl	4-morpholinocarbonyl
658	F	2-pyridyl	2-methyl-1-imidazolyl
659	F	2-pyridyl	5-methyl-1-imidazolyl
660	F	2-pyridyl	2-methylsulfonyl-1-imidazolyl
661	F	3-pyridyl	2-(aminosulfonyl)phenyl
662	F	3-pyridyl	2-(methylaminosulfonyl)phenyl
663	F	3-pyridyl	1-pyrrolidinocarbonyl
664	F	3-pyridyl	2-(methylsulfonyl)phenyl
665	F	3-pyridyl	4-morpholino
666	F	3-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
667	F	3-pyridyl	4-morpholinocarbonyl
668	F	3-pyridyl	2-methyl-1-imidazolyl
669	F	3-pyridyl	5-methyl-1-imidazolyl
670	F	3-pyridyl	2-methylsulfonyl-1-imidazolyl
671	F	2-pyrimidyl	2-(aminosulfonyl)phenyl
672	F	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
673	F	2-pyrimidyl	1-pyrrolidinocarbonyl
674	F	2-pyrimidyl	2-(methylsulfonyl)phenyl
675	F	2-pyrimidyl	4-morpholino
676	F	2-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
677	F	2-pyrimidyl	4-morpholinocarbonyl
678	F	2-pyrimidyl	2-methyl-1-imidazolyl
679	F	2-pyrimidyl	5-methyl-1-imidazolyl

680	F	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
681	F	5-pyrimidyl	2-(aminosulfonyl)phenyl
682	F	5-pyrimidyl	2-(methylaminosulfonyl)phenyl
683	F	5-pyrimidyl	1-pyrrolidinocarbonyl
684	F	5-pyrimidyl	2-(methylsulfonyl)phenyl
685	F	5-pyrimidyl	4-morpholino
686	F	5-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
687	F	5-pyrimidyl	4-morpholinocarbonyl
688	F	5-pyrimidyl	2-methyl-1-imidazolyl
689	F	5-pyrimidyl	5-methyl-1-imidazolyl
690	F	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
691	F	2-Cl-phenyl	2-(aminosulfonyl)phenyl
692	F	2-Cl-phenyl	2-(methylaminosulfonyl)phenyl
693	F	2-Cl-phenyl	1-pyrrolidinocarbonyl
694	F	2-Cl-phenyl	2-(methylsulfonyl)phenyl
695	F	2-Cl-phenyl	4-morpholino
696	F	2-Cl-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
697	F	2-Cl-phenyl	4-morpholinocarbonyl
698	F	2-Cl-phenyl	2-methyl-1-imidazolyl
699	F	2-Cl-phenyl	5-methyl-1-imidazolyl
700	F	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
701	F	2-F-phenyl	2-(aminosulfonyl)phenyl
702	F	2-F-phenyl	2-(methylaminosulfonyl)phenyl
703	F	2-F-phenyl	1-pyrrolidinocarbonyl
704	F	2-F-phenyl	2-(methylsulfonyl)phenyl
705	F	2-F-phenyl	4-morpholino
706	F	2-F-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
707	F	2-F-phenyl	4-morpholinocarbonyl
708	F	2-F-phenyl	2-methyl-1-imidazolyl
709	F	2-F-phenyl	5-methyl-1-imidazolyl
710	F	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
711	F	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
712	F	2,6-diF-phenyl	2-(methylaminosulfonyl)phenyl
713	F	2,6-diF-phenyl	1-pyrrolidinocarbonyl
714	F	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
715	F	2,6-diF-phenyl	4-morpholino
716	F	2,6-diF-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
717	F	2,6-diF-phenyl	4-morpholinocarbonyl
718	F	2,6-diF-phenyl	2-methyl-1-imidazolyl
719	F	2,6-diF-phenyl	5-methyl-1-imidazolyl
720	F	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
721	CO ₂ CH ₃	phenyl	2-(aminosulfonyl)phenyl
722	CO ₂ CH ₃	phenyl	2-(methylaminosulfonyl)phenyl
723	CO ₂ CH ₃	phenyl	1-pyrrolidinocarbonyl
724	CO ₂ CH ₃	phenyl	2-(methylsulfonyl)phenyl
725	CO ₂ CH ₃	phenyl	4-morpholino
726	CO ₂ CH ₃	phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
727	CO ₂ CH ₃	phenyl	4-morpholinocarbonyl
728	CO ₂ CH ₃	phenyl	2-methyl-1-imidazolyl
729	CO ₂ CH ₃	phenyl	5-methyl-1-imidazolyl
730	CO ₂ CH ₃	phenyl	2-methylsulfonyl-1-imidazolyl
731	CO ₂ CH ₃	2-pyridyl	2-(aminosulfonyl)phenyl
732	CO ₂ CH ₃	2-pyridyl	2-(methylaminosulfonyl)phenyl

733	CO ₂ CH ₃	2-pyridyl	1-pyrrolidinocarbonyl
734	CO ₂ CH ₃	2-pyridyl	2-(methylsulfonyl)phenyl
735	CO ₂ CH ₃	2-pyridyl	4-morpholino
736	CO ₂ CH ₃	2-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
737	CO ₂ CH ₃	2-pyridyl	4-morpholinocarbonyl
738	CO ₂ CH ₃	2-pyridyl	2-methyl-1-imidazolyl
739	CO ₂ CH ₃	2-pyridyl	5-methyl-1-imidazolyl
740	CO ₂ CH ₃	2-pyridyl	2-methylsulfonyl-1-imidazolyl
741	CO ₂ CH ₃	3-pyridyl	2-(aminosulfonyl)phenyl
742	CO ₂ CH ₃	3-pyridyl	2-(methylaminosulfonyl)phenyl
743	CO ₂ CH ₃	3-pyridyl	1-pyrrolidinocarbonyl
744	CO ₂ CH ₃	3-pyridyl	2-(methylsulfonyl)phenyl
745	CO ₂ CH ₃	3-pyridyl	4-morpholino
746	CO ₂ CH ₃	3-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
747	CO ₂ CH ₃	3-pyridyl	4-morpholinocarbonyl
748	CO ₂ CH ₃	3-pyridyl	2-methyl-1-imidazolyl
749	CO ₂ CH ₃	3-pyridyl	5-methyl-1-imidazolyl
750	CO ₂ CH ₃	3-pyridyl	2-methylsulfonyl-1-imidazolyl
751	CO ₂ CH ₃	2-pyrimidyl	2-(aminosulfonyl)phenyl
752	CO ₂ CH ₃	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
753	CO ₂ CH ₃	2-pyrimidyl	1-pyrrolidinocarbonyl
754	CO ₂ CH ₃	2-pyrimidyl	2-(methylsulfonyl)phenyl
755	CO ₂ CH ₃	2-pyrimidyl	4-morpholino
756	CO ₂ CH ₃	2-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
757	CO ₂ CH ₃	2-pyrimidyl	4-morpholinocarbonyl
758	CO ₂ CH ₃	2-pyrimidyl	2-methyl-1-imidazolyl
759	CO ₂ CH ₃	2-pyrimidyl	5-methyl-1-imidazolyl
760	CO ₂ CH ₃	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
761	CO ₂ CH ₃	5-pyrimidyl	2-(aminosulfonyl)phenyl
762	CO ₂ CH ₃	5-pyrimidyl	2-(methylaminosulfonyl)phenyl
763	CO ₂ CH ₃	5-pyrimidyl	1-pyrrolidinocarbonyl
764	CO ₂ CH ₃	5-pyrimidyl	2-(methylsulfonyl)phenyl
765	CO ₂ CH ₃	5-pyrimidyl	4-morpholino
766	CO ₂ CH ₃	5-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
767	CO ₂ CH ₃	5-pyrimidyl	4-morpholinocarbonyl
768	CO ₂ CH ₃	5-pyrimidyl	2-methyl-1-imidazolyl
769	CO ₂ CH ₃	5-pyrimidyl	5-methyl-1-imidazolyl
770	CO ₂ CH ₃	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
771	CO ₂ CH ₃	2-Cl-phenyl	2-(aminosulfonyl)phenyl
772	CO ₂ CH ₃	2-Cl-phenyl	2-(methylaminosulfonyl)phenyl
773	CO ₂ CH ₃	2-Cl-phenyl	1-pyrrolidinocarbonyl
774	CO ₂ CH ₃	2-Cl-phenyl	2-(methylsulfonyl)phenyl
775	CO ₂ CH ₃	2-Cl-phenyl	4-morpholino
776	CO ₂ CH ₃	2-Cl-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
777	CO ₂ CH ₃	2-Cl-phenyl	4-morpholinocarbonyl
778	CO ₂ CH ₃	2-Cl-phenyl	2-methyl-1-imidazolyl
779	CO ₂ CH ₃	2-Cl-phenyl	5-methyl-1-imidazolyl
780	CO ₂ CH ₃	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl

781	CO ₂ CH ₃	2-F-phenyl	2-(aminosulfonyl)phenyl
782	CO ₂ CH ₃	2-F-phenyl	2-(methylaminosulfonyl)phenyl
783	CO ₂ CH ₃	2-F-phenyl	1-pyrrolidinocarbonyl
784	CO ₂ CH ₃	2-F-phenyl	2-(methylsulfonyl)phenyl
785	CO ₂ CH ₃	2-F-phenyl	4-morpholino
786	CO ₂ CH ₃	2-F-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
787	CO ₂ CH ₃	2-F-phenyl	4-morpholinocarbonyl
788	CO ₂ CH ₃	2-F-phenyl	2-methyl-1-imidazolyl
789	CO ₂ CH ₃	2-F-phenyl	5-methyl-1-imidazolyl
790	CO ₂ CH ₃	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
791	CO ₂ CH ₃	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
792	CO ₂ CH ₃	2,6-diF-phenyl	2-(methylaminosulfonyl)phenyl
793	CO ₂ CH ₃	2,6-diF-phenyl	1-pyrrolidinocarbonyl
794	CO ₂ CH ₃	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
795	CO ₂ CH ₃	2,6-diF-phenyl	4-morpholino
796	CO ₂ CH ₃	2,6-diF-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
797	CO ₂ CH ₃	2,6-diF-phenyl	4-morpholinocarbonyl
798	CO ₂ CH ₃	2,6-diF-phenyl	2-methyl-1-imidazolyl
799	CO ₂ CH ₃	2,6-diF-phenyl	5-methyl-1-imidazolyl
800	CO ₂ CH ₃	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
801	CH ₂ OCH ₃	phenyl	2-(aminosulfonyl)phenyl
802	CH ₂ OCH ₃	phenyl	2-(methylaminosulfonyl)phenyl
803	CH ₂ OCH ₃	phenyl	1-pyrrolidinocarbonyl
804	CH ₂ OCH ₃	phenyl	2-(methylsulfonyl)phenyl
805	CH ₂ OCH ₃	phenyl	4-morpholino
806	CH ₂ OCH ₃	phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
807	CH ₂ OCH ₃	phenyl	4-morpholinocarbonyl
808	CH ₂ OCH ₃	phenyl	2-methyl-1-imidazolyl
809	CH ₂ OCH ₃	phenyl	5-methyl-1-imidazolyl
810	CH ₂ OCH ₃	phenyl	2-methylsulfonyl-1-imidazolyl
811	CH ₂ OCH ₃	2-pyridyl	2-(aminosulfonyl)phenyl
812	CH ₂ OCH ₃	2-pyridyl	2-(methylaminosulfonyl)phenyl
813	CH ₂ OCH ₃	2-pyridyl	1-pyrrolidinocarbonyl
814	CH ₂ OCH ₃	2-pyridyl	2-(methylsulfonyl)phenyl
815	CH ₂ OCH ₃	2-pyridyl	4-morpholino
816	CH ₂ OCH ₃	2-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
817	CH ₂ OCH ₃	2-pyridyl	4-morpholinocarbonyl
818	CH ₂ OCH ₃	2-pyridyl	2-methyl-1-imidazolyl
819	CH ₂ OCH ₃	2-pyridyl	5-methyl-1-imidazolyl
820	CH ₂ OCH ₃	2-pyridyl	2-methylsulfonyl-1-imidazolyl
821	CH ₂ OCH ₃	3-pyridyl	2-(aminosulfonyl)phenyl
822	CH ₂ OCH ₃	3-pyridyl	2-(methylaminosulfonyl)phenyl
823	CH ₂ OCH ₃	3-pyridyl	1-pyrrolidinocarbonyl
824	CH ₂ OCH ₃	3-pyridyl	2-(methylsulfonyl)phenyl
825	CH ₂ OCH ₃	3-pyridyl	4-morpholino
826	CH ₂ OCH ₃	3-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
827	CH ₂ OCH ₃	3-pyridyl	4-morpholinocarbonyl
828	CH ₂ OCH ₃	3-pyridyl	2-methyl-1-imidazolyl

829	CH ₂ OCH ₃	3-pyridyl	5-methyl-1-imidazolyl
830	CH ₂ OCH ₃	3-pyridyl	2-methylsulfonyl-1-imidazolyl
831	CH ₂ OCH ₃	2-pyrimidyl	2-(aminosulfonyl)phenyl
832	CH ₂ OCH ₃	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
833	CH ₂ OCH ₃	2-pyrimidyl	1-pyrrolidinocarbonyl
834	CH ₂ OCH ₃	2-pyrimidyl	2-(methylsulfonyl)phenyl
835	CH ₂ OCH ₃	2-pyrimidyl	4-morpholino
836	CH ₂ OCH ₃	2-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
837	CH ₂ OCH ₃	2-pyrimidyl	4-morpholinocarbonyl
838	CH ₂ OCH ₃	2-pyrimidyl	2-methyl-1-imidazolyl
839	CH ₂ OCH ₃	2-pyrimidyl	5-methyl-1-imidazolyl
840	CH ₂ OCH ₃	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
841	CH ₂ OCH ₃	5-pyrimidyl	2-(aminosulfonyl)phenyl
842	CH ₂ OCH ₃	5-pyrimidyl	2-(methylaminosulfonyl)phenyl
843	CH ₂ OCH ₃	5-pyrimidyl	1-pyrrolidinocarbonyl
844	CH ₂ OCH ₃	5-pyrimidyl	2-(methylsulfonyl)phenyl
845	CH ₂ OCH ₃	5-pyrimidyl	4-morpholino
846	CH ₂ OCH ₃	5-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
847	CH ₂ OCH ₃	5-pyrimidyl	4-morpholinocarbonyl
848	CH ₂ OCH ₃	5-pyrimidyl	2-methyl-1-imidazolyl
849	CH ₂ OCH ₃	5-pyrimidyl	5-methyl-1-imidazolyl
850	CH ₂ OCH ₃	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
851	CH ₂ OCH ₃	2-Cl-phenyl	2-(aminosulfonyl)phenyl
852	CH ₂ OCH ₃	2-Cl-phenyl	2-(methylaminosulfonyl)phenyl
853	CH ₂ OCH ₃	2-Cl-phenyl	1-pyrrolidinocarbonyl
854	CH ₂ OCH ₃	2-Cl-phenyl	2-(methylsulfonyl)phenyl
855	CH ₂ OCH ₃	2-Cl-phenyl	4-morpholino
856	CH ₂ OCH ₃	2-Cl-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
857	CH ₂ OCH ₃	2-Cl-phenyl	4-morpholinocarbonyl
858	CH ₂ OCH ₃	2-Cl-phenyl	2-methyl-1-imidazolyl
859	CH ₂ OCH ₃	2-Cl-phenyl	5-methyl-1-imidazolyl
860	CH ₂ OCH ₃	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
861	CH ₂ OCH ₃	2-F-phenyl	2-(aminosulfonyl)phenyl
862	CH ₂ OCH ₃	2-F-phenyl	2-(methylaminosulfonyl)phenyl
863	CH ₂ OCH ₃	2-F-phenyl	1-pyrrolidinocarbonyl
864	CH ₂ OCH ₃	2-F-phenyl	2-(methylsulfonyl)phenyl
865	CH ₂ OCH ₃	2-F-phenyl	4-morpholino
866	CH ₂ OCH ₃	2-F-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
867	CH ₂ OCH ₃	2-F-phenyl	4-morpholinocarbonyl
868	CH ₂ OCH ₃	2-F-phenyl	2-methyl-1-imidazolyl
869	CH ₂ OCH ₃	2-F-phenyl	5-methyl-1-imidazolyl
870	CH ₂ OCH ₃	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
871	CH ₂ OCH ₃	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
872	CH ₂ OCH ₃	2,6-diF-phenyl	2-(methylaminosulfonyl)phenyl
873	CH ₂ OCH ₃	2,6-diF-phenyl	1-pyrrolidinocarbonyl
874	CH ₂ OCH ₃	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
875	CH ₂ OCH ₃	2,6-diF-phenyl	4-morpholino
876	CH ₂ OCH ₃	2,6-diF-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl

877	CH ₂ OCH ₃	2,6-diF-phenyl	4-morpholinocarbonyl
878	CH ₂ OCH ₃	2,6-diF-phenyl	2-methyl-1-imidazolyl
879	CH ₂ OCH ₃	2,6-diF-phenyl	5-methyl-1-imidazolyl
880	CH ₂ OCH ₃	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
881	CONH ₂	phenyl	2-(aminosulfonyl)phenyl
882	CONH ₂	phenyl	2-(methylaminosulfonyl)phenyl
883	CONH ₂	phenyl	1-pyrrolidinocarbonyl
884	CONH ₂	phenyl	2-(methylsulfonyl)phenyl
885	CONH ₂	phenyl	4-morpholino
886	CONH ₂	phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
887	CONH ₂	phenyl	4-morpholinocarbonyl
888	CONH ₂	phenyl	2-methyl-1-imidazolyl
889	CONH ₂	phenyl	5-methyl-1-imidazolyl
890	CONH ₂	phenyl	2-methylsulfonyl-1-imidazolyl
891	CONH ₂	2-pyridyl	2-(aminosulfonyl)phenyl
892	CONH ₂	2-pyridyl	2-(methylaminosulfonyl)phenyl
893	CONH ₂	2-pyridyl	1-pyrrolidinocarbonyl
894	CONH ₂	2-pyridyl	2-(methylsulfonyl)phenyl
895	CONH ₂	2-pyridyl	4-morpholino
896	CONH ₂	2-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
897	CONH ₂	2-pyridyl	4-morpholinocarbonyl
898	CONH ₂	2-pyridyl	2-methyl-1-imidazolyl
899	CONH ₂	2-pyridyl	5-methyl-1-imidazolyl
900	CONH ₂	2-pyridyl	2-methylsulfonyl-1-imidazolyl
901	CONH ₂	3-pyridyl	2-(aminosulfonyl)phenyl
902	CONH ₂	3-pyridyl	2-(methylaminosulfonyl)phenyl
903	CONH ₂	3-pyridyl	1-pyrrolidinocarbonyl
904	CONH ₂	3-pyridyl	2-(methylsulfonyl)phenyl
905	CONH ₂	3-pyridyl	4-morpholino
906	CONH ₂	3-pyridyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
907	CONH ₂	3-pyridyl	4-morpholinocarbonyl
908	CONH ₂	3-pyridyl	2-methyl-1-imidazolyl
909	CONH ₂	3-pyridyl	5-methyl-1-imidazolyl
910	CONH ₂	3-pyridyl	2-methylsulfonyl-1-imidazolyl
911	CONH ₂	2-pyrimidyl	2-(aminosulfonyl)phenyl
912	CONH ₂	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
913	CONH ₂	2-pyrimidyl	1-pyrrolidinocarbonyl
914	CONH ₂	2-pyrimidyl	2-(methylsulfonyl)phenyl
915	CONH ₂	2-pyrimidyl	4-morpholino
916	CONH ₂	2-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
917	CONH ₂	2-pyrimidyl	4-morpholinocarbonyl
918	CONH ₂	2-pyrimidyl	2-methyl-1-imidazolyl
919	CONH ₂	2-pyrimidyl	5-methyl-1-imidazolyl
920	CONH ₂	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
921	CONH ₂	5-pyrimidyl	2-(aminosulfonyl)phenyl
922	CONH ₂	5-pyrimidyl	2-(methylaminosulfonyl)phenyl
923	CONH ₂	5-pyrimidyl	1-pyrrolidinocarbonyl
924	CONH ₂	5-pyrimidyl	2-(methylsulfonyl)phenyl

925	CONH ₂	5-pyrimidyl	4-morpholino
926	CONH ₂	5-pyrimidyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
927	CONH ₂	5-pyrimidyl	4-morpholinocarbonyl
928	CONH ₂	5-pyrimidyl	2-methyl-1-imidazolyl
929	CONH ₂	5-pyrimidyl	5-methyl-1-imidazolyl
930	CONH ₂	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
931	CONH ₂	2-Cl-phenyl	2-(aminosulfonyl)phenyl
932	CONH ₂	2-Cl-phenyl	2-(methylaminosulfonyl)phenyl
933	CONH ₂	2-Cl-phenyl	1-pyrrolidinocarbonyl
934	CONH ₂	2-Cl-phenyl	2-(methylsulfonyl)phenyl
935	CONH ₂	2-Cl-phenyl	4-morpholino
936	CONH ₂	2-Cl-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
937	CONH ₂	2-Cl-phenyl	4-morpholinocarbonyl
938	CONH ₂	2-Cl-phenyl	2-methyl-1-imidazolyl
939	CONH ₂	2-Cl-phenyl	5-methyl-1-imidazolyl
940	CONH ₂	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
941	CONH ₂	2-F-phenyl	2-(aminosulfonyl)phenyl
942	CONH ₂	2-F-phenyl	2-(methylaminosulfonyl)phenyl
943	CONH ₂	2-F-phenyl	1-pyrrolidinocarbonyl
944	CONH ₂	2-F-phenyl	2-(methylsulfonyl)phenyl
945	CONH ₂	2-F-phenyl	4-morpholino
946	CONH ₂	2-F-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
947	CONH ₂	2-F-phenyl	4-morpholinocarbonyl
948	CONH ₂	2-F-phenyl	2-methyl-1-imidazolyl
949	CONH ₂	2-F-phenyl	5-methyl-1-imidazolyl
950	CONH ₂	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
951	CONH ₂	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
952	CONH ₂	2,6-diF-phenyl	2-(methylaminosulfonyl)phenyl
953	CONH ₂	2,6-diF-phenyl	1-pyrrolidinocarbonyl
954	CONH ₂	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
955	CONH ₂	2,6-diF-phenyl	4-morpholino
956	CONH ₂	2,6-diF-phenyl	2-(1'-CF ₃ -tetrazol-2-yl)phenyl
957	CONH ₂	2,6-diF-phenyl	4-morpholinocarbonyl
958	CONH ₂	2,6-diF-phenyl	2-methyl-1-imidazolyl
959	CONH ₂	2,6-diF-phenyl	5-methyl-1-imidazolyl
960	CONH ₂	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl

Utility

The compounds of this invention are useful as anticoagulants for the treatment or prevention of thromboembolic disorders in mammals. The term "thromboembolic disorders" as used herein includes arterial or venous cardiovascular or cerebrovascular thromboembolic disorders, including, for example, unstable angina, first or recurrent myocardial infarction, ischemic sudden death, transient ischemic attack, stroke, atherosclerosis, venous thrombosis, deep vein thrombosis, thrombophlebitis, arterial embolism, coronary and cerebral arterial thrombosis, cerebral embolism, kidney embolisms, and pulmonary embolisms. The anticoagulant effect of compounds of the present invention is believed to be due to inhibition of factor Xa or thrombin.

The effectiveness of compounds of the present invention as inhibitors of factor Xa was determined using purified human factor Xa and synthetic substrate. The rate of factor Xa hydrolysis of chromogenic substrate S2222 (Kabi Pharmacia, Franklin, OH) was measured both in the absence and presence of compounds of the present invention. Hydrolysis of the substrate resulted in the release of pNA, which was monitored spectrophotometrically by measuring the increase in absorbance at 405 nm. A decrease in the rate of absorbance change at 405 nm in the presence of inhibitor is indicative of enzyme inhibition. The results of this assay are expressed as inhibitory constant, K_i .

Factor Xa determinations were made in 0.10 M sodium phosphate buffer, pH 7.5, containing 0.20 M NaCl, and 0.5 % PEG 8000. The Michaelis constant, K_m , for substrate hydrolysis was determined at 25°C using the method of Lineweaver and Burk. Values of K_i were determined by allowing 0.2-0.5 nM human factor Xa (Enzyme Research Laboratories, South Bend, IN) to react with the substrate (0.20 mM-1 mM) in the presence of inhibitor. Reactions were allowed to go for 30 minutes and the velocities (rate of absorbance change vs time) were measured in the time frame of 25-30 minutes. The following relationship was used to calculate K_i values:

$$(v_0 - v_s) / v_s = I / (K_i (1 + S / K_m))$$

where:

v_o is the velocity of the control in the absence of inhibitor;

v_s is the velocity in the presence of inhibitor;

I is the concentration of inhibitor;

K_i is the dissociation constant of the enzyme:inhibitor complex;

S is the concentration of substrate;

K_m is the Michaelis constant.

Using the methodology described above, a number of compounds of the present invention were found to exhibit a K_i of $\leq 10 \mu M$, thereby confirming the utility of the compounds of the present invention as effective Xa inhibitors.

The antithrombotic effect of compounds of the present invention can be demonstrated in a rabbit arterio-venous (AV) shunt thrombosis model. In this model, rabbits weighing 2-3 kg anesthetized with a mixture of xylazine (10 mg/kg i.m.) and ketamine (50 mg/kg i.m.) are used. A saline-filled AV shunt device is connected between the femoral arterial and the femoral venous cannulae. The AV shunt device consists of a piece of 6-cm tygon tubing which contains a piece of silk thread. Blood will flow from the femoral artery via the AV-shunt into the femoral vein. The exposure of flowing blood to a silk thread will induce the formation of a significant thrombus. After forty minutes, the shunt is disconnected and the silk thread covered with thrombus is weighed. Test agents or vehicle will be given (i.v., i.p., s.c., or orally) prior to the opening of the AV shunt. The percentage inhibition of thrombus formation is determined for each treatment group. The ID50 values (dose which produces 50% inhibition of thrombus formation) are estimated by linear regression.

The compounds of formula (I) may also be useful as inhibitors of serine proteases, notably human thrombin, plasma kallikrein and plasmin. Because of their inhibitory action, these compounds are indicated for use in the prevention or treatment of physiological reactions, blood coagulation and inflammation, catalyzed by the aforesaid class of enzymes. Specifically, the compounds have utility as drugs for the

treatment of diseases arising from elevated thrombin activity such as myocardial infarction, and as reagents used as anticoagulants in the processing of blood to plasma for diagnostic and other commercial purposes.

Some compounds of the present invention were shown to be direct acting inhibitors of the serine protease thrombin by their ability to inhibit the cleavage of small molecule substrates by thrombin in a purified system. *In vitro* inhibition constants were determined by the method described by Kettner et al. in *J. Biol. Chem.* **265**, 18289-18297 (1990), herein incorporated by reference. In these assays, thrombin-mediated hydrolysis of the chromogenic substrate S2238 (Helena Laboratories, Beaumont, TX) was monitored spectrophotometrically. Addition of an inhibitor to the assay mixture results in decreased absorbance and is indicative of thrombin inhibition. Human thrombin (Enzyme Research Laboratories, Inc., South Bend, IN) at a concentration of 0.2 nM in 0.10 M sodium phosphate buffer, pH 7.5, 0.20 M NaCl, and 0.5% PEG 6000, was incubated with various substrate concentrations ranging from 0.20 to 0.02 mM. After 25 to 30 minutes of incubation, thrombin activity was assayed by monitoring the rate of increase in absorbance at 405 nm which arises owing to substrate hydrolysis. Inhibition constants were derived from reciprocal plots of the reaction velocity as a function of substrate concentration using the standard method of Lineweaver and Burk. Using the methodology described above, some compounds of this invention were evaluated and found to exhibit a K_i of less than 10 μ M, thereby confirming the utility of the compounds of the present invention as effective Xa inhibitors.

The compounds of the present invention can be administered alone or in combination with one or more additional therapeutic agents. These include other anti-coagulant or coagulation inhibitory agents, anti-platelet or platelet inhibitory agents, thrombin inhibitors, or thrombolytic or fibrinolytic agents.

The compounds are administered to a mammal in a therapeutically effective amount. By "therapeutically

effective amount" it is meant an amount of a compound of Formula I that, when administered alone or in combination with an additional therapeutic agent to a mammal, is effective to prevent or ameliorate the thromboembolic disease condition or the progression of the disease.

By "administered in combination" or "combination therapy" it is meant that the compound of Formula I and one or more additional therapeutic agents are administered concurrently to the mammal being treated. When administered in combination each component may be administered at the same time or sequentially in any order at different points in time. Thus, each component may be administered separately but sufficiently closely in time so as to provide the desired therapeutic effect. Other anticoagulant agents (or coagulation inhibitory agents) that may be used in combination with the compounds of this invention include warfarin and heparin, as well as other factor Xa inhibitors such as those described in the publications identified above under Background of the Invention.

The term anti-platelet agents (or platelet inhibitory agents), as used herein, denotes agents that inhibit platelet function such as by inhibiting the aggregation, adhesion or granular secretion of platelets. Such agents include, but are not limited to, the various known non-steroidal anti-inflammatory drugs (NSAIDS) such as aspirin, ibuprofen, naproxen, sulindac, indomethacin, mefenamate, droxicam, diclofenac, sulfinpyrazone, and piroxicam, including pharmaceutically acceptable salts or prodrugs thereof. Of the NSAIDS, aspirin (acetylsalicyclic acid or ASA), and piroxicam are preferred. Other suitable anti-platelet agents include ticlopidine, including pharmaceutically acceptable salts or prodrugs thereof. Ticlopidine is also a preferred compound since it is known to be gentle on the gastro-intestinal tract in use. Still other suitable platelet inhibitory agents include IIb/IIIa antagonists, thromboxane-A₂-receptor antagonists and thromboxane-A₂-synthetase inhibitors, as well as pharmaceutically acceptable salts or prodrugs thereof.

The term thrombin inhibitors (or anti-thrombin agents), as used herein, denotes inhibitors of the serine protease thrombin. By inhibiting thrombin, various thrombin-mediated processes, such as thrombin-mediated platelet activation (that is, for example, the aggregation of platelets, and/or the granular secretion of plasminogen activator inhibitor-1 and/or serotonin) and/or fibrin formation are disrupted. A number of thrombin inhibitors are known to one of skill in the art and these inhibitors are contemplated to be used in combination with the present compounds. Such inhibitors include, but are not limited to, boroarginine derivatives, boro peptides, heparins, hirudin and argatroban, including pharmaceutically acceptable salts and prodrugs thereof. Boroarginine derivatives and boro peptides include N-acetyl and peptide derivatives of boronic acid, such as C-terminal α -aminoboronic acid derivatives of lysine, ornithine, arginine, homoarginine and corresponding isothiuronium analogs thereof. The term hirudin, as used herein, includes suitable derivatives or analogs of hirudin, referred to herein as hirulogs, such as disulfatohirudin. Boro peptide thrombin inhibitors include compounds described in Kettner et al., U.S. Patent No. 5,187,157 and European Patent Application Publication Number 293 881 A2, the disclosures of which are hereby incorporated herein by reference. Other suitable boroarginine derivatives and boro peptide thrombin inhibitors include those disclosed in PCT Application Publication Number 92/07869 and European Patent Application Publication Number 471,651 A2, the disclosures of which are hereby incorporated herein by reference.

The term thrombolytics (or fibrinolytic) agents (or thrombolytics or fibrinolytics), as used herein, denotes agents that lyse blood clots (thrombi). Such agents include tissue plasminogen activator, anistreplase, urokinase or streptokinase, including pharmaceutically acceptable salts or prodrugs thereof. The term anistreplase, as used herein, refers to anisoylated plasminogen streptokinase activator complex, as described, for example, in European Patent Application No. 028,489, the disclosure of which is hereby

incorporated herein by reference herein. The term urokinase, as used herein, is intended to denote both dual and single chain urokinase, the latter also being referred to herein as prourokinase.

Administration of the compounds of Formula I of the invention in combination with such additional therapeutic agent, may afford an efficacy advantage over the compounds and agents alone, and may do so while permitting the use of lower doses of each. A lower dosage minimizes the potential of side effects, thereby providing an increased margin of safety.

The compounds of the present invention are also useful as standard or reference compounds, for example as a quality standard or control, in tests or assays involving the inhibition of factor Xa. Such compounds may be provided in a commercial kit, for example, for use in pharmaceutical research involving factor Xa. For example, a compound of the present invention could be used as a reference in an assay to compare its known activity to a compound with an unknown activity. This would ensure the experimenter that the assay was being performed properly and provide a basis for comparison, especially if the test compound was a derivative of the reference compound. When developing new assays or protocols, compounds according to the present invention could be used to test their effectiveness.

The compounds of the present invention may also be used in diagnostic assays involving factor Xa. For example, the presence of factor Xa in an unknown sample could be determined by addition of chromogenic substrate S2222 to a series of solutions containing test sample and optionally one of the compounds of the present invention. If production of pNA is observed in the solutions containing test sample, but no compound of the present invention, then one would conclude factor Xa was present.

Dosage and Formulation

The compounds of this invention can be administered in such oral dosage forms as tablets, capsules (each of which includes sustained release or timed release formulations),

pills, powders, granules, elixirs, tinctures, suspensions, syrups, and emulsions. They may also be administered in intravenous (bolus or infusion), intraperitoneal, subcutaneous, or intramuscular form, all using dosage forms well known to those of ordinary skill in the pharmaceutical arts. They can be administered alone, but generally will be administered with a pharmaceutical carrier selected on the basis of the chosen route of administration and standard pharmaceutical practice.

The dosage regimen for the compounds of the present invention will, of course, vary depending upon known factors, such as the pharmacodynamic characteristics of the particular agent and its mode and route of administration; the species, age, sex, health, medical condition, and weight of the recipient; the nature and extent of the symptoms; the kind of concurrent treatment; the frequency of treatment; the route of administration, the renal and hepatic function of the patient, and the effect desired. A physician or veterinarian can determine and prescribe the effective amount of the drug required to prevent, counter, or arrest the progress of the thromboembolic disorder.

By way of general guidance, the daily oral dosage of each active ingredient, when used for the indicated effects, will range between about 0.001 to 1000 mg/kg of body weight, preferably between about 0.01 to 100 mg/kg of body weight per day, and most preferably between about 1.0 to 20 mg/kg/day. Intravenously, the most preferred doses will range from about 1 to about 10 mg/kg/minute during a constant rate infusion. Compounds of this invention may be administered in a single daily dose, or the total daily dosage may be administered in divided doses of two, three, or four times daily.

Compounds of this invention can be administered in intranasal form via topical use of suitable intranasal vehicles, or via transdermal routes, using transdermal skin patches. When administered in the form of a transdermal delivery system, the dosage administration will, of course, be continuous rather than intermittent throughout the dosage regimen.

The compounds are typically administered in admixture with suitable pharmaceutical diluents, excipients, or carriers (collectively referred to herein as pharmaceutical carriers) suitably selected with respect to the intended form of administration, that is, oral tablets, capsules, elixirs, syrups and the like, and consistent with conventional pharmaceutical practices.

For instance, for oral administration in the form of a tablet or capsule, the active drug component can be combined with an oral, non-toxic, pharmaceutically acceptable, inert carrier such as lactose, starch, sucrose, glucose, methyl cellulose, magnesium stearate, dicalcium phosphate, calcium sulfate, mannitol, sorbitol and the like; for oral administration in liquid form, the oral drug components can be combined with any oral, non-toxic, pharmaceutically acceptable inert carrier such as ethanol, glycerol, water, and the like. Moreover, when desired or necessary, suitable binders, lubricants, disintegrating agents, and coloring agents can also be incorporated into the mixture. Suitable binders include starch, gelatin, natural sugars such as glucose or beta-lactose, corn sweeteners, natural and synthetic gums such as acacia, tragacanth, or sodium alginate, carboxymethylcellulose, polyethylene glycol, waxes, and the like. Lubricants used in these dosage forms include sodium oleate, sodium stearate, magnesium stearate, sodium benzoate, sodium acetate, sodium chloride, and the like. Disintegrators include, without limitation, starch, methyl cellulose, agar, bentonite, xanthan gum, and the like.

The compounds of the present invention can also be administered in the form of liposome delivery systems, such as small unilamellar vesicles, large unilamellar vesicles, and multilamellar vesicles. Liposomes can be formed from a variety of phospholipids, such as cholesterol, stearylamine, or phosphatidylcholines.

Compounds of the present invention may also be coupled with soluble polymers as targetable drug carriers. Such polymers can include polyvinylpyrrolidone, pyran copolymer, polyhydroxypropylmethacrylamide-phenol,

polyhydroxyethylaspartamidephenol, or polyethyleneoxide-polylysine substituted with palmitoyl residues. Furthermore, the compounds of the present invention may be coupled to a class of biodegradable polymers useful in achieving controlled release of a drug, for example, polylactic acid, polyglycolic acid, copolymers of polylactic and polyglycolic acid, polyepsilon caprolactone, polyhydroxy butyric acid, polyorthoesters, polyacetals, polydihydropyrans, polycyanoacylates, and crosslinked or amphipathic block copolymers of hydrogels.

Dosage forms (pharmaceutical compositions) suitable for administration may contain from about 1 milligram to about 100 milligrams of active ingredient per dosage unit. In these pharmaceutical compositions the active ingredient will ordinarily be present in an amount of about 0.5-95% by weight based on the total weight of the composition.

Gelatin capsules may contain the active ingredient and powdered carriers, such as lactose, starch, cellulose derivatives, magnesium stearate, stearic acid, and the like. Similar diluents can be used to make compressed tablets. Both tablets and capsules can be manufactured as sustained release products to provide for continuous release of medication over a period of hours. Compressed tablets can be sugar coated or film coated to mask any unpleasant taste and protect the tablet from the atmosphere, or enteric coated for selective disintegration in the gastrointestinal tract.

Liquid dosage forms for oral administration can contain coloring and flavoring to increase patient acceptance.

In general, water, a suitable oil, saline, aqueous dextrose (glucose), and related sugar solutions and glycols such as propylene glycol or polyethylene glycols are suitable carriers for parenteral solutions. Solutions for parenteral administration preferably contain a water soluble salt of the active ingredient, suitable stabilizing agents, and if necessary, buffer substances. Antioxidizing agents such as sodium bisulfite, sodium sulfite, or ascorbic acid, either alone or combined, are suitable stabilizing agents. Also used are citric acid and its salts and sodium EDTA. In addition,

parenteral solutions can contain preservatives, such as benzalkonium chloride, methyl- or propyl-paraben, and chlorobutanol.

Suitable pharmaceutical carriers are described in Remington's Pharmaceutical Sciences, Mack Publishing Company, a standard reference text in this field.

Representative useful pharmaceutical dosage-forms for administration of the compounds of this invention can be illustrated as follows:

Capsules

A large number of unit capsules can be prepared by filling standard two-piece hard gelatin capsules each with 100 milligrams of powdered active ingredient, 150 milligrams of lactose, 50 milligrams of cellulose, and 6 milligrams magnesium stearate.

Soft Gelatin Capsules

A mixture of active ingredient in a digestable oil such as soybean oil, cottonseed oil or olive oil may be prepared and injected by means of a positive displacement pump into gelatin to form soft gelatin capsules containing 100 milligrams of the active ingredient. The capsules should be washed and dried.

Tablets

Tablets may be prepared by conventional procedures so that the dosage unit is 100 milligrams of active ingredient, 0.2 milligrams of colloidal silicon dioxide, 5 milligrams of magnesium stearate, 275 milligrams of microcrystalline cellulose, 11 milligrams of starch and 98.8 milligrams of lactose. Appropriate coatings may be applied to increase palatability or delay absorption.

Injectable

A parenteral composition suitable for administration by injection may be prepared by stirring 1.5% by weight of active ingredient in 10% by volume propylene glycol and water. The solution should be made isotonic with sodium chloride and sterilized.

Suspension

An aqueous suspension can be prepared for oral administration so that each 5 mL contain 100 mg of finely divided active ingredient, 200 mg of sodium carboxymethyl cellulose, 5 mg of sodium benzoate, 1.0 g of sorbitol solution, U.S.P., and 0.025 mL of vanillin.

Where the compounds of this invention are combined with other anticoagulant agents, for example, a daily dosage may be about 0.1 to 100 milligrams of the compound of Formula I and about 1 to 7.5 milligrams of the second anticoagulant, per kilogram of patient body weight. For a tablet dosage form, the compounds of this invention generally may be present in an amount of about 5 to 10 milligrams per dosage unit, and the second anti-coagulant in an amount of about 1 to 5 milligrams per dosage unit.

Where the compounds of Formula I are administered in combination with an anti-platelet agent, by way of general guidance, typically a daily dosage may be about 0.01 to 25 milligrams of the compound of Formula I and about 50 to 150 milligrams of the anti-platelet agent, preferably about 0.1 to 1 milligrams of the compound of Formula I and about 1 to 3 milligrams of antiplatelet agents, per kilogram of patient body weight.

Where the compounds of Formula I are administered in combination with thrombolytic agent, typically a daily dosage may be about 0.1 to 1 milligrams of the compound of Formula I, per kilogram of patient body weight and, in the case of the thrombolytic agents, the usual dosage of the thrombolytic agent when administered alone may be reduced by about 70-80% when administered with a compound of Formula I.

Where two or more of the foregoing second therapeutic agents are administered with the compound of Formula I, generally the amount of each component in a typical daily dosage and typical dosage form may be reduced relative to the usual dosage of the agent when administered alone, in view of the additive or synergistic effect of the therapeutic agents when administered in combination.

Particularly when provided as a single dosage unit, the potential exists for a chemical interaction between the combined active ingredients. For this reason, when the compound of Formula I and a second therapeutic agent are combined in a single dosage unit they are formulated such that although the active ingredients are combined in a single dosage unit, the physical contact between the active ingredients is minimized (that is, reduced). For example, one active ingredient may be enteric coated. By enteric coating one of the active ingredients, it is possible not only to minimize the contact between the combined active ingredients, but also, it is possible to control the release of one of these components in the gastrointestinal tract such that one of these components is not released in the stomach but rather is released in the intestines. One of the active ingredients may also be coated with a material which effects a sustained-release throughout the gastrointestinal tract and also serves to minimize physical contact between the combined active ingredients. Furthermore, the sustained-released component can be additionally enteric coated such that the release of this component occurs only in the intestine. Still another approach would involve the formulation of a combination product in which the one component is coated with a sustained and/or enteric release polymer, and the other component is also coated with a polymer such as a lowviscosity grade of hydroxypropyl methylcellulose (HPMC) or other appropriate materials as known in the art, in order to further separate the active components. The polymer coating serves to form an additional barrier to interaction with the other component.


These as well as other ways of minimizing contact between the components of combination products of the present invention, whether administered in a single dosage form or administered in separate forms but at the same time by the same manner, will be readily apparent to those skilled in the art, once armed with the present disclosure.

Obviously, numerous modifications and variations of the present invention are possible in light of the above teachings. It is therefore to be understood that within the

scope of the appended claims, the invention may be practiced otherwise than as specifically described herein.

WHAT IS CLAIMED AS NEW AND DESIRED TO BE SECURED BY
LETTER PATENT OF UNITED STATES IS:

1. A compound of formula I:



or a stereoisomer or pharmaceutically acceptable salt thereof,
wherein;

ring M contains, in addition to J, 0-2 N atoms;

J is O or S;

D is selected from CN, C(=NR⁸)NR⁷R⁹, NHC(=NR⁸)NR⁷R⁹,
NR⁸CH(=NR⁷), C(O)NR⁷R⁸, and (CR⁸R⁹)_tNR⁷R⁸, provided that D
is substituted meta or para to G on E;

E is selected from phenyl, pyridyl, pyrimidyl, pyrazinyl,
pyridazinyl, and piperidinyl substituted with 1 R;

alternatively, D-E-G together represent pyridyl substituted
with 1 R;

R is selected from H, halogen, (CH₂)_tOR³, C₁₋₄ alkyl, OCF₃, and
CF₃;

G is absent or is selected from NHCH₂, OCH₂, and SCH₂;

Z is selected from a C₁₋₄ alkylene, (CH₂)_rO(CH₂)_r,
(CH₂)_rNR³(CH₂)_r, (CH₂)_rC(O)(CH₂)_r, (CH₂)_rC(O)O(CH₂)_r,
(CH₂)_rOC(O)(CH₂)_r, (CH₂)_rC(O)NR³(CH₂)_r,
(CH₂)_rNR³C(O)(CH₂)_r, (CH₂)_rOC(O)O(CH₂)_r,
(CH₂)_rOC(O)NR³(CH₂)_r, (CH₂)_rNR³C(O)O(CH₂)_r,
(CH₂)_rNR³C(O)NR³(CH₂)_r, (CH₂)_rS(O)_p(CH₂)_r,
(CH₂)_rSO₂NR³(CH₂)_r, (CH₂)_rNR³SO₂(CH₂)_r, and

$(\text{CH}_2)_r\text{NR}^3\text{SO}_2\text{NR}^3(\text{CH}_2)_r$, provided that Z does not form a N-N, N-O, N-S, NCH_2N , NCH_2O , or NCH_2S bond with ring M or group A;

R^{1a} and R^{1b} are independently absent or selected from $-(\text{CH}_2)_r\text{R}^{1'}$, $\text{NCH}_2\text{R}^{1''}$, $\text{OCH}_2\text{R}^{1''}$, $\text{SCH}_2\text{R}^{1''}$, $\text{N}(\text{CH}_2)_2(\text{CH}_2)_t\text{R}^{1'}$, $\text{O}(\text{CH}_2)_2(\text{CH}_2)_t\text{R}^{1'}$, and $\text{S}(\text{CH}_2)_2(\text{CH}_2)_t\text{R}^{1'}$, or combined to form a 5-8 membered saturated, partially saturated or unsaturated ring substituted with 0-2 R^4 and which contains from 0-2 heteroatoms selected from the group consisting of N, O, and S;

$\text{R}^{1'}$ is selected from H, C_{1-3} alkyl, halo, $(\text{CF}_2)_r\text{CF}_3$, OR^2 , NR^2R^{2a} , $\text{C}(\text{O})\text{R}^{2c}$, $\text{OC}(\text{O})\text{R}^2$, $(\text{CF}_2)_r\text{CO}_2\text{R}^{2c}$, $\text{S}(\text{O})_p\text{R}^{2b}$, $\text{NR}^2(\text{CH}_2)_r\text{OR}^2$, $\text{NR}^2\text{C}(\text{O})\text{R}^{2b}$, $\text{NR}^2\text{C}(\text{O})\text{NHR}^{2b}$, $\text{NR}^2\text{C}(\text{O})_2\text{R}^{2a}$, $\text{OC}(\text{O})\text{NR}^{2b}$, $\text{C}(\text{O})\text{NR}^2\text{R}^{2a}$, $\text{SO}_2\text{NR}^2\text{R}^{2a}$, $\text{NR}^2\text{SO}_2\text{R}^{2b}$, C_{3-6} carbocyclic residue substituted with 0-2 R^4 , and 5-10 membered heterocyclic system containing from 1-4 heteroatoms selected from the group consisting of N, O, and S substituted with 0-2 R^4 ;

$\text{R}^{1''}$ is selected from H, $\text{C}(\text{O})\text{R}^{2b}$, $\text{C}(\text{O})\text{NR}^2\text{R}^{2a}$, $\text{S}(\text{O})\text{R}^{2b}$, $\text{S}(\text{O})_2\text{R}^{2b}$, and $\text{SO}_2\text{NR}^2\text{R}^{2a}$;

R^2 , at each occurrence, is selected from H, CF_3 , C_{1-6} alkyl, benzyl, C_{3-6} carbocyclic residue substituted with 0-2 R^{4b} , and 5-6 membered heterocyclic system containing from 1-4 heteroatoms selected from the group consisting of N, O, and S substituted with 0-2 R^{4b} ;

R^{2a} , at each occurrence, is selected from H, CF_3 , C_{1-6} alkyl, benzyl, C_{3-6} carbocyclic residue substituted with 0-2 R^{4b} , and 5-6 membered heterocyclic system containing from 1-4 heteroatoms selected from the group consisting of N, O, and S substituted with 0-2 R^{4b} ;

R^{2b} , at each occurrence, is selected from CF_3 , C_{1-4} alkoxy, C_{1-6} alkyl, benzyl, C_{3-6} carbocyclic residue substituted with

0-2 R^{4b} , and 5-6 membered heterocyclic system containing from 1-4 heteroatoms selected from the group consisting of N, O, and S substituted with 0-2 R^{4b} ;

R^{2c} , at each occurrence, is selected from CF_3 , OH, C_{1-4} alkoxy, C_{1-6} alkyl, benzyl, C_{3-6} carbocyclic residue substituted with 0-2 R^{4b} , and 5-6 membered heterocyclic system containing from 1-4 heteroatoms selected from the group consisting of N, O, and S substituted with 0-2 R^{4b} ;

alternatively, R^2 and R^{2a} combine to form a 5 or 6 membered saturated, partially saturated or unsaturated ring substituted with 0-2 R^{4b} which contains from 0-1 additional heteroatoms selected from the group consisting of N, O, and S;

R^3 , at each occurrence, is selected from H, C_{1-4} alkyl, and phenyl;

R^{3a} , at each occurrence, is selected from H, C_{1-4} alkyl, and phenyl;

A is selected from:

C_{3-10} carbocyclic residue substituted with 0-2 R^4 , and 5-10 membered heterocyclic system containing from 1-4 heteroatoms selected from the group consisting of N, O, and S substituted with 0-2 R^4 ;

B is selected from:

X-Y, NR^2R^{2a} , $C(=NR^2)NR^2R^{2a}$, $NR^2C(=NR^2)NR^2R^{2a}$, C_{3-10} carbocyclic residue substituted with 0-2 R^{4a} , and 5-10 membered heterocyclic system containing from 1-4 heteroatoms selected from the group consisting of N, O, and S substituted with 0-2 R^{4a} ;

X is selected from C_{1-4} alkylene, $-CR^2(CR^2R^{2b})(CH_2)_t-$, $-C(O)-$, $-C(=NR)-$, $-CR^2(NR^1R^2)-$, $-CR^2(OR^2)-$, $-CR^2(SR^2)-$, $-C(O)CR^2R^{2a}-$, $-CR^2R^{2a}C(O)-$, $-S(O)_p-$, $-S(O)_pCR^2R^{2a}-$,

-CR²R^{2a}S(O)_p-, -S(O)₂NR²-, -NR²S(O)₂-, -NR²S(O)₂CR²R^{2a}-,
 -CR²R^{2a}S(O)₂NR²-, -NR²S(O)₂NR²-, -C(O)NR²-, -NR²C(O)-,
 -C(O)NR²CR²R^{2a}-, -NR²C(O)CR²R^{2a}-, -CR²R^{2a}C(O)NR²-,
 -CR²R^{2a}NR²C(O)-, -NR²C(O)O-, -OC(O)NR²-, -NR²C(O)NR²-,
 -NR²-, -NR²CR²R^{2a}-, -CR²R^{2a}NR²-, O, -CR²R^{2a}O-, and
 -OCR²R^{2a}-;

Y is selected from:

(CH₂)_rNR²R^{2a}, provided that X-Y do not form a N-N, O-N, or S-N bond,

C₃₋₁₀ carbocyclic residue substituted with 0-2 R^{4a}, and
 5-10 membered heterocyclic system containing from 1-4 heteroatoms selected from the group consisting of N, O, and S substituted with 0-2 R^{4a};

R⁴, at each occurrence, is selected from =O, (CH₂)_rOR², halo, C₁₋₄ alkyl, -CN, NO₂, (CH₂)_rNR²R^{2a}, (CH₂)_rC(O)R^{2b}, NR²C(O)R^{2b}, C(O)NR²R^{2a}, NR²C(O)NR²R^{2a}, CH(=NR²)NR²R^{2a}, NHC(=NR²)NR²R^{2a}, SO₂NR²R^{2a}, NR²SO₂NR²R^{2a}, NR²SO₂-C₁₋₄ alkyl, NR²SO₂R⁵, S(O)_pR⁵, (CF₂)_rCF₃, NCH₂R^{1''}, OCH₂R^{1''}, SCH₂R^{1''}, N(CH₂)₂(CH₂)_tR^{1'}, O(CH₂)₂(CH₂)_tR^{1'}, and S(CH₂)₂(CH₂)_tR^{1'},

alternatively, one R⁴ is a 5-6 membered aromatic heterocycle containing from 1-4 heteroatoms selected from the group consisting of N, O, and S;

R^{4a}, at each occurrence, is selected from =O, (CH₂)_rOR², halo, C₁₋₄ alkyl, -CN, NO₂, (CH₂)_rNR²R^{2a}, (CH₂)_rC(O)R^{2b}, NR²C(O)R^{2b}, C(O)NR²R^{2a}, NR²C(O)NR²R^{2a}, CH(=NR²)NR²R^{2a}, NHC(=NR²)NR²R^{2a}, SO₂NR²R^{2a}, NR²SO₂NR²R^{2a}, NR²SO₂-C₁₋₄ alkyl, NR²SO₂R⁵, S(O)_pR⁵, and (CF₂)_rCF₃;

alternatively, one R^{4a} is a 5-6 membered aromatic heterocycle containing from 1-4 heteroatoms selected from the group consisting of N, O, and S substituted with 0-1 R⁵;

R^{4b}, at each occurrence, is selected from =O, (CH₂)_rOR³, halo, C₁₋₄ alkyl, -CN, NO₂, (CH₂)_rNR³R^{3a}, (CH₂)_rC(O)R³,

NR³C(O)R^{3a}, C(O)NR³R^{3a}, NR³C(O)NR³R^{3a}, CH(=NR³)NR³R^{3a}, NH³C(=NR³)NR³R^{3a}, SO₂NR³R^{3a}, NR³SO₂NR³R^{3a}, NR³SO₂-C₁₋₄ alkyl, NR³SO₂CF₃, NR³SO₂-phenyl, S(O)_pCF₃, S(O)_p-C₁₋₄ alkyl, S(O)_p-phenyl, and (CF₂)_rCF₃;

R⁵, at each occurrence, is selected from CF₃, C₁₋₆ alkyl, phenyl substituted with 0-2 R⁶, and benzyl substituted with 0-2 R⁶;

R⁶, at each occurrence, is selected from H, OH, (CH₂)_rOR², halo, C₁₋₄ alkyl, CN, NO₂, (CH₂)_rNR²R^{2a}, (CH₂)_rC(O)R^{2b}, NR²C(O)R^{2b}, NR²C(O)NR²R^{2a}, CH(=NH)NH₂, NHC(=NH)NH₂, SO₂NR²R^{2a}, NR²SO₂NR²R^{2a}, and NR²SO₂C₁₋₄ alkyl;

R⁷, at each occurrence, is selected from H, OH, C₁₋₆ alkyl, C₁₋₆ alkylcarbonyl, C₁₋₆ alkoxy, C₁₋₄ alkoxycarbonyl, (CH₂)_n-phenyl, C₆₋₁₀ aryloxy, C₆₋₁₀ aryloxycarbonyl, C₆₋₁₀ arylmethylcarbonyl, C₁₋₄ alkylcarbonyloxy C₁₋₄ alkoxycarbonyl, C₆₋₁₀ arylcarbonyloxy C₁₋₄ alkoxycarbonyl, C₁₋₆ alkylaminocarbonyl, phenylaminocarbonyl, and phenyl C₁₋₄ alkoxycarbonyl;

R⁸, at each occurrence, is selected from H, C₁₋₆ alkyl and (CH₂)_n-phenyl;

alternatively, R⁷ and R⁸ combine to form a 5 or 6 membered saturated, ring which contains from 0-1 additional heteroatoms selected from the group consisting of N, O, and S;

R⁹, at each occurrence, is selected from H, C₁₋₆ alkyl and (CH₂)_n-phenyl;


n, at each occurrence, is selected from 0, 1, 2, and 3;

m, at each occurrence, is selected from 0, 1, and 2;

p, at each occurrence, is selected from 0, 1, and 2;

r, at each occurrence, is selected from 0, 1, 2, and 3;
 s, at each occurrence, is selected from 0, 1, and 2; and,
 t, at each occurrence, is selected from 0 and 1;
 provided that D-E-G-(CH₂)_s- and -Z-A-B are not both
 benzamidines.

2. A compound according to Claim 1, wherein the compound
 is of formulae Ia-If:



wherein, groups D-E- and -Z-A-B are attached to adjacent atoms
 on the ring;

Z is selected from a CH₂O, OCH₂, CH₂NH, NHCH₂, C(O), CH₂C(O),
 C(O)CH₂, NHC(O), C(O)NH, CH₂S(O)₂, S(O)₂(CH₂), SO₂NH, and
 NHSO₂, provided that Z does not form a N-N, N-O, NCH₂N, or
 NCH₂O bond with ring M or group A;

A is selected from one of the following carbocyclic and
 heterocyclic systems which are substituted with 0-2 R⁴;
 phenyl, piperidiny, piperaziny, pyridyl,
 pyrimidyl, furanyl, morpholinyl, thiophenyl, pyrrolyl,
 pyrrolidinyl, oxazolyl, isoxazolyl, thiazolyl,
 isothiazolyl, pyrazolyl, imidazolyl, oxadiazolyl,
 thiadiazolyl, triazolyl, 1,2,3-oxadiazolyl,

1,2,4-oxadiazolyl, 1,2,5-oxadiazolyl, 1,3,4-oxadiazolyl, 1,2,3-thiadiazolyl, 1,2,4-thiadiazolyl, 1,2,5-thiadiazolyl, 1,3,4-thiadiazolyl, 1,2,3-triazolyl, 1,2,4-triazolyl, 1,2,5-triazolyl, 1,3,4-triazolyl, benzofuranyl, benzothiofuranyl, indolyl, benzimidazolyl, benzoxazolyl, benzthiazolyl, indazolyl, benzisoxazolyl, benzisothiazolyl, and isoindazolyl;

B is selected from: Y, X-Y, NR^2R^{2a} , $\text{C}(=\text{NR}^2)\text{NR}^2\text{R}^{2a}$, and $\text{NR}^2\text{C}(=\text{NR}^2)\text{NR}^2\text{R}^{2a}$;


X is selected from C_{1-4} alkylene, $-\text{C}(\text{O})-$, $-\text{C}(=\text{NR})-$, $-\text{CR}^2(\text{NR}^2\text{R}^{2a})-$, $-\text{C}(\text{O})\text{CR}^2\text{R}^{2a}-$, $-\text{CR}^2\text{R}^{2a}\text{C}(\text{O})-$, $-\text{C}(\text{O})\text{NR}^2-$, $-\text{NR}^2\text{C}(\text{O})-$, $-\text{C}(\text{O})\text{NR}^2\text{CR}^2\text{R}^{2a}-$, $-\text{NR}^2\text{C}(\text{O})\text{CR}^2\text{R}^{2a}-$, $-\text{CR}^2\text{R}^{2a}\text{C}(\text{O})\text{NR}^2-$, $-\text{CR}^2\text{R}^{2a}\text{NR}^2\text{C}(\text{O})-$, $-\text{NR}^2\text{C}(\text{O})\text{NR}^2-$, $-\text{NR}^2-$, $-\text{NR}^2\text{CR}^2\text{R}^{2a}-$, $-\text{CR}^2\text{R}^{2a}\text{NR}^2-$, O, $-\text{CR}^2\text{R}^{2a}\text{O}-$, and $-\text{OCR}^2\text{R}^{2a}-$;

Y is NR^2R^{2a} , provided that X-Y do not form a N-N or O-N bond;

alternatively, Y is selected from one of the following carbocyclic and heterocyclic systems which are substituted with 0-2 R^{4a} ;


cyclopropyl, cyclopentyl, cyclohexyl, phenyl, piperidiny, piperaziny, pyridyl, pyrimidyl, furanyl, morpholinyl, thiophenyl, pyrrolyl, pyrrolidinyl, oxazolyl, isoxazolyl, isoxazolinyl, thiazolyl, isothiazolyl, pyrazolyl, imidazolyl, oxadiazolyl, thiadiazolyl, triazolyl, 1,2,3-oxadiazolyl, 1,2,4-oxadiazolyl, 1,2,5-oxadiazolyl, 1,3,4-oxadiazolyl, 1,2,3-thiadiazolyl, 1,2,4-thiadiazolyl, 1,2,5-thiadiazolyl, 1,3,4-thiadiazolyl, 1,2,3-triazolyl, 1,2,4-triazolyl, 1,2,5-triazolyl, 1,3,4-triazolyl, benzofuranyl, benzothiofuranyl, indolyl, benzimidazolyl, benzoxazolyl, benzthiazolyl, indazolyl, benzisoxazolyl, benzisothiazolyl, and isoindazolyl;

alternatively, Y is selected from the following bicyclic heteroaryl ring systems:



K is selected from O, S, NH, and N.

3. A compound according to Claim 2, wherein the compound is of formulae Ib and Ic:



wherein;

J is O or S; and,

Z is selected from a C(O), CH₂C(O), C(O)CH₂, NHC(O), C(O)NH, C(O)N(CH₃), CH₂S(O)₂, S(O)₂(CH₂), SO₂NH, and NHSO₂, provided that Z does not form a N-N or NCH₂N bond with ring M or group A.

4. A compound according to Claim 3, wherein the compound is of formulae Ib and Ic:

E is phenyl substituted with R or 2-pyridyl substituted with R;

D is selected from NH_2 , C(O)NH_2 , C(=NH)NH_2 , CH_2NH_2 , CH_2NHCH_3 , $\text{CH(CH}_3\text{)NH}_2$, and $\text{C(CH}_3\text{)}_2\text{NH}_2$, provided that D is substituted meta or para to ring M on E; and,

R is selected from H, OCH_3 , Cl, and F.

5. A compound according to Claim 4, wherein the compound is of formulae Ib and Ic:

D-E is selected from 3-aminophenyl, 3-amidinophenyl, 3-aminomethylphenyl, 3-aminocarbonylphenyl, 3-(methylaminomethyl)phenyl, 3-(1-aminoethyl)phenyl, 3-(2-amino-2-propyl)phenyl, 4-chloro-3-aminophenyl, 4-chloro-3-amidinophenyl, 4-chloro-3-aminomethylphenyl, 4-chloro-3-(methylaminomethyl)phenyl, 4-fluoro-3-aminophenyl, 4-fluoro-3-amidinophenyl, 4-fluoro-3-aminomethylphenyl, 4-fluoro-3-(methylaminomethyl)phenyl, 6-aminopyrid-2-yl, 6-amidinopyrid-2-yl, 6-aminomethylpyrid-2-yl, 6-aminocarbonylpyrid-2-yl, 6-(methylaminomethyl)pyrid-2-yl, 6-(1-aminoethyl)pyrid-2-yl, and 6-(2-amino-2-propyl)pyrid-2-yl.

6. A compound according to Claim 3, wherein the compound is of formulae Ib and Ic:

Z is C(O)CH_2 and CONH , provided that Z does not form a N-N bond with group A;

A is selected from phenyl, pyridyl, and pyrimidyl, and is substituted with 0-2 R^4 ; and,

B is selected from X-Y, phenyl, pyrrolidino, morpholino, 1,2,3-triazolyl, and imidazolyl, and is substituted with 0-1 R^{4a};

R⁴, at each occurrence, is selected from OH, (CH₂)_rOR², halo, C₁₋₄ alkyl, (CH₂)_rNR²R^{2a}, and (CF₂)_rCF₃;

R^{4a} is selected from C₁₋₄ alkyl, CF₃, S(O)_pR⁵, SO₂NR²R^{2a}, and 1-CF₃-tetrazol-2-yl;

R⁵, at each occurrence, is selected from CF₃, C₁₋₆ alkyl, phenyl, and benzyl;

X is CH₂ or C(O); and,

Y is selected from pyrrolidino and morpholino.

7. A compound according to Claim 6, wherein the compound is of formulae Ib and Ic:

A is selected from the group: phenyl, 2-pyridyl, 3-pyridyl, 2-pyrimidyl, 2-Cl-phenyl, 3-Cl-phenyl, 2-F-phenyl, 3-F-phenyl, 2-methylphenyl, 2-aminophenyl, and 2-methoxyphenyl; and,

B is selected from the group: 2-CF₃-phenyl, 2-(aminosulfonyl)phenyl, 2-(methylaminosulfonyl)phenyl, 2-(dimethylaminosulfonyl)phenyl, 1-pyrrolidinocarbonyl, 2-(methylsulfonyl)phenyl, 4-morpholino, 2-(1'-CF₃-tetrazol-2-yl)phenyl, 4-morpholinocarbonyl, 2-methyl-1-imidazolyl, 5-methyl-1-imidazolyl, 2-methylsulfonyl-1-imidazolyl and, 5-methyl-1,2,3-triazolyl.

8. A compound according to Claim 3, wherein the compound is of formulae Ib and Ic:

E is phenyl substituted with R or 2-pyridyl substituted with R;

D is selected from NH_2 , $\text{C}(\text{O})\text{NH}_2$, $\text{C}(=\text{NH})\text{NH}_2$, CH_2NH_2 , CH_2NHCH_3 , $\text{CH}(\text{CH}_3)\text{NH}_2$, and $\text{C}(\text{CH}_3)_2\text{NH}_2$, provided that D is substituted meta or para to ring M on E; and,

R is selected from H, OCH_3 , Cl, and F;

Z is $\text{C}(\text{O})\text{CH}_2$ and CONH , provided that Z does not form a N-N bond with group A;

A is selected from phenyl, pyridyl, and pyrimidyl, and is substituted with 0-2 R^4 ; and,

B is selected from X-Y, phenyl, pyrrolidino, morpholino, 1,2,3-triazolyl, and imidazolyl, and is substituted with 0-1 R^{4a} ;

R^4 , at each occurrence, is selected from OH, $(\text{CH}_2)_r\text{OR}^2$, halo, C_{1-4} alkyl, $(\text{CH}_2)_r\text{NR}^2\text{R}^{2a}$, and $(\text{CF}_2)_r\text{CF}_3$;

R^{4a} is selected from C_{1-4} alkyl, CF_3 , $\text{S}(\text{O})_p\text{R}^5$, $\text{SO}_2\text{NR}^2\text{R}^{2a}$, and 1- CF_3 -tetrazol-2-yl;

R^5 , at each occurrence, is selected from CF_3 , C_{1-6} alkyl, phenyl, and benzyl;

X is CH_2 or $\text{C}(\text{O})$; and,

Y is selected from pyrrolidino and morpholino.

9. A compound according to Claim 8, wherein the compound is of formulae Ib and Ic:

D-E is selected from 3-aminophenyl, 3-amidinophenyl, 3-aminomethylphenyl, 3-aminocarbonylphenyl, 3-

(methylaminomethyl)phenyl, 3-(1-aminoethyl)phenyl, 3-(2-amino-2-propyl)phenyl, 4-chloro-3-aminophenyl, 4-chloro-3-amidinophenyl, 4-chloro-3-aminomethylphenyl, 4-chloro-3-(methylaminomethyl)phenyl, 4-fluoro-3-aminophenyl, 4-fluoro-3-amidinophenyl, 4-fluoro-3-aminomethylphenyl, 4-fluoro-3-(methylaminomethyl)phenyl, 6-aminopyrid-2-yl, 6-amidinopyrid-2-yl, 6-aminomethylpyrid-2-yl, 6-aminocarbonylpyrid-2-yl, 6-(methylaminomethyl)pyrid-2-yl, 6-(1-aminoethyl)pyrid-2-yl, 6-(2-amino-2-propyl)pyrid-2-yl;

A is selected from the group: phenyl, 2-pyridyl, 3-pyridyl, 2-pyrimidyl, 2-Cl-phenyl, 3-Cl-phenyl, 2-F-phenyl, 3-F-phenyl, 2-methylphenyl, 2-aminophenyl, and 2-methoxyphenyl; and,

B is selected from the group: 2-CF₃-phenyl, 2-(aminosulfonyl)phenyl, 2-(methylaminosulfonyl)phenyl, 2-(dimethylaminosulfonyl)phenyl, 1-pyrrolidinocarbonyl, 2-(methylsulfonyl)phenyl, 4-morpholino, 2-(1'-CF₃-tetrazol-2-yl)phenyl, 4-morpholinocarbonyl, 2-methyl-1-imidazolyl, 5-methyl-1-imidazolyl, 2-methylsulfonyl-1-imidazolyl and, 5-methyl-1,2,3-triazolyl.

10. A compound according to Claim 9, wherein the compound is of formula Ib₁.

11. A compound according to Claim 9, wherein the compound is of formula Ib₂.

12. A compound according to Claim 9, wherein the compound is of formula Ib₃.

13. A compound according to Claim 9, wherein the compound is of formula Ib₄.

14. A compound according to Claim 9, wherein the compound is of formula Ic₁.

15. A compound according to Claim 9, wherein the compound is of formula Ic₂.

16. A compound according to Claim 3, wherein the compound is of formulae Ib and Ic:

D is selected from C(=NR⁸)NR⁷R⁹, C(O)NR⁷R⁸, NR⁷R⁸, and CH₂NR⁷R⁸, provided that D is substituted meta or para to ring M on E;

E is phenyl substituted with R or pyridyl substituted with R;

R is selected from H, Cl, F, OR³, CH₃, CH₂CH₃, OCF₃, and CF₃;

Z is selected from C(O), CH₂C(O), C(O)CH₂, NHC(O), and C(O)NH, provided that Z does not form a N-N bond with ring M or group A;

R^{1a} and R^{1b} are independently absent or selected from -(CH₂)_r-R^{1'}, NCH₂R^{1''}, OCH₂R^{1''}, SCH₂R^{1''}, N(CH₂)₂(CH₂)_tR^{1'}, O(CH₂)₂(CH₂)_tR^{1'}, and S(CH₂)₂(CH₂)_tR^{1'}, or combined to form a 5-8 membered saturated, partially saturated or unsaturated ring substituted with 0-2 R⁴ and which contains from 0-2 heteroatoms selected from the group consisting of N, O, and S;

R^{1'}, at each occurrence, is selected from H, C₁₋₃ alkyl, halo, (CF₂)_rCF₃, OR², NR²R^{2a}, C(O)R^{2c}, (CF₂)_rCO₂R^{2c}, S(O)_pR^{2b},

$\text{NR}^2(\text{CH}_2)_r\text{OR}^2$, $\text{NR}^2\text{C}(\text{O})\text{R}^{2b}$, $\text{NR}^2\text{C}(\text{O})_2\text{R}^{2b}$, $\text{C}(\text{O})\text{NR}^2\text{R}^{2a}$,
 $\text{SO}_2\text{NR}^2\text{R}^{2a}$, and $\text{NR}^2\text{SO}_2\text{R}^{2b}$;

A is selected from one of the following carbocyclic and heterocyclic systems which are substituted with 0-2 R^4 ;
 phenyl, piperidinyl, piperazinyl, pyridyl,
 pyrimidyl, furanyl, morpholinyl, thiophenyl, pyrrolyl,
 pyrrolidinyl, oxazolyl, isoxazolyl, thiazolyl,
 isothiazolyl, pyrazolyl, and imidazolyl;

B is selected from: Y, X-Y, NR^2R^{2a} , $\text{C}(=\text{NR}^2)\text{NR}^2\text{R}^{2a}$, and
 $\text{NR}^2\text{C}(=\text{NR}^2)\text{NR}^2\text{R}^{2a}$;

X is selected from CH_2 , $-\text{CR}^2(\text{CR}^2\text{R}^{2b})(\text{CH}_2)_t-$, $-\text{C}(\text{O})-$, $-\text{C}(=\text{NR})-$,
 $-\text{CH}(\text{NR}^2\text{R}^{2a})-$, $-\text{C}(\text{O})\text{NR}^2-$, $-\text{NR}^2\text{C}(\text{O})-$, $-\text{NR}^2\text{C}(\text{O})\text{NR}^2-$, $-\text{NR}^2-$,
 and O;

Y is NR^2R^{2a} , provided that X-Y do not form a N-N or O-N bond;

alternatively, Y is selected from one of the following carbocyclic and heterocyclic systems which are substituted with 0-2 R^{4a} ;

phenyl, piperidinyl, piperazinyl, pyridyl,
 pyrimidyl, furanyl, morpholinyl, thiophenyl, pyrrolyl,
 pyrrolidinyl, oxazolyl, isoxazolyl, isoxazolinyll, thiazolyl,
 isothiazolyl, pyrazolyl, imidazolyl,
 oxadiazolyl, thiadiazolyl, triazolyl, 1,2,3-oxadiazolyl,
 1,2,4-oxadiazolyl, 1,2,5-oxadiazolyl, 1,3,4-oxadiazolyl,
 1,2,3-thiadiazolyl, 1,2,4-thiadiazolyl,
 1,2,5-thiadiazolyl, 1,3,4-thiadiazolyl, 1,2,3-triazolyl,
 1,2,4-triazolyl, 1,2,5-triazolyl, and 1,3,4-triazolyl;

R^4 , at each occurrence, is selected from =O, OH, Cl, F, C_{1-4} alkyl,
 $(\text{CH}_2)_r\text{NR}^2\text{R}^{2a}$, $(\text{CH}_2)_r\text{C}(\text{O})\text{R}^{2b}$, $\text{NR}^2\text{C}(\text{O})\text{R}^{2b}$, $\text{C}(\text{O})\text{NR}^2\text{R}^{2a}$,
 $\text{CH}(=\text{NH})\text{NH}_2$, $\text{NHC}(=\text{NH})\text{NH}_2$, $\text{SO}_2\text{NR}^2\text{R}^{2a}$, $\text{NR}^2\text{SO}_2\text{-C}_{1-4}$ alkyl,
 $\text{NR}^2\text{SO}_2\text{R}^5$, $\text{S}(\text{O})_p\text{R}^5$, and $(\text{CF}_2)_r\text{CF}_3$;

R^{4a}, at each occurrence, is selected from =O, OH, Cl, F, C₁₋₄ alkyl, (CH₂)_rNR²R^{2a}, (CH₂)_rC(O)R^{2b}, NR²C(O)R^{2b}, C(O)NR²R^{2a}, CH(=NH)NH₂, NHC(=NH)NH₂, SO₂NR²R^{2a}, NR²SO₂-C₁₋₄ alkyl, NR²SO₂R⁵, S(O)_pR⁵, (CF₂)_rCF₃, and 1-CF₃-tetrazol-2-yl;

R⁵, at each occurrence, is selected from CF₃, C₁₋₆ alkyl, phenyl substituted with 0-2 R⁶, and benzyl substituted with 0-2 R⁶;

R⁶, at each occurrence, is selected from H, =O, OH, OR², Cl, F, CH₃, CN, NO₂, (CH₂)_rNR²R^{2a}, (CH₂)_rC(O)R^{2b}, NR²C(O)R^{2b}, CH(=NH)NH₂, NHC(=NH)NH₂, and SO₂NR²R^{2a};

R⁷, at each occurrence, is selected from H, OH, C₁₋₆ alkyl, C₁₋₆ alkylcarbonyl, C₁₋₆ alkoxy, C₁₋₄ alkoxycarbonyl, benzyl, C₆₋₁₀ aryloxy, C₆₋₁₀ aryloxycarbonyl, C₆₋₁₀ arylmethylcarbonyl, C₁₋₄ alkylcarbonyloxy C₁₋₄ alkoxycarbonyl, C₆₋₁₀ arylcarbonyloxy C₁₋₄ alkoxycarbonyl, C₁₋₆ alkylaminocarbonyl, phenylaminocarbonyl, and phenyl C₁₋₄ alkoxycarbonyl;

R⁸, at each occurrence, is selected from H, C₁₋₆ alkyl and benzyl; and

alternatively, R⁷ and R⁸ combine to form a morpholino group; and,

R⁹, at each occurrence, is selected from H, C₁₋₆ alkyl and benzyl.

17. A compound according to Claim 16, wherein the compound is of formulae Ib and Ic:

E is phenyl substituted with R or 2-pyridyl substituted with R;

R is selected from H, Cl, F, OCH₃, CH₃, OCF₃, and CF₃;

Z is selected from a $C(O)CH_2$ and $C(O)NH$, provided that Z does not form a N-N bond with group A;

R^{1a} is selected from H, CH_3 , CH_2CH_3 , Cl, F, CF_3 , OCH_3 , NR^2R^{2a} , $S(O)_pR^{2b}$, $CH_2S(O)_pR^{2b}$, $CH_2NR^2S(O)_pR^{2b}$, $C(O)R^{2c}$, $CH_2C(O)R^{2c}$, $C(O)NR^2R^{2a}$, and $SO_2NR^2R^{2a}$;

R^{1b} is selected from H, CH_3 , CH_2CH_3 , Cl, F, CF_3 , OCH_3 , NR^2R^{2a} , $S(O)_pR^{2b}$, $CH_2S(O)_pR^{2b}$, $CH_2NR^2S(O)_pR^{2b}$, $C(O)R^{2c}$, $CH_2C(O)R^{2c}$, $C(O)NR^2R^{2a}$, and $SO_2NR^2R^{2a}$;

A is selected from one of the following carbocyclic and heterocyclic systems which are substituted with 0-2 R^4 ;
phenyl, pyridyl, pyrimidyl, furanyl, thiophenyl, pyrrolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, pyrazolyl, and imidazolyl;

B is selected from: Y and X-Y;

X is selected from CH_2 , $-CR^2(CR^2R^{2b})-$, $-C(O)-$, $-C(=NR)-$, $-CH(NR^2R^{2a})-$, $-C(O)NR^2-$, $-NR^2C(O)-$, $-NR^2C(O)NR^2-$, $-NR^2-$, and O;

Y is NR^2R^{2a} , provided that X-Y do not form a N-N or O-N bond;

alternatively, Y is selected from one of the following carbocyclic and heterocyclic systems which are substituted with 0-2 R^{4a} ;

phenyl, piperidinyl, piperazinyl, pyridyl, pyrimidyl, furanyl, morpholinyl, thiophenyl, pyrrolyl, pyrrolidinyl, oxazolyl, isoxazolyl, isoxazolinyl, thiazolyl, isothiazolyl, pyrazolyl, imidazolyl, oxadiazolyl, thiadiazolyl, triazolyl, 1,2,3-oxadiazolyl, 1,2,4-oxadiazolyl, 1,2,5-oxadiazolyl, 1,3,4-oxadiazolyl, 1,2,3-thiadiazolyl, 1,2,4-thiadiazolyl, 1,2,5-thiadiazolyl, 1,3,4-thiadiazolyl, 1,2,3-triazolyl, 1,2,4-triazolyl, 1,2,5-triazolyl, and 1,3,4-triazolyl;

R², at each occurrence, is selected from H, CF₃, CH₃, benzyl, and phenyl;

R^{2a}, at each occurrence, is selected from H, CF₃, CH₃, benzyl, and phenyl;

R^{2b}, at each occurrence, is selected from CF₃, OCH₃, CH₃, benzyl, and phenyl;

R^{2c}, at each occurrence, is selected from CF₃, OH, OCH₃, CH₃, benzyl, and phenyl;

alternatively, R² and R^{2a} combine to form a 5 or 6 membered saturated, partially unsaturated, or unsaturated ring which contains from 0-1 additional heteroatoms selected from the group consisting of N, O, and S;

R³, at each occurrence, is selected from H, CH₃, CH₂CH₃, and phenyl;

R^{3a}, at each occurrence, is selected from H, CH₃, CH₂CH₃, and phenyl;

R⁴, at each occurrence, is selected from OH, Cl, F, CH₃, CH₂CH₃, NR²R^{2a}, CH₂NR²R^{2a}, C(O)R^{2b}, NR²C(O)R^{2b}, C(O)NR²R^{2a}, and CF₃;

R^{4a}, at each occurrence, is selected from OH, Cl, F, CH₃, CH₂CH₃, NR²R^{2a}, CH₂NR²R^{2a}, C(O)R^{2b}, C(O)NR²R^{2a}, SO₂NR²R^{2a}, S(O)_pR⁵, CF₃, and 1-CF₃-tetrazol-2-yl;

R⁵, at each occurrence, is selected from CF₃, C₁₋₆ alkyl, phenyl substituted with 0-2 R⁶, and benzyl substituted with 1 R⁶;

R⁶, at each occurrence, is selected from H, OH, OCH₃, Cl, F, CH₃, CN, NO₂, NR²R^{2a}, CH₂NR²R^{2a}, and SO₂NR²R^{2a};

R⁷, at each occurrence, is selected from H, OH, C₁₋₃ alkyl, C₁₋₃ alkylcarbonyl, C₁₋₃ alkoxy, C₁₋₄ alkoxycarbonyl, benzyl, phenoxy, phenoxycarbonyl, benzylcarbonyl, C₁₋₄ alkylcarbonyloxy C₁₋₄ alkoxycarbonyl, phenylcarbonyloxy C₁₋₄ alkoxycarbonyl, C₁₋₆ alkylaminocarbonyl, phenylaminocarbonyl, and phenyl C₁₋₄ alkoxycarbonyl;

R⁸, at each occurrence, is selected from H, CH₃, and benzyl; and,

alternatively, R⁷ and R⁸ combine to form a morpholino group;

R⁹, at each occurrence, is selected from H, CH₃, and benzyl.

18. A compound according to Claim 17, wherein the compound is of formulae Ib and Ic:

R^{1a} is absent or is selected from H, CH₃, CH₂CH₃, Cl, F, CF₃, OCH₃, NR²R^{2a}, S(O)_pR^{2b}, C(O)NR²R^{2a}, CH₂S(O)_pR^{2b}, CH₂NR²S(O)_pR^{2b}, C(O)R^{2c}, CH₂C(O)R^{2c}, and SO₂NR²R^{2a};

R^{1b} is absent or is selected from H, CH₃, CH₂CH₃, Cl, F, CF₃, OCH₃, NR²R^{2a}, S(O)_pR^{2b}, C(O)NR²R^{2a}, CH₂S(O)_pR^{2b}, CH₂NR²S(O)_pR^{2b}, C(O)R^{2b}, CH₂C(O)R^{2b}, and SO₂NR²R^{2a};

A is selected from one of the following carbocyclic and heterocyclic systems which are substituted with 0-2 R⁴; phenyl, pyridyl, and pyrimidyl;

B is selected from: Y and X-Y;

X is selected from -C(O)- and O;

Y is NR²R^{2a}, provided that X-Y do not form a O-N bond;

alternatively, Y is selected from one of the following carbocyclic and heterocyclic systems which are substituted with 0-2 R^{4a};

phenyl, piperazinyl, pyridyl, pyrimidyl, morpholinyl, pyrrolidinyl, imidazolyl, and 1,2,3-triazolyl;

R², at each occurrence, is selected from H, CF₃, CH₃, benzyl, and phenyl;

R^{2a}, at each occurrence, is selected from H, CF₃, CH₃, benzyl, and phenyl;

R^{2b}, at each occurrence, is selected from CF₃, OCH₃, CH₃, benzyl, and phenyl;

R^{2c}, at each occurrence, is selected from CF₃, OH, OCH₃, CH₃, benzyl, and phenyl;

alternatively, R² and R^{2a} combine to form a ring system selected from pyrrolidinyl, piperazinyl and morpholino;

R⁴, at each occurrence, is selected from Cl, F, CH₃, NR²R^{2a}, and CF₃;

R^{4a}, at each occurrence, is selected from Cl, F, CH₃, SO₂NR²R^{2a}, S(O)_pR⁵, and CF₃; and,

R⁵, at each occurrence, is selected from CF₃ and CH₃.

19. A compound according to Claim 1, wherein the compound is selected from the group:

3-(3-amidinophenyl)-4-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-5-(hydroxymethyl)isoxazole;

- 3-(3-amidinophenyl)-4-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]isoxazole;
- 3-(3-amidinophenyl)-4-[(2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]isoxazole;
- 3-(3-amidinophenyl)-4-[5-(2-aminosulfonyl)phenylpyrid-2-yl)aminocarbonyl]-5-(methoxymethyl)isoxazole;
- 3-(3-amidinophenyl)-4-[(2'-trifluoromethyl-[1,1']-biphen-4-yl)aminocarbonyl]isoxazole;
- 3-(3-amidinophenyl)-4-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-5-(trifluoromethyl)isoxazole;
- 2-acetylamino-4-(3-amidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole;
- 2-amino-4-(3-amidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole;
- 2-methyl-4-(3-amidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole;
- 5-(3-amidinophenyl)-4-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]oxazole;
- 3-(3-amidinophenyl)-4-[(2'-t-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]isoxazole;
- 3-(3-amidinophenyl)-4-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-5-(methoxymethyl)-isoxazole;
- 3-(3-amidinophenyl)-4-[(2'-t-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]isoxazole;
- 3-(3-amidinophenyl)-4-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-5-(methoxymethyl)isoxazole;

2-methyl-4-(3-amidinophenyl)-5-[(2'-trifluoromethyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole;

2-phenyl-4-(3-amidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole;

3-(3-amidinophenyl)-4-[(3-fluoro-2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]isoxazole;

3-(3-amidinophenyl)-4-[(2'-trifluoromethylthio-[1,1']-biphen-4-yl)aminocarbonyl]isoxazole;

3-(3-amidinophenyl)-5-amino-4-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]isoxazole;

2-(phenylamino)-4-(3-amidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole;

2-(benzylamino)-4-(3-amidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole;

2-(methylamino)-4-(3-amidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole;

2-(methylamino)-4-(3-carboxamidophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole;

2-methyl-4-(3-amidinophenyl)-5-[[5-(2'-aminosulfonylphenyl-1-yl)pyridin-2-yl]aminocarbonyl]thiazole;

2-methyl-4-(3-(carboxamido)phenyl)-5-[[5-(2'-aminosulfonylphenyl-1-yl)pyridin-2-yl]aminocarbonyl]thiazole;

2-(3-pyridyl)-4-(3-amidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole;

2-(3-pyridyl)-4-(3-carboxamidophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole;

2-chloro-4-(3-amidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole;

2-chloro-4-(3-carboxamidophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole;

2-chloro-4-(3-amidinophenyl)-5-[[5-(2'-aminosulfonylphenyl-1-yl)pyridin-2-yl]aminocarbonyl]thiazole;

2-chloro-4-(3-(carboxamido)phenyl)-5-[[5-(2'-aminosulfonylphenyl-1-yl)pyridin-2-yl]aminocarbonyl]thiazole;

2-hydroxy-4-(3-amidinophenyl)-5-[[5-(2'-aminosulfonylphenyl-1-yl)pyridin-2-yl]aminocarbonyl]thiazole;

2-chloro-4-(3-aminophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole;

2-amino-4-[(3-amino-4-chloro)phenyl]-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole;

2-chloro-4-[(3-amino-4-chloro)phenyl]-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole; and,

2-amino-4-[(3-aminomethyl)phenyl]-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]thiazole;

and a pharmaceutically acceptable salt thereof.

20. A pharmaceutical composition, comprising: a pharmaceutically acceptable carrier and a therapeutically effective amount of a compound according to Claim 1 or a pharmaceutically acceptable salt thereof.

21. A method for treating or preventing a thromboembolic disorder, comprising: administering to a patient in need thereof a therapeutically effective amount of a compound according to Claim 1 or a pharmaceutically acceptable salt thereof.

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6 : C07D 261/18, A61K 31/42, C07D 275/02, A61K 31/425, C07D 417/12, A61K 31/41, C07D 413/12, A61K 31/44, C07D 417/04		A3	(11) International Publication Number: WO 98/28282 (43) International Publication Date: 2 July 1998 (02.07.98)
(21) International Application Number: PCT/US97/23470 (22) International Filing Date: 18 December 1997 (18.12.97) (30) Priority Data: 08/771,814 23 December 1996 (23.12.96) US 08/879,763 20 June 1997 (20.06.97) US (71) Applicant: THE DU PONT MERCK PHARMACEUTICAL COMPANY [US/US]; 1007 Market Street, Wilmington, DE 19898 (US). (72) Inventors: PRUITT, James, Russell; 237 Skycrest Drive, Landenberg, PA 19350 (US). FEVIG, John, Matthew; 987 Church Road, Lincoln University, PA 19352 (US). QUAN, Mimi, Lifen; 113 Venus Drive, Newark, DE 19711 (US). PINTO, Donald, Joseph, Phillip; 39 Whitson Road, Newark, DE 19702 (US). (74) Agent: VANCE, David, H.; The Du Pont Merck Pharmaceu- tical Company, Legal Patent Records Center, 1007 Market Street, Wilmington, DE 19898 (US).		(81) Designated States: AM, AU, AZ, BR, BY, CA, CN, CZ, EE, HU, IL, JP, KG, KR, KZ, LT, LV, MD, MX, NO, NZ, PL, RO, RU, SG, SI, SK, TJ, TM, UA, VN, Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i> (88) Date of publication of the international search report: 17 September 1998 (17.09.98)	

(54) Title: OXYGEN OR SULFUR CONTAINING 5-MEMBERED HETEROAROMATICS AS FACTOR Xa INHIBITORS

(57) Abstract

The present application describes oxygen and sulfur containing heteroaromatics and derivatives thereof of formula (I), or pharmaceutically acceptable salt or prodrug forms thereof, wherein J is O or S and D may be C(=NH)NH₂, which are useful as inhibitors of factor Xa.



FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 97/23470

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07D261/18 A61K31/42 C07D275/02 A61K31/425 C07D417/12
A61K31/41 C07D413/12 A61K31/44 C07D417/04

According to International Patent Classification(IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,A	US 5 668 159 A (CONFALONE PASQUALE NICHOLAS ET AL) 16 September 1997 see claim 1 ---	1
A	WO 96 38426 A (DU PONT MERCK PHARMA) 5 December 1996 cited in the application see claim 1 ---	1
A	WO 95 18111 A (DU PONT MERCK PHARMA) 6 July 1995 cited in the application see claim 1 ---	1
A	US 5 317 103 A (BAKER RAYMOND ET AL) 31 May 1994 cited in the application see claim 1 ---	1
-/--		

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

28 May 1998

Date of mailing of the international search report

20.07.98

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Gettins, M

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 97/23470

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>EP 0 513 387 A (OTSUKA PHARMA CO LTD) 19 November 1992 see claim 1</p> <p>-----</p>	1

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US 97/23470

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Although claim 21 is directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound.
2. ☒ Claims Nos.: 1-18,20,21 (all partially)
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International Application No. PCT/US 97/23470

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Claims Nos.: 1-18,20,21 (all partially)

The scope of the claims and the thousands of examples are such that it is impossible to carry out a complete search on structural and economic grounds. It is noted that in all of the examples there appears to be a common structural element which appears to represent an essential technical (structural feature) It appears that G is always a bond, S is always zero, Z is always -C(O)-NH- , R1b is always hydrogen and the -E-D and -Z-A-B are attached to adjacent carbon atoms of the M ring. The search has been listed to these particular embodiments.

INTERNATIONAL SEARCH REPORT

Information on patent family members

Inter Jnal Application No

PCT/US 97/23470

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 5668159 A	16-09-1997	NONE	
WO 9638426 A	05-12-1996	AU 6024396 A EP 0832076 A	18-12-1996 01-04-1998
WO 9518111 A	06-07-1995	US 5563158 A AU 1400095 A US 5691329 A	08-10-1996 17-07-1995 25-11-1997
US 5317103 A	31-05-1994	NONE	
EP 0513387 A	19-11-1992	AU 656930 B AU 8936791 A CA 2074933 A WO 9209586 A JP 10101562 A JP 5051318 A US 5643932 A US 5677319 A	23-02-1995 25-06-1992 31-05-1992 11-06-1992 21-04-1998 02-03-1993 01-07-1997 14-10-1997

THIS PAGE BLANK (USPTO)